

Research Article

Some Geometric Inequalities in a New Banach Sequence Space

M. Mursaleen, Rifat Çolak, and Mikail Et

Received 11 July 2007; Accepted 18 November 2007

Recommended by Peter Yu Hin Pang

The difference sequence space $m(\phi, p, \Delta^{(r)})$, which is a generalization of the space $m(\phi)$ introduced and studied by Sargent (1960), was defined by Çolak and Et (2005). In this paper we establish some geometric inequalities for this space.

Copyright © 2007 M. Mursaleen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction and preliminaries

Let \mathcal{C} denote the space whose elements are finite sets of distinct positive integers. Given an element $\sigma \in \mathcal{C}$, we write $c(\sigma)$ for the sequence $(c_n(\sigma))$ such that $c_n(\sigma) = 1$ for $n \in \sigma$, and $c_n(\sigma) = 0$, otherwise. Further

$$\mathcal{C}_s = \left\{ \sigma \in \mathcal{C} : \sum_{n=1}^{\infty} c_n(\sigma) \leq s \right\}, \quad (1.1)$$

that is, \mathcal{C}_s is the set of those σ whose support has cardinality at most s , where s is a natural number.

Let w be the set of all real sequences and

$$\Phi = \left\{ \phi = (\phi_n) \in w : \phi_1 > 0, \nabla \phi_k \geq 0, \nabla \left(\frac{\phi_k}{k} \right) \leq 0 (k = 1, 2, \dots) \right\}, \quad (1.2)$$

where $\nabla \phi_k = \phi_k - \phi_{k-1}$. For $\phi \in \Phi$, Sargent [1] introduced the following sequence space:

$$m(\phi) = \left\{ x = (x_n) \in w : \sup_{s \geq 1} \sup_{\sigma \in \mathcal{C}_s} \left(\frac{1}{\phi_s} \sum_{n \in \sigma} |x_n| \right) < \infty \right\}. \quad (1.3)$$

In [2], the space $m(\phi)$ has been considered for matrix transformations and in [3] some of its geometric properties have been considered. Tripathy and Sen [4] extended $m(\phi)$ to $m(\phi, p)$, $1 \leq p < \infty$. Recently, Çolak and Et [5] defined the space $m(\phi, p, \Delta^{(r)})$ by using the idea of difference sequences (see [6–8]).

Let r be a positive integer throughout. The operators $\Delta^{(r)}, \Sigma^{(r)} : w \rightarrow w$ are defined by

$$\begin{aligned} (\Delta^{(1)}x)_k &= (\Delta x)_k = x_k - x_{k+1}, \\ (\Sigma^{(1)}x)_k &= (\Sigma x)_k = \sum_{j=k}^{\infty} x_j \quad (k = 1, 2, \dots), \\ \Delta^{(r)} &= \Delta^{(1)} \circ \Delta^{(r-1)}, \quad \Sigma^{(r)} = \Sigma^{(1)} \circ \Sigma^{(r-1)}, \quad (r \geq 2), \\ \Sigma^{(r)} \circ \Delta^{(r)} &= \Delta^{(r)} \circ \Sigma^{(r)} = id, \quad \text{the identity on } w. \end{aligned} \tag{1.4}$$

For $0 \leq p < \infty$, the space $m(\phi, p, \Delta^{(r)})$ is defined as follows:

$$m(\phi, p, \Delta^{(r)}) = \left\{ x \in w : \sup_{s \geq 1, \sigma \in \mathcal{C}_s} \left(\frac{1}{\phi_s} \sum_{n \in \sigma} |\Delta^{(r)}x_n|^p \right) < \infty \right\}, \tag{1.5}$$

which is a Banach space ($1 \leq p < \infty$) with the norm

$$\|x\|_{m(\phi, p, \Delta^{(r)})} = \sum_{i=1}^r |x_i| + \sup_{s \geq 1, \sigma \in \mathcal{C}_s} \frac{1}{\phi_s} \left(\sum_{n \in \sigma} |\Delta^{(r)}x_n|^p \right)^{1/p}, \tag{1.6}$$

and a complete p -normed space ($0 < p < 1$) with the p -norm

$$\|x\|_{m_p(\phi, \Delta^{(r)})} = \sum_{i=1}^r |x_i|^p + \sup_{s \geq 1, \sigma \in \mathcal{C}_s} \frac{1}{\phi_s} \sum_{n \in \sigma} |\Delta^{(r)}x_n|^p. \tag{1.7}$$

In this paper, we will consider the case $1 < p < \infty$ to study some geometric properties of $m(\phi, p, \Delta^{(r)})$. We will examine the Banach-Saks property of type p , strict convexity and uniform convexity. The space $m(\phi, p)$, $1 \leq p < \infty$ was defined by Tripathy and Sen [4] which is in fact $m(\phi, p, \Delta)$ with Δ replaced by id .

Let $1 < p < \infty$. A Banach space X is said to have the *Banach-Saks property of type p or property $(BS)_p$* if every weakly null-sequence (x_k) has a subsequence (x_{k_i}) such that for some $C > 0$, the inequality

$$\left\| \sum_{i=0}^n x_{k_i} \right\|_X \leq c(n+1)^{1/p}, \quad n = 1, 2, 3, \dots, \tag{1.8}$$

holds.

The property $(BS)_p$ for a Cesàro sequence space was considered in [9].

We find uniform convexity and strict convexity of our space through the Gurarii's modulus of convexity (see [10, 11]).

For a normed linear space X , the modulus of convexity defined by

$$\beta_X(\varepsilon) = \inf \left\{ 1 - \inf_{0 \leq \alpha \leq 1} \|\alpha x + (1 - \alpha)y\| : x, y \in S(X), \|x - y\| = \varepsilon \right\}, \tag{1.9}$$

is called the Gurarii's modulus of convexity, where $S(X)$ denotes the unit sphere in X and $0 < \varepsilon \leq 2$. If $0 < \beta_X(\varepsilon) < 1$, then X is uniformly convex and if $\beta_X(\varepsilon) \leq 1$, then X is strictly convex.

2. Main results

THEOREM 2.1. *The space $m(\phi, p, \Delta^{(r)})$ has the Banach-Saks property of type p .*

Proof. We will prove the case $r = 1$ and the general case can be followed on the same lines. \square

Let (ε_n) be a sequence of positive numbers for which $\sum_{n=1}^{\infty} \varepsilon_n \leq 1/2$. Let (x_n) be a weakly null sequence in $B(m(\phi, p, \Delta))$, the unit ball in $m(\phi, p, \Delta)$. Set $x_0 = 0$ and $z_1 = x_{n_1} = \Delta x_1$. Then there exists $s_1 \in \mathbb{N}$ such that

$$\left\| \sum_{i \in \tau_1} z_1(i) e_i \right\|_{m(\phi, p, \Delta)} < \varepsilon_1, \quad (2.1)$$

where τ_1 consists of the elements of σ which exceed s_1 . Since $x_n \xrightarrow{w} 0 \Rightarrow x_n \rightarrow 0$ coordinate-wise, there is $n_2 \in \mathbb{N}$ such that

$$\left\| \sum_{i=1}^{s_1} x_n(i) e_i \right\|_{m(\phi, p, \Delta)} < \varepsilon_1, \quad \text{when } n \geq n_2. \quad (2.2)$$

Set $z_2 = x_{n_2} = \Delta x_2$. Then there exists $s_2 > s_1$ such that

$$\left\| \sum_{i \in \tau_2} z_2(i) e_i \right\|_{m(\phi, p, \Delta)} < \varepsilon_2, \quad (2.3)$$

where τ_2 consists of the elements of σ which exceed s_2 . Again using the fact $x_n \rightarrow 0$ coordinate-wise, there exists $n_3 > n_2$ such that

$$\left\| \sum_{i=1}^{s_2} x_n(i) e_i \right\|_{m(\phi, p, \Delta)} < \varepsilon_2, \quad \text{when } n \geq n_3. \quad (2.4)$$

Continuing this process, we can find two increasing sequences (s_j) and (n_j) such that

$$\begin{aligned} \left\| \sum_{i=1}^{s_j} x_n(i) e_i \right\|_{m(\phi, p, \Delta)} &< \varepsilon_j, \quad \text{when } n \geq n_{j+1}, \\ \left\| \sum_{i \in \tau_j} z_j(i) e_i \right\|_{m(\phi, p, \Delta)} &< \varepsilon_j, \end{aligned} \quad (2.5)$$

where $z_j = x_{n_j} = \Delta x_j$ and τ_j consists of the elements of σ which exceed s_j . Note that $z_j(i)$ is a term in the sequence with fixed j and running i .

Since $\varepsilon_{j-1} + \varepsilon_j < 1$, we have

$$\left(\frac{1}{\phi_s} \sum_{n \in \sigma} |z_j(n)| \right) \leq (\varepsilon_{j-1} + \varepsilon_j) < 1, \tag{2.6}$$

for all $j \in \mathbb{N}$ and $s \geq 1$. Hence

$$\begin{aligned} \left\| \sum_{j=1}^n z_j \right\|_{m(\phi, p, \Delta)} &= \left\| \sum_{j=1}^n \left(\sum_{i=1}^{s_{j-1}} z_j(i) e_i + \sum_{i=s_{j-1}+1}^{s_j} z_j(i) e_i + \sum_{i \in \tau_j} z_j(i) e_i \right) \right\|_{m(\phi, p, \Delta)} \\ &\leq \left\| \sum_{j=1}^n \left(\sum_{i=1}^{s_{j-1}} z_j(i) e_i \right) \right\|_{m(\phi, p, \Delta)} + \left\| \sum_{j=1}^n \left(\sum_{i=s_{j-1}+1}^{s_j} z_j(i) e_i \right) \right\|_{m(\phi, p, \Delta)} \\ &\quad + \left\| \sum_{j=1}^n \left(\sum_{i \in \tau_j} z_j(i) e_i \right) \right\|_{m(\phi, p, \Delta)} \\ &\leq \sum_{j=1}^n \left\| \left(\sum_{i=s_{j-1}+1}^{s_j} z_j(i) e_i \right) \right\|_{m(\phi, p, \Delta)} + 2 \sum_{j=1}^n \varepsilon_j, \\ \sum_{j=1}^n \left\| \sum_{i=s_{j-1}+1}^{s_j} z_j(i) e_i \right\|_{m(\phi, p, \Delta)}^p &= \sum_{j=1}^n \sup_{s \geq 1} \sup_{\tau_{j-1} \in \mathcal{C}_s} \left(\frac{1}{\phi_s} \sum_{i \in \tau_{j-1}} |z_j(i)|^p \right) \\ &\leq \sum_{j=1}^n \sup_{s \geq 1} \sup_{\sigma \in \mathcal{C}_s} \left(\frac{1}{\phi_s} \sum_{i \in \sigma} |z_j(i)|^p \right) \leq n. \end{aligned} \tag{2.7}$$

Therefore by (2.7)

$$\left\| \sum_{j=1}^n z_j \right\|_{m(\phi, p, \Delta)} \leq n^{1/p} + 1 \leq 2n^{1/p} \tag{2.8}$$

since $\sum_{j=1}^n \varepsilon_j \leq 1/2$.

Hence $m(\phi, p, \Delta)$ has the Banach-Saks property of type p .

Remark 2.2. The above result can also be extended to the case when $r \neq 1$ and so the proof should also work for a more general case with Δ replaced by a matrix operator (transformation).

THEOREM 2.3. *The Gurarii’s modulus of convexity for the space $X = m(\phi, p, \Delta)$ is*

$$\beta_X(\varepsilon) \leq 1 - \left[1 - \left(\frac{\varepsilon}{2} \right)^p \right]^{1/p}, \tag{2.9}$$

where $0 < \varepsilon \leq 2$.

Proof. Let $x \in m(\phi, p, \Delta)$. Then

$$\|x\|_{m(\phi, p, \Delta)} = \|\Delta x\|_{m(\phi, p)} = |x_1| + \sup_{s \geq 1, \sigma \in \mathcal{C}_s} \frac{1}{\phi_s} \left[\sum_{n \in \sigma} |\Delta x_n|^p \right]^{1/p}. \quad (2.10)$$

Let $0 < \varepsilon \leq 2$ and consider the sequences

$$\begin{aligned} u = (u_n) &= \left(\left(\sum \left(1 - \left(\frac{\varepsilon}{2} \right)^p \right) \right)^{1/p}, \sum \left(\frac{\varepsilon}{2} \right), 0, 0, \dots \right), \\ v = (v_n) &= \left(\left(\sum \left(1 - \left(\frac{\varepsilon}{2} \right)^p \right) \right)^{1/p}, \sum \left(-\frac{\varepsilon}{2} \right), 0, 0, \dots \right). \end{aligned} \quad (2.11)$$

Then $\|\Delta u\|_{m(\phi, p)} = \|u\|_{m(\phi, p, \Delta)} = 1$, $\|\Delta v\|_{m(\phi, p)} = \|v\|_{m(\phi, p, \Delta)} = 1$, that is, $u, v \in S(m(\phi, p, \Delta))$ and $\|\Delta u - \Delta v\|_{m(\phi, p)} = \|u - v\|_{m(\phi, p, \Delta)} = \varepsilon$.

For $0 \leq \alpha \leq 1$,

$$\|\alpha u + (1 - \alpha)v\|_{m(\phi, p, \Delta)}^p = \|\alpha \Delta u + (1 - \alpha)\Delta v\|_{m(\phi, p)}^p = 1 - \left(\frac{\varepsilon}{2} \right)^p + |2\alpha - 1| \left(\frac{\varepsilon}{2} \right)^p. \quad (2.12)$$

Hence

$$\inf_{0 \leq \alpha \leq 1} \|\alpha u + (1 - \alpha)v\|_{m(\phi, p, \Delta)}^p = 1 - \left(\frac{\varepsilon}{2} \right)^p. \quad (2.13)$$

Therefore, for $p \geq 1$

$$\beta_X(\varepsilon) \leq 1 - \left[1 - \left(\frac{\varepsilon}{2} \right)^p \right]^{1/p}. \quad (2.14)$$

This completes the proof of the theorem. \square

COROLLARY 2.4. (i) If $\varepsilon = 2$, then $\beta_X(\varepsilon) \leq 1$ and hence $m(\phi, p, \Delta)$ is strictly convex.

(ii) If $0 < \varepsilon < 2$, then $0 < \beta_X(\varepsilon) < 1$ and hence $m(\phi, p, \Delta)$ is uniformly convex.

Remark 2.5. Note that these results are best possible for the time being, that is, they cannot be readily generalized to the general case because our results also hold for general matrix transformation.

Acknowledgments

The present paper was completed when Professor Mursaleen visited Firat University (May-June, 2007). The author is very much grateful to the Firat University for providing hospitalities. This research was supported by FUBAP (The Management Union of the Scientific Research Projects of Firat University) when the first author visited Firat University under the Project no. 1179.

References

- [1] W. L. C. Sargent, "Some sequence spaces related to the l^p spaces," *Journal of the London Mathematical Society*, vol. 35, no. 2, pp. 161–171, 1960.
- [2] E. Malkowsky and M. Mursaleen, "Matrix transformations between FK-spaces and the sequence spaces $m(\phi)$ and $n(\phi)$," *Journal of Mathematical Analysis and Applications*, vol. 196, no. 2, pp. 659–665, 1995.
- [3] M. Mursaleen, "Some geometric properties of a sequence space related to l^p ," *Bulletin of the Australian Mathematical Society*, vol. 67, no. 2, pp. 343–347, 2003.
- [4] B. C. Tripathy and M. Sen, "On a new class of sequences related to the space l^p ," *Tamkang Journal of Mathematics*, vol. 33, no. 2, pp. 167–171, 2002.
- [5] R. Çolak and M. Et, "On some difference sequence sets and their topological properties," *Bulletin of the Malaysian Mathematical Sciences Society*, vol. 28, no. 2, pp. 125–130, 2005.
- [6] M. Et and R. Çolak, "On some generalized difference sequence spaces," *Soochow Journal of Mathematics*, vol. 21, no. 4, pp. 377–386, 1995.
- [7] H. Kizmaz, "On certain sequence spaces," *Canadian Mathematical Bulletin*, vol. 24, no. 2, pp. 169–176, 1981.
- [8] E. Malkowsky, M. Mursaleen, and S. Suantai, "The dual spaces of sets of difference sequences of order m and matrix transformations," *Acta Mathematica Sinica*, vol. 23, no. 3, pp. 521–532, 2007.
- [9] Y. Cui and H. Hudzik, "On the Banach-Saks and weak Banach-Saks properties of some Banach sequence spaces," *Acta Scientiarum Mathematicarum*, vol. 65, no. 1-2, pp. 179–187, 1999.
- [10] V. I. Gurarii, "Differential properties of the convexity moduli of Banach spaces," *Matematicheskie Issledovaniya*, vol. 2, no. 1, pp. 141–148, 1967.
- [11] L. Sánchez and A. Ullán, "Some properties of Gurarii's modulus of convexity," *Archiv der Mathematik*, vol. 71, no. 5, pp. 399–406, 1998.

M. Mursaleen: Department of Mathematics, Aligarh Muslim University, Aligarh 202002, India
Email address: mursaleenm@gmail.com

Rifat Çolak: Department of Mathematics, Firat University, 23119 Elazığ, Turkey
Email address: rcolak@firat.edu.tr

Mikail Et: Department of Mathematics, Firat University, 23119 Elazığ, Turkey
Email address: mikaillet@yahoo.com