

Research Article

Rearrangement and Convergence in Spaces of Measurable Functions

D. Caponetti, A. Trombetta, and G. Trombetta

Received 3 November 2006; Accepted 25 February 2007

Recommended by Nikolaos S. Papageorgiou

We prove that the convergence of a sequence of functions in the space L_0 of measurable functions, with respect to the topology of convergence in measure, implies the convergence μ -almost everywhere (μ denotes the Lebesgue measure) of the sequence of rearrangements. We obtain nonexpansivity of rearrangement on the space L_∞ , and also on Orlicz spaces L_N with respect to a finitely additive extended real-valued set function. In the space L_∞ and in the space E_Φ , of finite elements of an Orlicz space L_Φ of a σ -additive set function, we introduce some parameters which estimate the Hausdorff measure of noncompactness. We obtain some relations involving these parameters when passing from a bounded set of L_∞ , or L_Φ , to the set of rearrangements.

Copyright © 2007 D. Caponetti et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The notion of rearrangement of a real-valued μ -measurable function was introduced by Hardy et al. in [1]. It has been studied by many authors and leads to interesting results in Lebesgue spaces and, more generally, in Orlicz spaces (see, e.g., [2–5]). The space L_0 is a space of real-valued *measurable functions*, defined on a nonempty set Ω , in which we can give a natural generalization of the topology of convergence in measure using a group pseudonorm which depends on a submeasure defined on the power set $\mathcal{P}(\Omega)$ of Ω (see [6, 7] and the references given there). In the second section of this note we study rearrangements of functions of the space L_0 . The rearrangements belong to the space $T_0([0, +\infty))$ of all real-valued totally μ -measurable functions defined on $[0, +\infty)$. We extend to this setting some convergence results (see, e.g., [3, 5]). Precisely, we prove that the convergence in the space L_0 implies the convergence μ -almost everywhere of rearrangements. Moreover, by the convergence in L_0 of a nondecreasing sequence of nonnegative

functions, we obtain the convergence in measure of the corresponding nondecreasing sequence of rearrangements. In the third section we introduce, in a natural manner, the space L_∞ as the closure of the subspace of all simple functions of L_0 with respect to the essentially supremum norm. The space L_∞ so defined is contained in L_0 , and we prove nonexpansivity of rearrangement on this space. In the last section we obtain nonexpansivity of rearrangement on Orlicz spaces L_N of a finitely additive extended real-valued set function.

We recall (see [8]) that for a bounded subset Y of a normed space $(X, \|\cdot\|)$ the Hausdorff measure of noncompactness $\gamma_X(Y)$ of Y is defined by

$$\gamma_X(Y) = \inf \{ \varepsilon > 0 : \text{there is a finite subset } F \text{ of } X \text{ such that } Y \subseteq \cup_{f \in F} B_X(f, \varepsilon) \}, \tag{1.1}$$

where $B_X(f, \varepsilon) = \{g \in X : \|f - g\| \leq \varepsilon\}$. In sections 3 and 4 we introduce, respectively, a parameter ω_{L_∞} in L_∞ and a parameter ω_{E_Φ} in the space E_Φ of finite elements of a classical Orlicz space L_Φ of a σ -additive set function. By means of these parameters, we derive an exact formula in L_∞ and an estimate in E_Φ for the Hausdorff measure of noncompactness. Then as a consequence of nonexpansivity of rearrangement we obtain inequalities involving such parameters, when passing from a set of functions in L_∞ , or in L_Φ , to the set of rearrangements. We denote by \mathbb{N} , \mathbb{Q} , and \mathbb{R} the set of all natural, rational, and real numbers, respectively.

2. Rearrangements of functions and convergence in the space L_0

Let Ω be a nonempty set and \mathbb{R}^Ω the set of all real-valued functions on Ω with its natural Riesz space structure. Let \mathcal{A} be an algebra in the power set $\mathcal{P}(\Omega)$ of Ω and let $\eta : \mathcal{P}(\Omega) \rightarrow [0, +\infty]$ be a submeasure (i.e., a monotone, subadditive function with $\eta(\emptyset) = 0$). Then

$$\|f\|_0 = \inf \{ a > 0 : \eta(\{|f| > a\}) < a \}, \tag{2.1}$$

where $\{|f| > a\} = \{x \in \Omega : |f(x)| > a\}$ and where $\inf \emptyset = +\infty$ defines a group pseudo-norm on \mathbb{R}^Ω (i.e., $\|0\|_0 = 0$, $\|f\|_0 = \|-f\|_0$ and $\|f + g\|_0 \leq \|f\|_0 + \|g\|_0$ for all $f, g \in \mathbb{R}^\Omega$). We denote by

$$S(\Omega, \mathcal{A}) = \left\{ \sum_{i=1}^n a_i \chi_{A_i} : n \in \mathbb{N}, a_i \in \mathbb{R}, A_i \in \mathcal{A} \right\} \tag{2.2}$$

the space of all real-valued \mathcal{A} -simple functions on Ω ; hereby χ_A denotes the characteristic function of A defined on Ω . By $L_0 := L_0(\Omega, \mathcal{A}, \eta)$ we denote the closure of the space $S(\Omega, \mathcal{A})$ in $(\mathbb{R}^\Omega, \|\cdot\|_0)$.

For each function $f \in \mathbb{R}^\Omega$, set $|f|_\infty = \sup_\Omega |f|$ and denote by $B(\Omega, \mathcal{A})$ the closure of the space $S(\Omega, \mathcal{A})$ in $(\mathbb{R}^\Omega, |\cdot|_\infty)$. As $\|f\|_0 \leq |f|_\infty$, we have $B(\Omega, \mathcal{A}) \subseteq L_0$. If for $M \in \mathcal{P}(\Omega)$ we set $\eta(M) = 0$ if $M = \emptyset$ and $\eta(M) = +\infty$ if $M \neq \emptyset$, then $(L_0, \|\cdot\|_0) = (B(\Omega, \mathcal{A}), |\cdot|_\infty)$. We point out that the space $B(\Omega, \mathcal{P}(\Omega))$ coincides with the space of all real-valued bounded functions defined on Ω , and clearly $B(\Omega, \mathcal{A}) \subseteq B(\Omega, \mathcal{P}(\Omega))$.

Throughout this note, given a finitely additive set function $\nu : \mathcal{A} \rightarrow [0, +\infty]$, we denote by $\nu^* : \mathcal{P}(\Omega) \rightarrow [0, +\infty]$ the submeasure defined by $\nu^*(E) = \inf\{\nu(A) : A \in \mathcal{A} \text{ and } E \subseteq A\}$. Moreover, whenever Ω is a Lebesgue measurable subset of \mathbb{R}^n , we denote by μ the Lebesgue measure on the σ -algebra of all Lebesgue measurable subsets of Ω , we write μ -a.e. for μ -almost everywhere.

Example 2.1 (see [9, Chapter III]). Let Ω be a Lebesgue measurable subset of \mathbb{R}^n , \mathcal{A} the σ -algebra of all Lebesgue measurable subsets of Ω and $\eta = \mu^*$. If $\eta(\Omega) < +\infty$, then L_0 coincides with the space $M(\Omega)$ of all real-valued μ -measurable functions defined on Ω . If $\eta(\Omega) = +\infty$, then L_0 coincides with the space $T_0(\Omega)$ of all real-valued totally μ -measurable functions defined on Ω .

The following definitions are adapted from [10, Chapter 4].

Definition 2.2.

- (i) A subset A of Ω is said to be an η -null set if $\eta(A) = 0$.
- (ii) A function $f \in \mathbb{R}^\Omega$ is said to be an η -null function if $\eta(\{|f| > a\}) = 0$ for every $a > 0$.
- (iii) Two functions $f, g \in \mathbb{R}^\Omega$ are said to be *equal η -almost everywhere*, and is used the notation $f = g$ η -a.e. if $f - g$ is an η -null function.
- (iv) A function $f \in \mathbb{R}^\Omega$ is said to be *dominated η -almost everywhere* by a function g , and is used the notation $f \leq g$ η -a.e. if there exists an η -null function $h \in \mathbb{R}^\Omega$ such that $f \leq g + h$.

Observe that a function $f \in \mathbb{R}^\Omega$ is an η -null function if and only if $\|f\|_0 = 0$.

The *distribution function* η_f of a function $f \in L_0$ is defined by

$$\eta_f(\lambda) = \eta(\{|f| > \lambda\}) \quad (\lambda \geq 0). \quad (2.3)$$

Observe that $\eta_f = \eta_{|f|}$ and η_f may assume the value $+\infty$. In the next proposition, we state some elementary properties of the distribution function η_f (see [2, Chapter 2]).

PROPOSITION 2.3. *Let $f, g \in L_0$ and $a \neq 0$. Then the distribution function η_f of f is non-negative and decreasing. Moreover,*

- (i) $\eta_{af}(\lambda) = \eta_f(\lambda/|a|)$ for each $\lambda \geq 0$,
- (ii) $\eta_{f+g}(\lambda_1 + \lambda_2) \leq \eta_f(\lambda_1) + \eta_g(\lambda_2)$ for each $\lambda_1, \lambda_2 \geq 0$.

PROPOSITION 2.4. *Let $f, g \in L_0$. If $\|f - g\|_0 = 0$ then $\eta_f = \eta_g$ μ -a.e.*

Proof. Let $f, g \in L_0$ and $h \in L_0$ be an η -null function such that $g = f + h$. Let I and J denote the intervals $\{\lambda \geq 0 : \eta_f(\lambda) = +\infty\}$ and $\{\lambda \geq 0 : \eta_g(\lambda) = +\infty\}$, respectively. We start by proving that $\mu(I) = \mu(J)$. Assume $\mu(I) \neq \mu(J)$ and $\mu(I) < \mu(J)$. Then $I \subset J$ and $\mu(J \setminus I) > 0$. Denoted by $\text{int}(J \setminus I)$ the interior of the interval $J \setminus I$, we have $\eta_g(\lambda) = +\infty$ and $\eta_f(\lambda) < +\infty$ for each $\lambda \in \text{int}(J \setminus I)$. Fix $\lambda_1 \in \text{int}(J \setminus I)$ and $\lambda_2 > 0$ such that $\lambda_1 + \lambda_2 \in \text{int}(J \setminus I)$. By property (ii) of Proposition 2.3, we have

$$+\infty = \eta_g(\lambda_1 + \lambda_2) = \eta_{f+h}(\lambda_1 + \lambda_2) \leq \eta_f(\lambda_1) + \eta_h(\lambda_2) = \eta_f(\lambda_1) < +\infty, \quad (2.4)$$

that is a contradiction. Set $\bar{\lambda} = \sup I = \sup J$ and let $\lambda_0 \in [\bar{\lambda}, +\infty)$ be a point of continuity of both the functions η_f and η_g . By property (ii) of Proposition 2.3, it follows that

$$\eta_f(\lambda_0) = \lim_n \eta_{g-h}\left(\lambda_0 + \frac{1}{n}\right) \leq \eta_g(\lambda_0) + \lim_n \eta_h\left(\frac{1}{n}\right) = \eta_g(\lambda_0). \tag{2.5}$$

Similarly, we find $\eta_g(\lambda_0) \leq \eta_f(\lambda_0)$. Hence $\eta_f = \eta_g$ μ -a.e. □

PROPOSITION 2.5. *Let $f, g \in L_0$. If $|f| \leq |g|$ η -a.e., then $\eta_f \leq \eta_g$ μ -a.e.*

Proof. Let $h \in L_0$ be an η -null function such that $|f| \leq |g| + h$. Then $\eta_{|f|} \leq \eta_{|g|+h}$ and, by Proposition 2.4, $\eta_{|g|} = \eta_{|g|+h}$ μ -a.e. Hence $\eta_{|f|} \leq \eta_{|g|}$ μ -a.e., which gives the assert. □

Observe that, when $(\Omega, \mathcal{A}, \nu)$ is a totally σ -finite measure space and $\eta = \nu^*$, the distribution function η_f of $f \in L_0$ is right continuous (see [2]). In our setting this is not true anymore, as the following example shows.

Example 2.6 (see [9, Chapter III, page 103]). Let $\Omega = [0, 1)$ and let \mathcal{A} be the algebra of all finite unions of right-open intervals contained in Ω . Denote again by μ the Lebesgue measure μ restricted to \mathcal{A} . Let $\eta = \mu^*$. Consider the function $f : [0, 1) \rightarrow \mathbb{R}$ defined as $f(x) = 0$, if $x \in [0, 1) \setminus \mathbb{Q}$, and as $f(x) = 1/q$, if $x = p/q \in [0, 1) \cap \mathbb{Q}$ in lowest terms. Then $\|f\|_0 = 0$ and so f is an η -null function but f is not null μ -a.e. since $\eta(\{|f| > 0\}) = 1$. Moreover, $\eta_f(\lambda) = 0$ if $\lambda > 0$ and $\eta_f(0) = 1$. Then η_f is not right continuous in 0.

Throughout, without loss of generality, we will assume that the distribution function η_f of a function $f \in L_0$ is right continuous, which together with Proposition 2.4 yields $\eta_f = \eta_g$ whenever $f, g \in L_0$ and $\|f - g\|_0 = 0$.

The *decreasing rearrangement* f^* of a function $f \in L_0$ is defined by

$$f^*(t) = \inf \{ \lambda \geq 0 : \eta_f(\lambda) \leq t \} \quad (t \geq 0). \tag{2.6}$$

Clearly, by the above assumption on η_f , $f^* = g^*$ if $f, g \in L_0$ with $\|f - g\|_0 = 0$.

PROPOSITION 2.7. *Let $f \in L_0$. If $f^*(t) = +\infty$, then $t = 0$.*

Proof. Assume that $f^*(t) = +\infty$. Then $\eta_f(\lambda) > t$ for all $\lambda \geq 0$. Since $\|f\|_0 < +\infty$, for some $\bar{\lambda} \geq 0$ we have $\eta_f(\bar{\lambda}) < +\infty$. Hence, as η_f is decreasing, there exists finite $\lim_{\lambda \rightarrow +\infty} \eta_f(\lambda) = l \geq 0$. The thesis follows by proving that $l = 0$. Assume $l > 0$ and choose a function $s \in S(\Omega, \mathcal{A})$ such that $\|f - s\|_0 \leq l/2$.

Fix $\lambda > l + \max_{\Omega} |s|$ and put $A = \{|f| > \lambda\}$, then $\eta(A) = \eta_f(\lambda) \geq l$ and

$$|f(x) - s(x)| \geq ||f(x)| - |s(x)|| \geq l \tag{2.7}$$

for each $x \in A$. So that $\|f - s\|_0 \geq l$. So we obtain $l \leq \|f - s\|_0 \leq l/2$: a contradiction. □

The following proposition contains some properties of rearrangements of functions of L_0 . The proofs of (i)–(iv) (except some slight modifications) are identical to that of [2] for rearrangements of functions of a Banach function space, and we omit them.

PROPOSITION 2.8. Let $f, g \in L_0$ and $a \in \mathbb{R}$. Then f^* is nonnegative, decreasing, and right continuous. Moreover,

- (i) $(af)^* = |a|f^*$;
- (ii) $f^*(\eta_f(\lambda)) \leq \lambda$, ($\eta_f(\lambda) < +\infty$) and $\eta_f(f^*(t)) \leq t$, ($f^*(t) < +\infty$);
- (iii) $(f+g)^*(t_1+t_2) \leq f^*(t_1) + g^*(t_2)$ for each $t_1, t_2 \geq 0$;
- (iv) if $|f| \leq |g|$ η -a.e., then $f^* \leq g^*$ μ -a.e.

Proof. Clearly f^* is nonnegative and decreasing. We prove that f^* is right continuous. Fix $t_0 \geq 0$ and assume that $\lim_{t \rightarrow t_0^+} f^*(t) = a < f^*(t_0) < +\infty$. Choose $b \in (a, f^*(t_0))$. Observe that, since $b < f^*(t_0)$, we have that $\eta_f(b) > t_0$ by the definition of f^* . Moreover, since $\lim_{t \rightarrow t_0^+} f^*(t) = a$, there exists $t_1 > 0$ such that $t_0 < t_1 < \eta_f(b)$ and $f^*(t_1) < b$. From the definition of f^* we obtain that $\eta_f(b) \leq t_1$. It follows that $t_1 < \eta_f(b) \leq t_1$ which is a contradiction. Then $\lim_{t \rightarrow t_0^+} f^*(t) = f^*(t_0)$.

To complete the proof, suppose that $f^*(0) = +\infty$, and assume that $\lim_{t \rightarrow 0^+} f^*(t) = a < +\infty$. Choose $b > a$. Then $\eta_f(b) > 0$ and since $\lim_{t \rightarrow 0^+} f^*(t) = a$ we have that there exists $t_2 > 0$ such that $t_2 < \eta_f(b)$ and $f^*(t_2) < b$. From the definition of f^* we obtain that $\eta_f(b) \leq t_2$. It follows that $t_2 < \eta_f(b) \leq t_2$ which is contradiction. Hence $\lim_{t \rightarrow 0^+} f^*(t) = +\infty$. \square

Now we show that the rearrangement of a function of L_0 is a function of the space $T_0([0, +\infty))$ of all real-valued totally μ -measurable functions defined on $[0, +\infty)$, introduced in [9, Chapter III, Definition 10] (see also Example 2.1). In $T_0([0, +\infty))$, we write $|\cdot|_0$ instead of $\|\cdot\|_0$.

THEOREM 2.9. Let $f \in L_0$. Then

- (i) f and f^* are equimeasurable, that is, $\eta_f(\lambda) = \mu_{f^*}(\lambda)$ for all $\lambda \geq 0$;
- (ii) $f^* \in T_0([0, +\infty))$ and $|f^*|_0 = \|f\|_0$.

Proof. (i) Fixed $\lambda \geq 0$ such that $\eta_f(\lambda) < +\infty$, by the first inequality of property (ii) of Proposition 2.8, we have that $f^*(\eta_f(\lambda)) \leq \lambda$. Moreover, since f^* is decreasing, we have $f^*(t) \leq \lambda$ for each t such that $\eta_f(\lambda) < t$. It follows that $\mu_{f^*}(\lambda) = \sup\{f^* > \lambda\} \leq \eta_f(\lambda)$. It remains to prove that $\eta_f(\lambda) \leq \mu_{f^*}(\lambda)$. Suppose that $f^*(0) = +\infty$. Then $\mu_{f^*}(\lambda) = \sup\{f^* > \lambda\}$ for all $\lambda \geq 0$. Assume that there exists $\lambda_0 \geq 0$ such that $\eta_f(\lambda_0) > \mu_{f^*}(\lambda_0)$. Fixed $t \in (\mu_{f^*}(\lambda_0), \eta_f(\lambda_0))$, we have that $f^*(t) \leq \lambda_0$ since $t > \mu_{f^*}(\lambda_0) = \sup\{f^* > \lambda_0\}$. On the other hand, since $t < \eta_f(\lambda_0)$, by the definition of f^* , we obtain $f^*(t) > \lambda_0$ which is a contradiction. The same proof breaks down if $f^*(0) < +\infty$ and $\lambda < f^*(0)$. If $f^*(0) < +\infty$ and $\lambda \geq f^*(0)$ then $\mu_{f^*}(\lambda) = 0$. Moreover, by the second part of the property (ii) of Proposition 2.8, it follows that $\eta_f(f^*(0)) = 0$ and then $\eta_f(\lambda) = 0$ for all $\lambda \geq f^*(0)$. This completes the proof.

(ii) is an immediate consequence of (i). \square

The next theorem states two well-known convergence results (see, e.g., [5, Lemma 1.1] and [3, Lemma 2], resp.).

THEOREM 2.10. Let Ω be a Lebesgue measurable subset of \mathbb{R}^n , and let $\{f_n\}$ be a sequence of elements of the space $T_0(\Omega)$ of all real-valued totally μ -measurable functions defined on Ω .

- (i) If $\{f_n\}$ converges in measure to f , then $f_n^*(t)$ converges to $f^*(t)$ in each point t of continuity of f^* .
- (ii) If $\{f_n\}$ is a nondecreasing sequence of nonnegative functions convergent to f μ -a.e, then f_n^* is a nondecreasing sequence convergent to f^* pointwise.

The remainder of this section will be devoted to extend these convergence results to the general setting of the space L_0 . We need the following lemma.

LEMMA 2.11. Let $f_n, f \in L_0$ ($n = 1, 2, \dots$) be such that $\|f_n - f\|_0 \rightarrow 0$. Then $\eta_{f_n}(\lambda) \rightarrow \eta_f(\lambda)$ for each point λ of continuity of η_f . Moreover, if $\lim_{\lambda \rightarrow \lambda_0^+} \eta_f(\lambda) = +\infty$ then $\lim_{n \rightarrow +\infty} \eta_{f_n}(\lambda_0) = +\infty$.

Proof. Let $\lambda > 0$ be a point of continuity of η_f and assume $\eta_{f_n}(\lambda) \not\rightarrow \eta_f(\lambda)$. Then there are $\varepsilon_0 > 0$ and a subsequence $(\eta_{f_{n_k}})$ of (η_{f_n}) such that $|\eta_{f_{n_k}}(\lambda) - \eta_f(\lambda)| > \varepsilon_0$ for each $k \in \mathbb{N}$. Put

$$I_1 = \{k \in \mathbb{N} : \eta_{f_{n_k}}(\lambda) > \eta_f(\lambda) + \varepsilon_0\}, \quad I_2 = \{k \in \mathbb{N} : \eta_{f_{n_k}}(\lambda) < \eta_f(\lambda) - \varepsilon_0\}. \quad (2.8)$$

Either I_1 or I_2 is infinite. Let $h > 0$ such that

$$\eta_f(\lambda - h) < \eta_f(\lambda) + \frac{\varepsilon_0}{2}, \quad \eta_f(\lambda + h) > \eta_f(\lambda) - \frac{\varepsilon_0}{2}. \quad (2.9)$$

Suppose I_1 is infinite and let $k \in I_1$. Consider the sets

$$\begin{aligned} A_{\lambda-h} &= \{x \in \Omega : |f(x)| > \lambda - h\}, \\ A_{n_k, \lambda} &= \{x \in \Omega : |f_{n_k}(x)| > \lambda\}. \end{aligned} \quad (2.10)$$

Then $\eta(A_{\lambda-h}) = \eta_f(\lambda - h)$ and $\eta(A_{n_k, \lambda}) = \eta_{f_{n_k}}(\lambda)$. We have that $\eta_{f_{n_k}}(\lambda) - \eta_f(\lambda - h) > \varepsilon_0/2$. Moreover,

$$\eta(A_{n_k, \lambda} \setminus A_{\lambda-h}) \geq \eta(A_{n_k, \lambda}) - \eta(A_{\lambda-h}) > \frac{\varepsilon_0}{2}. \quad (2.11)$$

Let $x \in A_{n_k, \lambda} \setminus A_{\lambda-h}$. Then $|f(x)| \leq \lambda - h$ and $|f_{n_k}(x)| > \lambda$. Therefore $|f_{n_k}(x)| - |f(x)| > h$. Hence

$$\begin{aligned} \eta(\{x \in \Omega : |f_{n_k}(x) - f(x)| > h\}) &\geq \eta(\{x \in \Omega : |f_{n_k}(x)| - |f(x)| > h\}) \\ &\geq \eta(A_{n_k, \lambda} \setminus A_{\lambda-h}) > \frac{\varepsilon_0}{2}, \end{aligned} \quad (2.12)$$

and this is a contradiction since $\|f_n - f\|_0 \rightarrow 0$. The proof is similar in the case the set I_2 is infinite. The second part of the proposition follows analogously. \square

THEOREM 2.12. Let $f_n, f \in L_0$ ($n = 1, 2, \dots$) be such that $\|f_n - f\|_0 \rightarrow 0$. Then $f_n^*(t) \rightarrow f^*(t)$ for each point t of continuity of f^* . Moreover, if $\lim_{t \rightarrow 0^+} f^*(t) = +\infty$ then $\lim_{n \rightarrow +\infty} f_n^*(0) = +\infty$.

Proof. Let $t_0 > 0$ be a point of continuity of f^* and assume $f_n^*(t_0) \not\rightarrow f^*(t_0)$. Then there are $\varepsilon_0 > 0$ and a subsequence $(f_{n_k}^*)$ of (f_n^*) such that $|f_{n_k}^*(t_0) - f^*(t_0)| > \varepsilon_0$ for each $k \in \mathbb{N}$.

Put

$$I_1 = \{k \in \mathbb{N} : f_{n_k}^*(t_0) > f^*(t_0) + \varepsilon_0\}, \quad I_2 = \{k \in \mathbb{N} : f_{n_k}^*(t_0) < f^*(t_0) - \varepsilon_0\}. \quad (2.13)$$

Either I_1 or I_2 is infinite. Let $h > 0$ such that

$$f^*(t_0 - h) < f^*(t_0) + \frac{\varepsilon_0}{2}, \quad f^*(t_0 + h) > f^*(t_0) - \frac{\varepsilon_0}{2}. \quad (2.14)$$

Suppose I_1 is infinite. Fix $k \in I_1$, $t \in [t_0 - h, t_0]$ and $\sigma \in [f^*(t_0) + \varepsilon_0/2, f^*(t_0) + \varepsilon_0]$. Then

$$\begin{aligned} f^*(t) &\leq f^*(t_0) + \frac{\varepsilon_0}{2} \leq \sigma, \\ f_{n_k}^*(t) &> f^*(t_0) + \varepsilon_0 \geq \sigma. \end{aligned} \quad (2.15)$$

Hence $\eta_f(\sigma) \leq t_0 - h < t_0$ and $\eta_{f_k}(\sigma) \geq t_0$. This shows that $\eta_{f_n}(\sigma) \not\rightarrow \eta_f(\sigma)$ for all $k \in I_1$ and $\sigma \in [f^*(t_0) + \varepsilon_0/2, f^*(t_0) + \varepsilon_0]$ which by Lemma 2.11 is a contradiction. The second implication follows similarly. \square

LEMMA 2.13. *Let $f_n, f \in L_0$ ($n = 1, 2, \dots$) be such that $\{f_n\}$ is a nondecreasing sequence of nonnegative functions and $\|f_n - f\|_0 \rightarrow 0$. Then $|\eta_{f_n} - \eta_f|_0 \rightarrow 0$.*

Proof. Assume by contradiction $|\eta_{f_n} - \eta_f|_0 \not\rightarrow 0$. Since $\eta_{f_n} \leq \eta_{f_{n+1}} \leq \eta_f$, we find $\varepsilon_0 > 0$, $\sigma_0 > 0$ and $\bar{n} \in \mathbb{N}$ such that

$$\mu(\{\lambda \geq 0 : \eta_f(\lambda) - \eta_{f_n}(\lambda) > \varepsilon_0\}) > \sigma_0 \quad (2.16)$$

for all $n \in \mathbb{N}$ with $n \geq \bar{n}$. Set $B_n = \{\lambda \geq 0 : \eta_f(\lambda) - \eta_{f_n}(\lambda) > \varepsilon_0\}$, then $\cap_{n \geq \bar{n}} B_n$ is nonempty, and for $\lambda_0 \in \cap_{n \geq \bar{n}} B_n$ we have

$$\sup_{n \geq \bar{n}} \eta_{f_n}(\lambda_0) \leq \eta_f(\lambda_0) - \varepsilon_0. \quad (2.17)$$

Then we choose $h > 0$ such that

$$\eta_f(\lambda_1) - \eta_{f_n}(\lambda_2) \geq \frac{\varepsilon_0}{2} \quad (2.18)$$

for all $\lambda_1, \lambda_2 \in [\lambda_0, \lambda_0 + h]$ and all $n \geq \bar{n}$. In particular, we have

$$\eta_f(\lambda_0 + h) - \eta_{f_n}(\lambda_0) \geq \frac{\varepsilon_0}{2}. \quad (2.19)$$

Then using the same notations and considerations similar to that of Lemma 2.11, we find

$$\begin{aligned} \{x \in \Omega : f(x) - f_n(x) > h\} &\supseteq A_{\lambda_0+h} \setminus A_{n, \lambda_0}, \\ \eta(A_{\lambda_0+h} \setminus A_{n, \lambda_0}) &\geq \eta_f(\lambda_0 + h) - \eta_{f_n}(\lambda_0) \geq \frac{\varepsilon_0}{2} \end{aligned} \quad (2.20)$$

which is a contradiction since $\|f_n - f\|_0 \rightarrow 0$. \square

THEOREM 2.14. *Let $f_n, f \in L_0$ ($n = 1, 2, \dots$) be such that $\{f_n\}$ is a nondecreasing sequence of nonnegative functions and $\|f_n - f\|_0 \rightarrow 0$. Then $|f_n^* - f^*|_0 \rightarrow 0$.*

Proof. The proof, using Lemma 2.13, is analogous to the proof of Theorem 2.12. \square

We remark that if $\{f_n\}$ is a sequence of elements of the space $T_0(\Omega)$, Theorem 2.14 yields (ii) of Theorem 2.10.

3. Nonexpansivity of rearrangement in the space L_∞

We introduce the notion of essentially boundedness, following [10]. For $f \in \mathbb{R}^\Omega$, set

$$\|f\|_\infty = \inf_{A \subseteq \Omega, \eta(A) = 0} \sup_{\Omega \setminus A} |f|, \tag{3.1}$$

then $\|\cdot\|_\infty$ defines a group pseudonorm on \mathbb{R}^Ω , for each submeasure η on $\mathcal{P}(\Omega)$.

We recall that, if ν is a finitely additive extended real-valued set function on an algebra $\mathcal{A} \subseteq \mathcal{P}(\Omega)$ and $\eta = \nu^*$, the space $\mathcal{L}_\infty(\Omega, \mathcal{A}, \nu)$ of all real-valued essentially bounded functions introduced in [10] is defined by

$$\mathcal{L}_\infty(\Omega, \mathcal{A}, \nu) = \{f \in \mathbb{R}^\Omega : \|f\|_\infty < +\infty\}. \tag{3.2}$$

In our setting it is natural to define a space $L_\infty(\Omega, \mathcal{A}, \eta)$ of all real-valued essentially bounded functions as follows.

Definition 3.1. The space $L_\infty := L_\infty(\Omega, \mathcal{A}, \eta)$ is the closure of the space $S(\Omega, \mathcal{A})$ in $(\mathbb{R}^\Omega, \|\cdot\|_\infty)$.

Let $f \in \mathbb{R}^\Omega$. Since $\|f\|_0 \leq \|f\|_\infty$, we have $L_\infty \subseteq L_0$. Moreover, $\|f\|_0 = 0$ if and only if $\|f\|_\infty = 0$. In the remainder part of this note we will identify functions $f, g \in \mathbb{R}^\Omega$ for which $\|f - g\|_0 = 0$. Then $(L_0, \|\cdot\|_0)$ and $(L_\infty, \|\cdot\|_\infty)$ become an F -normed space (in the sense of [11]) and a normed space, respectively.

PROPOSITION 3.2. *Let ν be a finitely additive extended real-valued set function on an algebra \mathcal{A} in $\mathcal{P}(\Omega)$ and $\eta = \nu^*$. Then the space $\mathcal{L}_\infty(\Omega, \mathcal{A}, \nu)$ coincides with the space $L_\infty(\Omega, \mathcal{P}(\Omega), \eta)$.*

Proof. Given $f \in L_\infty(\Omega, \mathcal{P}(\Omega), \eta)$, find a simple function $s \in S(\Omega, \mathcal{P}(\Omega))$ such that $\|f - s\|_\infty < +\infty$. From $\|f\|_\infty \leq \|f - s\|_\infty + \|s\|_\infty$, we get $f \in \mathcal{L}_\infty(\Omega, \mathcal{A}, \nu)$. On the other hand, if $f \in \mathcal{L}_\infty(\Omega, \mathcal{A}, \nu)$ then there exists $A \subseteq \Omega$ such that $\eta(A) = 0$ and such that $\sup_{\Omega \setminus A} |f| < +\infty$. Consider the real function g on Ω defined by $g = f$ on $\Omega \setminus A$ and by $g = 0$ on A . Of course $g \in \mathcal{L}_\infty(\Omega, \mathcal{A}, \nu)$ and $\|f - g\|_\infty = 0$. Moreover, $g \in B(\Omega, \mathcal{P}(\Omega)) \subseteq L_\infty(\Omega, \mathcal{P}(\Omega), \eta)$. Then there exists a sequence (s_n) in $S(\Omega, \mathcal{P}(\Omega))$ such that $\|g - s_n\|_\infty \rightarrow 0$. Since $\|f - s_n\|_\infty \leq \|f - g\|_\infty + \|g - s_n\|_\infty = \|g - s_n\|_\infty$, we have that $f \in L_\infty(\Omega, \mathcal{P}(\Omega), \eta)$. \square

We write briefly $B([0, +\infty))$ instead of $B(\Omega, \mathcal{A})$, when $\Omega = [0, +\infty)$, \mathcal{A} is the σ -algebra of all Lebesgue measurable subsets of Ω and $\eta = \mu^*$. The next proposition establishes that the rearrangement of a function of L_∞ is a function of $B([0, +\infty))$.

PROPOSITION 3.3. *Let $f \in L_\infty$. Then $f^* \in B([0, +\infty))$ and $\|f^*\|_\infty = f^*(0) = \|f\|_\infty$.*

Proof. Let $\varepsilon > 0$. Then there is $A \subseteq \Omega$ such that $\eta(A) = 0$ and $\sup_{\Omega \setminus A} |f| < \|f\|_\infty + \varepsilon$. Hence $\{|f| > \|f\|_\infty + \varepsilon\} \subseteq A$, so that $\eta(\{|f| > \|f\|_\infty + \varepsilon\}) = 0$.

Therefore $|f^*|_\infty = f^*(0) \leq \|f\|_\infty + \varepsilon$ so that $|f^*|_\infty \leq \|f\|_\infty$. Now we have to prove that $\|f\|_\infty \leq |f^*|_\infty$. Assume $|f^*|_\infty < c < \|f\|_\infty$. Then for each $A \subseteq \Omega$ such that $\eta(A) = 0$ we have $\sup_{\Omega \setminus A} |f| > c$ and $\eta_f(c) = \eta(\{|f| > c\}) > 0$. For $t \in [0, \eta_f(c))$, by the definition of the function f^* , we obtain $f^*(t) \geq c > |f^*|_\infty = f^*(0)$ which is a contradiction, since f^* is decreasing. \square

Our next aim is to prove nonexpansivity of rearrangement on L_∞ . We need the following two lemmas.

LEMMA 3.4. *Let $s_1, s_2 \in S(\Omega, \mathcal{A})$. Then $|s_1^* - s_2^*|_\infty \leq \|s_1 - s_2\|_\infty$.*

Proof. Let $s_1, s_2 \in S(\Omega, \mathcal{A})$ and put $\|s_1 - s_2\|_\infty = \varepsilon$. Let $\{A_1, \dots, A_n\}$ be a finite partition of Ω in \mathcal{A} such that $s_1 = \sum_{i=1}^n a_i \chi_{A_i}$ and $s_2 = \sum_{i=1}^n b_i \chi_{A_i}$. Set

$$s = \sum_{i=1}^n \min\{|a_i|, |b_i|\} \chi_{A_i \setminus A}, \quad (3.3)$$

where $\eta(A) = 0$ and $|s_1(x) - s_2(x)| \leq \varepsilon$ for all $x \in \Omega \setminus A$. It suffices to prove that

$$s(x) \leq |s_1(x)| \leq s_\varepsilon(x), \quad s(x) \leq |s_2(x)| \leq s_\varepsilon(x), \quad (3.4)$$

for all $x \in \Omega \setminus A$, where $s_\varepsilon = |s| + \varepsilon$. In fact, from this and from property (iv) of Proposition 2.8, it follows that

$$s^* \leq s_1^* \leq s_\varepsilon^* \mu\text{-a.e.}, \quad s^* \leq s_2^* \leq s_\varepsilon^* \mu\text{-a.e.}, \quad (3.5)$$

and thus $|s_1^* - s_2^*|_\infty \leq |s_\varepsilon^* - s^*|_\infty = \varepsilon$. Fix $x \in \Omega \setminus A$ and let $i \in \{1, \dots, n\}$ such that $x \in A_i \setminus A$. Now, if $s(x) = |a_i|$ we have

$$s(x) = |s_1(x)| \leq |a_i| + \varepsilon = s_\varepsilon(x). \quad (3.6)$$

If $s(x) = |b_i|$, since $\|s_1 - s_2\|_\infty = \varepsilon$ implies $0 \leq |a_i| - |b_i| \leq |a_i - b_i| \leq \varepsilon$, we have

$$s(x) \leq |a_i| = |s_1(x)| \leq |b_i| + \varepsilon = s_\varepsilon(x). \quad (3.7)$$

Analogously we obtain $s(x) \leq |s_2(x)| \leq s_\varepsilon(x)$ for $x \in \Omega \setminus A$, and the lemma follows. \square

LEMMA 3.5. *Let $f \in L_\infty$. Then for each $\varepsilon > 0$ there exists a function $s \in S(\Omega, \mathcal{A})$ such that $\|f - s\|_\infty \leq \varepsilon/2$ and $|f^* - s^*|_\infty \leq \varepsilon$.*

Proof. Fix $\varepsilon > 0$. Then similar to [10, page 101] (see Theorem 3.10), we have that there is a finite partition $\{A_1, \dots, A_n\}$ of Ω in \mathcal{A} and $A \subseteq \Omega$ with $\eta(A) = 0$ such that

$$\sup_{x, y \in A_i \setminus A} |f(x) - f(y)| \leq \varepsilon \quad (3.8)$$

for each $i \in \{1, \dots, n\}$. Set

$$\lambda_i = \inf_{x \in A_i \setminus A} |f(x)|, \quad \Lambda_i = \sup_{x \in A_i \setminus A} |f(x)|, \quad a_i = \frac{\lambda_i + \Lambda_i}{2}, \quad (3.9)$$

for each $i \in \{1, \dots, n\}$. Define the simple function

$$s = \sum_{i=1}^n a_i \chi_{A_i}. \tag{3.10}$$

Then for each $i \in \{1, \dots, n\}$ and for each $x \in A_i \setminus A$ we have $|f(x) - s(x)| \leq \varepsilon/2$. Hence $\|f - s\|_\infty \leq \varepsilon/2$. Now consider the simple function φ defined by

$$\varphi(x) = \begin{cases} \left| a_i + \frac{\varepsilon}{2} \right|, & \text{if } x \in A_i, a_i < -\frac{\varepsilon}{2}, \\ 0, & \text{if } x \in A_i, -\frac{\varepsilon}{2} \leq a_i \leq \frac{\varepsilon}{2}, \\ \left| a_i - \frac{\varepsilon}{2} \right|, & \text{if } x \in A_i, a_i > \frac{\varepsilon}{2}. \end{cases} \tag{3.11}$$

Then a direct computation shows that

$$\varphi(x) \leq |f(x)| \leq \varphi_\varepsilon(x), \quad \varphi(x) \leq |s(x)| \leq \varphi_\varepsilon(x), \tag{3.12}$$

for all $x \in \Omega \setminus A$, where $\varphi_\varepsilon = |\varphi| + \varepsilon$. Put $h(x) = (\max |a_i|) \chi_A(x)$ and $k(x) = |f(x)| \chi_A(x)$. Then $\varphi \leq |f| + h$ and $|f| \leq \varphi_\varepsilon + k$. As h and k are both η -null functions, from the property (iv) of Proposition 2.8 it follows that $\varphi^* \leq f^* \leq \varphi_\varepsilon^* \mu$ -a.e., and analogously $\varphi^* \leq s^* \leq \varphi_\varepsilon^* \mu$ -a.e., hence $|f^* - s^*|_\infty \leq |\varphi_\varepsilon^* - \varphi^*|_\infty = \varepsilon$. \square

THEOREM 3.6. *Let $f, g \in L_\infty$. Then $|f^* - g^*|_\infty \leq \|f - g\|_\infty$.*

Proof. Let $\varepsilon > 0$. By Lemma 3.5 we can find $s, u \in S(\Omega, \mathcal{A})$ such that

$$\begin{aligned} \|f - s\|_\infty &\leq \frac{\varepsilon}{4}, & \|g - u\|_\infty &\leq \frac{\varepsilon}{4}, \\ |f^* - s^*|_\infty &\leq \frac{\varepsilon}{2}, & |g^* - u^*|_\infty &\leq \frac{\varepsilon}{2}. \end{aligned} \tag{3.13}$$

We have that

$$\|s - u\|_\infty \leq \|f - s\|_\infty + \|f - g\|_\infty + \|g - u\|_\infty \leq \|f - g\|_\infty + \frac{\varepsilon}{2}. \tag{3.14}$$

Then the last inequality and Lemma 3.4 imply $|s^* - u^*|_\infty \leq \|f - g\|_\infty + \varepsilon/2$.

Consequently we have

$$|f^* - g^*|_\infty \leq |f^* - s^*|_\infty + |s^* - u^*|_\infty + |g^* - u^*|_\infty \leq \|f - g\|_\infty + \varepsilon, \tag{3.15}$$

and by the arbitrariness of ε the theorem follows. \square

Remark 3.7. We observe that Theorem 3.6 does not hold in every space L_0 . In fact, let $L_0 = M([0, 1])$ (see Example 2.1) and set

$$s_n = \sum_{i=0}^{n-1} (n - i) \chi_{[i/n, (i+1)/n)}, \quad t_n = \sum_{i=1}^{n-1} (n - i) \chi_{[i/n, (i+1)/n)}, \tag{3.16}$$

for $n = 2, 3, \dots$. Then for each n we have $t_n = s_n \chi_{[1/n, 1]}$, $s_n - t_n = n \chi_{[0, 1/n]}$, and $|s_n - t_n|_0 = 1/n$. On the other hand, since $s_n^* = s_n$ and $t_n^* = \sum_{i=0}^{n-1} (n-1-i) \chi_{[i/n, (i+1)/n]}$, we have that $s_n^* - t_n^* = \chi_{[0, 1]}$ and then $|s_n^* - t_n^*|_0 = 1$.

Throughout for a set M in L_0 , we put $M^* = \{f^* : f \in M\}$. The following inequality between the Hausdorff measure of noncompactness of a bounded subset M of L_∞ and that of M^* is an immediate consequence of nonexpansivity of rearrangement on L_∞ .

COROLLARY 3.8. *Let M be a bounded subset of L_∞ . Then*

$$\gamma_{B((0, +\infty))}(M^*) \leq \gamma_{L_\infty}(M). \quad (3.17)$$

The following example shows that there is not any constant c such that $\gamma_{L_\infty}(M) \leq c \gamma_{B((0, +\infty))}(M^*)$.

Example 3.9. Let $M = \{\chi_I : I \subseteq [0, 1], \mu(I) = 1/2\}$. Then $M^* = \{\chi_{[0, 1/2]}\}$ and we have that $\gamma_{B((0, +\infty))}(M^*) = 0$ while $\gamma_{L_\infty}(M) > 0$.

In order to obtain a precise formula for the Hausdorff measure of noncompactness in the space L_∞ , we consider for any bounded subset M of L_∞ the following parameter:

$$\omega_{L_\infty}(M) = \inf \left\{ \varepsilon > 0 : \text{there exists a finite partition } \{A_1, \dots, A_n\} \right. \\ \left. \text{of } \Omega \text{ in } \mathcal{A} \text{ such that for all } f \in M \text{ there is } A_f \subseteq \Omega \right. \\ \left. \text{with } \eta(A_f) = 0 \text{ and } \sup_{x, y \in A_i \setminus A_f} |f(x) - f(y)| \leq \varepsilon \text{ for all } i = 1, \dots, n \right\}. \quad (3.18)$$

The proof of the following result is similar to that of [12, Theorem 2.1].

THEOREM 3.10. *Let M be a bounded subset of L_∞ . Then*

$$\gamma_{L_\infty}(M) = \frac{1}{2} \omega_{L_\infty}(M). \quad (3.19)$$

Proof. Fix $a > \gamma_{L_\infty}(M)$. Then we can find $s_1, \dots, s_n \in S(\Omega, \mathcal{A})$ such that for each $f \in M$ there is $i \in \{1, \dots, n\}$ with $\|f - s_i\|_\infty \leq a$. Let $\{A_1, \dots, A_m\}$ be a partition of Ω in \mathcal{A} such that the restriction $s_i|_{A_j}$ is constant for all $i \in \{1, \dots, n\}$ and for all $j \in \{1, \dots, m\}$. Let $f \in M$, $i \in \{1, \dots, n\}$, and $A_f \subseteq \Omega$ such that $\eta(A_f) = 0$ and $\sup_{\Omega \setminus A_f} |f - s_i| \leq a$. For each $j \in \{1, \dots, m\}$, we have that

$$\sup_{x, y \in A_i \setminus A_f} |f(x) - f(y)| \leq 2a, \quad (3.20)$$

hence $\omega_{L_\infty}(M) \leq 2\gamma_{L_\infty}(M)$ and $(1/2)\omega_{L_\infty}(M) \leq \gamma_{L_\infty}(M)$.

Now fix $a > \omega_{L_\infty}(M)$ and let $c > 0$ such that $\|f\|_\infty \leq c$ for each $f \in M$. Then there is a finite partition $\{A_1, \dots, A_n\}$ of Ω in \mathcal{A} such that for all $f \in M$ there is $A_f \subseteq \Omega$ with $\eta(A_f) = 0$ and $\sup_{x, y \in A_i \setminus A_f} |f(x) - f(y)| \leq a$ for all $i = 1, \dots, n$. Moreover, for all $f \in M$

there is $B_f \subseteq \Omega$ with $\eta(B_f) = 0$ such that $\sup_{\Omega \setminus B_f} |f| \leq c$. Set $C_f = A_f \cup B_f$ for each $f \in M$. Fix $\varepsilon > 0$. Let $k, m \in \mathbb{N}$ such that $1/m < \varepsilon$ and $-c + k/m > c$. Set $X = \{-c + i/m : i = 0, \dots, k\}$ and $F = \{\sum_{i=1}^n a_i \chi_{A_i} : a_i \in X\}$. Then for each $f \in M$ there is a function $s \in F$ such that $\sup_{\Omega \setminus C_f} |f - s| \leq a/2 + 1/m \leq a/2 + \varepsilon$. Since F is finite it follows that $\gamma_{L_\infty}(M) \leq (1/2)\omega_{L_\infty}(M)$. This completes the proof. \square

Observe that as a particular case of [12, Theorem 2.1], for a bounded subset T of $B([0, +\infty))$ we have

$$\gamma_{B([0, +\infty))}(T) = \frac{1}{2} \omega_{B([0, +\infty))}(T), \tag{3.21}$$

where

$$\omega_{B([0, +\infty))}(T) = \inf \left\{ \varepsilon > 0 : \text{there exists a finite partition } \{A_1, \dots, A_n\} \right. \\ \left. \text{of } [0, +\infty) \text{ of Lebesgue measurable sets such that for all } f \in T \right. \\ \left. \sup_{x, y \in A_i \setminus A_f} |f(x) - f(y)| \leq \varepsilon \text{ for all } i = 1, \dots, n \right\}. \tag{3.22}$$

In view of the formulas we have obtained, by Corollary 3.8 we have the following.

COROLLARY 3.11. *Let M be a bounded subset of L_∞ . Then*

$$\omega_{B([0, +\infty))}(M^*) \leq \omega_{L_\infty}(M). \tag{3.23}$$

4. Nonexpansivity of rearrangement in Orlicz spaces L_N

In this section, as a particular case of [6] (see also [13]), we consider Orlicz spaces L_N of finitely additive extended real-valued set functions defined on algebras of sets. The space L_N has been introduced in [6] in the same way as Dunford and Schwartz [9, page 112] define the space of integrable functions and the integral for integrable functions, and generalize the Orlicz spaces of σ -additive measures defined on σ -algebras of sets.

As in the previous sections, Ω is a nonempty set and \mathcal{A} is an algebra in $\mathcal{P}(\Omega)$. Let $\nu : \mathcal{A} \rightarrow [0, +\infty]$ be a finitely additive set function. Throughout we assume that each simple function $s \in S(\Omega, \mathcal{A})$ is ν -integrable, that is, $s = \sum_{i=1}^n a_i \chi_{A_i}$ with $a_i \in \mathbb{R}$, $A_i \in \mathcal{A}$ and $a_i = 0$ if $\nu(A_i) = \infty$ (with $0 \cdot \infty = 0$). Denote by $(L_1(\Omega, \mathcal{A}, \nu), \|\cdot\|_1)$ the Lebesgue space defined in [9], then $\|f\|_1 = \int_\Omega |f| d\nu$ is a Riesz pseudonorm in the sense of [14]. Let $\eta = \nu^*$ and $N : [0, +\infty) \rightarrow [0, +\infty)$ be a continuous, strictly increasing function such that $N(0) = 0$ and $N(s + t) \leq k(N(s) + N(t))$ ($k \in \mathbb{N}$) for all $s, t \geq 0$. The latter condition holds if and only if N satisfies the Δ_2 -condition, that is, there is a constant $c \in [0, +\infty[$ with $N(2t) \leq cN(t)$ for all $t \geq 0$ (see [6, page 90]).

Then, for $s \in S(\Omega, \mathcal{A})$, $\|s\|_N$ is defined by $\|s\|_N = \|N \circ |s|\|_1$, and the space E_N is defined as follows.

Definition 4.1 (see [6, page 92]). The space $L_N := L_N(\Omega, \mathcal{A}, \eta)$ is the space of all functions $f \in L_0$, for which there is a $\|\cdot\|_N$ -Cauchy sequence (s_n) in $S(\Omega, \mathcal{A})$ converging to f with respect to $\|\cdot\|_0$, and $\|f\|_N = \lim_n \|s_n\|_N$, the sequence (s_n) is said to *determine* f .

PROPOSITION 4.2 (see [6, Proposition 2.6 (c)]). *If (s_n) is a sequence in $S(\Omega, \mathcal{A})$ determining $f \in L_N$, then (s_n) converges to f with respect to $\|\cdot\|_N$.*

We will call *convergence in N -mean* the convergence with respect to $\|\cdot\|_N$.

PROPOSITION 4.3 (see [6, Proposition 2.10 (b)]). *For all $f \in L_N$, $\|f\|_N = \|N \circ |f|\|_1$.*

In the following if $\Omega = [0, +\infty)$, \mathcal{A} is the σ -algebra of all Lebesgue measurable subsets of $[0, +\infty)$ and $\eta = \mu^*$, we will write $L_N([0, +\infty))$ instead of L_N . For $f \in L_N([0, +\infty))$, we denote $\|f\|_N$ by $|f|_N$.

In order to consider rearrangements of functions of L_N to any function $s = \sum_{i=1}^n a_i \chi_{A_i}$ in $S(\Omega, \mathcal{A})$, we associate the simple function $\bar{s}: [0, +\infty) \rightarrow \mathbb{R}$ defined by

$$\bar{s} = \sum_{i=1}^n a_i \chi_{[\sum_{i=1}^{n-1} \nu(A_i), \sum_{i=1}^n \nu(A_i)]}. \quad (4.1)$$

We immediately find $\|s\|_N = |\bar{s}|_N$ and $s^* = (\bar{s})^*$.

LEMMA 4.4. *Let $s \in S(\Omega, \mathcal{A})$. Then $\|s\|_N = |s^*|_N$.*

Proof. An easy computation shows that $\int_{[0, +\infty)} N(|\bar{s}(t)|) d\mu = \int_{[0, +\infty)} N((\bar{s})^*(t)) d\mu$. Therefore, we obtain

$$\|s\|_N = \int_{[0, +\infty)} N(|\bar{s}(t)|) d\mu = \int_{[0, +\infty)} N((\bar{s})^*(t)) d\mu = \int_{[0, +\infty)} N(s^*(t)) d\mu = |s^*|_N. \quad (4.2)$$

□

LEMMA 4.5. *Let $s_1, s_2 \in S(\Omega, \mathcal{A})$. Then $|s_1^* - s_2^*|_N \leq \|s_1 - s_2\|_N$.*

Proof. By [3, (6), page 24] we have

$$\int_{[0, +\infty)} N(|(\bar{s}_1)^*(t) - (\bar{s}_2)^*(t)|) d\mu \leq \int_{[0, +\infty)} N(||\bar{s}_1(t)| - |\bar{s}_2(t)||) d\mu. \quad (4.3)$$

Since

$$\int_{[0, +\infty)} N(||\bar{s}_1(t)| - |\bar{s}_2(t)||) d\mu = \int_{\Omega} N(||s_1| - |s_2||) d\nu, \quad (4.4)$$

we get

$$|s_1^* - s_2^*|_N \leq \int_{\Omega} N(||s_1| - |s_2||) d\nu \leq \|s_1 - s_2\|_N. \quad (4.5)$$

□

LEMMA 4.6. *Let (s_n) be a sequence in $S(\Omega, \mathcal{A})$ such that $\|s_n - f\|_N \rightarrow 0$. Then*

$$|s_n^* - f^*|_N \rightarrow 0, \|f\|_N = |f^*|_N. \quad (4.6)$$

Proof. Since $\|s_n - f\|_N \rightarrow 0$ by [6, Theorem 2.7], we have $\|s_n - f\|_0 \rightarrow 0$. Then by Theorem 2.14 it follows that $|s_n^* - f|_0 \rightarrow 0$, and so we can choose a subsequence $(s_{n_k}^*)$ of (s_n^*) which converges to f^* μ -a.e. On the other hand by Lemma 4.4 since (s_n) is a $\|\cdot\|_N$ -Cauchy sequence we have that (s_n^*) is a $|\cdot|_N$ -Cauchy. Then there is a function $g \in L_N([0, +\infty))$ such that $|s_n^* - g|_N \rightarrow 0$. Therefore $|s_{n_k}^* - g|_0 \rightarrow 0$ and so we can find a subsequence $(s_{n_l}^*)$ of $(s_{n_k}^*)$ which converges to g μ -a.e. Then $f^* = g$ μ -a.e. and $|s_{n_l}^* - f^*|_N \rightarrow 0$. Finally

$$\begin{aligned} \left| \|f\|_N - |f^*|_N \right| &\leq \left| \|f\|_N - \|s_n\|_N \right| + \left| \|s_n\|_N - |s_n^*|_N \right| + \left| |s_n^*|_N - |f^*|_N \right| \\ &\leq \|s_n - f\|_N + |s_n^* - f^*|_N. \end{aligned} \tag{4.7}$$

Hence $\|f\|_N = |f^*|_N$ and this proves the lemma. □

We omit the proof of nonexpansivity of rearrangement on L_N , which is analogous to that of Theorem 3.6, when we use the above lemma.

THEOREM 4.7. *Let $f, g \in L_N$. Then $|f^* - g^*|_N \leq \|f - g\|_N$.*

COROLLARY 4.8. *Let M be a bounded set in L_N . Then*

$$\gamma_{L_N([0, +\infty))}(M^*) \leq \gamma_{L_N}(M). \tag{4.8}$$

Now let Ω be an open bounded subset of the n -dimensional Euclidean space \mathbb{R}^n (with norm $\|\cdot\|_n$), and let \mathcal{A} be the σ -algebra of all Lebesgue measurable subsets of Ω and $\eta = \mu^*$. Now we assume that Φ is a Young function and we consider the space E_Φ of finite elements of the Orlicz space L_Φ generated by Φ . In this situation, we introduce a parameter ω_{E_Φ} to estimate the Hausdorff measure of noncompactness.

Recall that Φ is a *Young function* if $\Phi(t) = \int_0^t \varphi(s) ds$ ($t \geq 0$), where $\varphi : [0, +\infty) \rightarrow [0, +\infty)$ is such that

- (i) $\varphi(0) = 0$;
- (ii) $\varphi(s) > 0, s > 0$;
- (iii) φ is right continuous at any point $s \geq 0$;
- (iv) φ is nondecreasing on $[0, +\infty)$;
- (v) $\lim_{s \rightarrow +\infty} \varphi(s) = +\infty$.

In particular, Φ is continuous, nonnegative, strictly increasing, convex on $[0, +\infty)$ and $\Phi(0) = 0$.

By $L_\Phi(\Omega)$ we denote the Orlicz space generated by Φ , that is,

$$L_\Phi(\Omega) = \left\{ f \in L_0 : \lim_{\lambda \rightarrow 0^+} \|\Phi \circ (\lambda|f|)\|_1 = 0 \right\}. \tag{4.9}$$

We equip $L_\Phi(\Omega)$ with the Luxemburg norm

$$\|f\|_\Phi = \inf \left\{ k > 0 : \left\| \Phi \circ \left(\frac{|f|}{k} \right) \right\|_1 \leq 1 \right\}. \tag{4.10}$$

By $E_\Phi(\Omega)$ we denote the space of finite elements, that is,

$$E_\Phi(\Omega) = \{ f \in L_0 : \|\Phi \circ (\lambda|f|)\|_1 < +\infty, \text{ for any } \lambda > 0 \}. \tag{4.11}$$

The space $E_\Phi(\Omega)$ is a closed subspace of $L_\Phi(\Omega)$ and $E_\Phi(\Omega) = L_\Phi(\Omega)$ if the Δ_2 -condition holds. For details on Orlicz spaces see [15, 16].

We recall that the convergence with respect to the Luxemburg norm $||| \cdot |||_\Phi$ implies Φ -mean convergence, for $\Phi \in \Delta_2$ the two types of convergence are equivalent.

For $r > 0$, $x \in \mathbb{R}^n$, and $f \in L_\Phi(\Omega)$ let us put $f(y) = 0$ if $y \notin \Omega$. The so called *Steklow function* $S_r(f)$ corresponding to f is defined as follows:

$$S_r(f)(x) = \frac{1}{\mu(B(x,r))} \int_{B(x,r)} f(y) d\mu = \frac{1}{\mu(B(x,r))} \int_{\|y\|_n < r} f(x+y) d\mu. \quad (4.12)$$

$S_r(f)$ is continuous on \mathbb{R}^n , has compact support and $|||S_r(f)|||_\Phi \leq |||f|||_\Phi$ (cfr., [16, Theorem 9.10]).

THEOREM 4.9 (see [15, (ii) page 173]). *Let M be a bounded subset of $L_\Phi(\Omega)$. Set $M_r = \{S_r(f) : f \in M\}$. Then*

- (1) $M_r \subset C_o^\infty(\mathbb{R}^n)$;
- (2) M_r is relatively compact in $C(\overline{\Omega})$ with respect to $\| \cdot \|_\infty$.

Now for any bounded subset M of $E_\Phi(\Omega)$, generalizing an analogous parameter defined in the case of Lebesgue spaces $L_p[0, 1]$, we put

$$\omega_{E_\Phi}(M) = \limsup_{\delta \rightarrow 0} \max_{f \in M} \max_{0 < r \leq \delta} |||f - S_r(f)|||_\Phi. \quad (4.13)$$

The following theorem gives an estimate of the Hausdorff measure of noncompactness γ_{E_Φ} by means of the parameter ω_{E_Φ} . We observe that the theorem is an extension of the compactness criterion given in [15, Theorem 3.14.6], which is the analogous in $E_\Phi(\Omega)$ of the Kolmogorov compactness criterion in the Lebesgue spaces $L_p[0, 1]$.

THEOREM 4.10. *Let M be a bounded set of $E_\Phi(\Omega)$. Then*

$$\frac{1}{2} \omega_{E_\Phi}(M) \leq \gamma_{E_\Phi}(M) \leq \omega_{E_\Phi}(M). \quad (4.14)$$

Proof. Let $\alpha > \omega_{E_\Phi}(M)$. For some $0 < r \leq \delta$ we have that $|||f - S_r(f)|||_\Phi \leq \alpha$ for all $f \in M$. Since M_r is compact in $C(\overline{\Omega})$ with respect to $\| \cdot \|_\infty$, for all $\varepsilon > 0$ we can choose an ε -net $\{S_r(f_1), S_r(f_2), \dots, S_r(f_n)\}$ for M_r in M_r . Then for any $f \in M$ there exists $i \in \{1, \dots, n\}$ such that $|S_r(f)(t) - S_r(f_i)(t)| \leq \varepsilon$ for all $t \in \overline{\Omega}$, so that $|||S_r(f) - S_r(f_i)|||_\Phi \leq \varepsilon |||\chi_\Omega|||_\Phi$. Hence

$$|||f - S_r(f_i)|||_\Phi \leq |||f - S_r(f)|||_\Phi + |||S_r(f) - S_r(f_i)|||_\Phi \leq \alpha + \varepsilon |||\chi_\Omega|||_\Phi \quad (4.15)$$

and consequently $\gamma_{E_\Phi}(M) \leq \omega_{E_\Phi}(M)$.

We now prove the left inequality. Let $\alpha > \gamma_{E_\Phi}(M)$. Fix an α -net $\{f_1, f_2, \dots, f_n\}$ for M in E_Φ . Since $M \subset E_\Phi$ we can assume that the functions f_i ($i = 1, 2, \dots, n$) are in $C(\overline{\Omega})$. By the uniform continuity of each f_i on $\overline{\Omega}$, there is some $\delta > 0$ such that $|f_i(t) - f_i(x)| \leq \varepsilon$ holds for each $i \in \{1, \dots, n\}$ whenever $t, x \in \overline{\Omega}$ satisfy $\|x - t\|_n < \delta$. Then, if $0 < r < \delta$ we

obtain $|f_i(t) - S_r(f_i)(t)| \leq \varepsilon$ for all $t \in \overline{\Omega}$. The latter inequality implies $\|f_i - S_r(f_i)\|_\Phi \leq \varepsilon \|\chi_\Omega\|_\Phi$. Moreover $\|S_r(f) - S_r(f_i)\|_\Phi = \|S_r(f - f_i)\|_\Phi \leq \|f - f_i\|_\Phi$. Therefore

$$\begin{aligned} \|f - S_r(f)\|_\Phi &\leq \|f - f_i\|_\Phi + \|f_i - S_r(f_i)\|_\Phi + \|S_r(f_i) - S_r(f)\|_\Phi \\ &\leq 2\|f - f_i\|_\Phi + \|f_i - S_r(f_i)\|_\Phi \leq 2\alpha + \varepsilon \|\chi_\Omega\|_\Phi \end{aligned} \quad (4.16)$$

holds for all $f \in M$ and $0 < r < \delta$. Hence $\omega_{E_\Phi}(M) \leq 2\gamma_{E_\Phi}(M)$. \square

From the last result and Corollary 4.8 we get the following.

COROLLARY 4.11. *Assume that the Young function Φ satisfies the Δ_2 -condition, and let M be a bounded subset of $L_\Phi(\Omega)$. Then*

$$\omega_{L_\Phi[0,+\infty)}(M^*) \leq 2\omega_{L_\Phi}(M). \quad (4.17)$$

Remark 4.12. We observe that in the Lebesgue space $L_p[0, 1]$ ($1 \leq p < \infty$)

$$\omega_p(f^*; \delta) \leq 2\omega_p(f; \delta) \quad (4.18)$$

for $0 \leq \delta \leq 1/2$, where $\omega_p(f; \delta) = \sup_{0 \leq h \leq \delta} (\int_{[0, 1-h]} |f(x) - f(x+h)|^p d\mu)^{1/p}$ is the modulus of continuity of a given function $f \in L_p[0, 1]$ (see [5, Theorem 3.1]).

Acknowledgment

This work was supported by MIUR of Italy.

References

- [1] G. H. Hardy, J. E. Littlewood, and G. Polya, *Inequalities*, Cambridge University Press, Cambridge, UK, 1934.
- [2] C. Bennett and R. Sharpley, *Interpolation of Operators*, vol. 129 of *Pure and Applied Mathematics*, Academic Press, Boston, Mass, USA, 1988.
- [3] G. Chiti, "Rearrangements of functions and convergence in Orlicz spaces," *Applicable Analysis*, vol. 9, no. 1, pp. 23–27, 1979.
- [4] M. G. Crandall and L. Tartar, "Some relations between nonexpansive and order preserving mappings," *Proceedings of the American Mathematical Society*, vol. 78, no. 3, pp. 385–390, 1980.
- [5] V. I. Kolyada, "Rearrangements of functions, and embedding theorems," *Russian Mathematical Surveys*, vol. 44, no. 5, pp. 73–117, 1989.
- [6] P. de Lucia and H. Weber, "Completeness of function spaces," *Ricerche di Matematica*, vol. 39, no. 1, pp. 81–97, 1990.
- [7] G. Trombetta and H. Weber, "The Hausdorff measure of noncompactness for balls of F -normed linear spaces and for subsets of L_0 ," *Bollettino dell'Unione Matematica Italiana. Serie VI. C.*, vol. 5, no. 1, pp. 213–232, 1986.
- [8] J. Banaś and K. Goebel, *Measures of Noncompactness in Banach Spaces*, vol. 60 of *Lecture Notes in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 1980.
- [9] N. Dunford and J. T. Schwartz, *Linear Operators—Part I*, John Wiley & Sons, New York, NY, USA, 1958.
- [10] K. P. S. Bhaskara Rao and M. Bhaskara Rao, *Theory of Charges*, vol. 109 of *Pure and Applied Mathematics*, Academic Press, New York, NY, USA, 1983.
- [11] H. Jarchow, *Locally Convex Spaces*, B. G. Teubner, Stuttgart, Germany, 1981.

- [12] A. Avallone and G. Trombetta, “Measures of noncompactness in the space L_0 and a generalization of the Arzelà–Ascoli theorem,” *Bollettino dell’Unione Matematica Italiana. Serie VII. B*, vol. 5, no. 3, pp. 573–587, 1991.
- [13] H. Weber, “Generalized Orlicz spaces. Locally solid group topologies,” *Mathematische Nachrichten*, vol. 145, pp. 201–215, 1990.
- [14] C. D. Aliprantis and O. Burkinshaw, *Locally Solid Riesz Spaces*, Academic Press, New York, NY, USA, 1978.
- [15] A. Kufner, O. John, and S. Fučík, *Function Spaces*, Noordhoff, Leyden, The Netherlands, 1977.
- [16] J. Musielak, *Orlicz Spaces and Modular Spaces*, vol. 1034 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1983.

D. Caponetti: Department of Mathematics, University of Palermo, 90123 Palermo, Italy
Email address: d.caponetti@math.unipa.it

A. Trombetta: Department of Mathematics, University of Calabria, 87036 Rende (CS), Italy
Email address: aletromb@unical.it

G. Trombetta: Department of Mathematics, University of Calabria, 87036 Rende (CS), Italy
Email address: trombetta@unical.it