

Research Article

On Stability of a Functional Equation Connected with the Reynolds Operator

Adam Najdecki

Received 18 July 2006; Revised 30 November 2006; Accepted 3 December 2006

Recommended by Saburou Saitoh

Let (X, \circ) be an Abelian semigroup, $g : X \rightarrow X$, and let \mathbb{K} be either \mathbb{R} or \mathbb{C} . We prove superstability of the functional equation $f(x \circ g(y)) = f(x)f(y)$ in the class of functions $f : X \rightarrow \mathbb{K}$. We also show some stability results of the equation in the class of functions $f : X \rightarrow \mathbb{K}^n$.

Copyright © 2007 Adam Najdecki. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Throughout this paper n is a positive integer, (X, \circ) is a commutative semigroup, \mathbb{K} is either the field of reals \mathbb{R} or the field of complex numbers \mathbb{C} , and $g : X \rightarrow X$ is an arbitrary function. We study stability of the functional equation

$$f(x \circ g(y)) = f(x)f(y) \quad \text{for } x, y \in X, \quad (1)$$

in the class of functions $f : X \rightarrow \mathbb{K}^n$, where $(a_1, a_2, \dots, a_n)(b_1, b_2, \dots, b_n) = (a_1b_2, a_2b_2, \dots, a_nb_n)$ for $(a_1, a_2, \dots, a_n), (b_1, b_2, \dots, b_n) \in \mathbb{K}^n$. (For details concerning the problem of stability of functional equations we refer to, e.g., [1].)

Particular cases of (1) are the well-known multiplicative Cauchy equation $f(xy) = f(x)f(y)$, exponential equation $f(x + y) = f(x)f(y)$ (see, e.g., [2]) and the equation

$$f(xf(y)) = f(x)f(y). \quad (2)$$

The origin of (2) is in the averaging theory applied to turbulent fluid motion. This equation is connected with some linear operators, that is, the Reynolds operator (see [3] and [4]), the averaging operator, the multiplicatively symmetric operator (see [2]).

Ger and Šemrl in [5] (cf. [6], [7]) considered the problem of stability for the exponential equation in the class of functions mapping X into a semisimple complex commutative

Banach algebra \mathcal{A} . They have shown that if a mapping $f : X \rightarrow \mathcal{A}$ satisfies

$$\|f(x \circ y) - f(x)f(y)\| \leq \epsilon \tag{3}$$

with some $\epsilon > 0$, then there exist a commutative C^* -algebra \mathcal{B} and a continuous monomorphism Λ of \mathcal{A} into \mathcal{B} such that \mathcal{B} is represented as a direct sum $\mathcal{B} = I \oplus J$ where I and J are closed ideals and $P\Lambda f$ is exponential, and $Q\Lambda f$ is norm-bounded where P and Q are projections corresponding to the direct sum decomposition $\mathcal{B} = I \oplus J$. We present a very short and simple proof that a similar result is valid for function $F : X \rightarrow \mathbb{K}^n$ satisfying (with any norm in \mathbb{K}^n) the following more general condition:

$$\|F(x \circ g(y)) - F(x)F(y)\| \leq \epsilon \quad \text{for } x, y \in X. \tag{4}$$

Let us start with the following theorem, showing superstability of (1).

THEOREM 1. *Let $f : X \rightarrow \mathbb{K}$ be a function satisfying*

$$|f(x \circ g(y)) - f(x)f(y)| \leq \epsilon \quad \text{for } x, y \in X. \tag{5}$$

Then either f is bounded or (1) holds.

Proof. Suppose that f is unbounded. Take a sequence $(x_n : n \in \mathbb{N})$ of elements of X with $|f(x_n)| \rightarrow \infty$. Replace in (5) x by $x \circ g(x_n)$. Then for $x, y \in X$, we have

$$|f(x \circ g(x_n) \circ g(y)) - f(x \circ g(x_n))f(y)| \leq \epsilon. \tag{6}$$

Next (5) implies

$$f(x) = \lim_{n \rightarrow \infty} \frac{f(x \circ g(x_n))}{f(x_n)} \quad \text{for } x \in X. \tag{7}$$

Thus from (6) and (7), for every $x, y \in X$, we obtain

$$\begin{aligned} f(x \circ g(y)) &= \lim_{n \rightarrow \infty} \frac{f(x \circ g(y) \circ g(x_n))}{f(x_n)} \\ &= \lim_{n \rightarrow \infty} \frac{f(x \circ g(x_n) \circ g(y)) - f(x \circ g(x_n))f(y)}{f(x_n)} + \lim_{n \rightarrow \infty} \frac{f(x \circ g(x_n))}{f(x_n)} f(y) \\ &= f(x)f(y). \end{aligned} \tag{8}$$

□

Remark 2. If $f : X \rightarrow \mathbb{K}$ is a bounded function satisfying (5), then

$$|f(x)| \leq \frac{1 + \sqrt{1 + 4\epsilon}}{2} \quad \text{for } x \in X. \tag{9}$$

In fact, suppose that $f : X \rightarrow \mathbb{K}$ satisfies (5) and

$$M := \sup \{|f(x)| : x \in X\} > \frac{1 + \sqrt{1 + 4\epsilon}}{2}. \tag{10}$$

There exists a sequence $(x_n : n \in \mathbb{N})$ of elements of X such that $\lim_{n \rightarrow \infty} |f(x_n)| = M$. Then for sufficiently large $n \in \mathbb{N}$, we have

$$|f(x_n \circ g(x_n)) - f(x_n)^2| \geq ||f(x_n)|^2 - |f(x_n \circ g(x_n))|| \geq |f(x_n)|^2 - M. \quad (11)$$

Moreover

$$\lim_{n \rightarrow \infty} (|f(x_n)|^2 - M) = M^2 - M > \epsilon. \quad (12)$$

Thus $|f(x_n \circ g(x_n)) - f(x_n)^2| > \epsilon$ for some $n \in \mathbb{N}$, which contradicts (5).

THEOREM 3. *Let $F : X \rightarrow \mathbb{K}^n$, $F = (f_1, f_2, \dots, f_n)$ be a function satisfying (4). Then there exist ideals $I, J \subset \mathbb{K}^n$ such that $\mathbb{K}^n = I \oplus J$, PF is bounded, and QF satisfies (1) where $P : \mathbb{K}^n \rightarrow I$ and $Q : \mathbb{K}^n \rightarrow J$ are natural projections.*

Proof. Since every two norms in \mathbb{K}^n are equivalent, (4) implies that there is $\eta > 0$ such that

$$\sum_{i=1}^n |f_i(x \circ g(y)) - f_i(x)f_i(y)| \leq \eta \|F(x \circ g(y)) - F(x)F(y)\| \leq \eta \epsilon \quad \text{for } x, y \in X. \quad (13)$$

Let $M := \{i \in \{1, \dots, n\} : f_i \text{ is an unbounded solution of (1)}\}$ and $L := \{i \in \{1, \dots, n\} : f_i \text{ is bounded}\}$. By Theorem 1, $L \cup M = \{1, \dots, n\}$. Now it is enough to write $I = \{(a_1, \dots, a_n) \in \mathbb{K}^n : a_i = 0 \text{ for } i \in M\}$ and $J = \{(a_1, \dots, a_n) \in \mathbb{K}^n : a_i = 0 \text{ for } i \in L\}$. \square

References

- [1] D. H. Hyers, G. Isac, and Th. M. Rassias, *Stability of Functional Equations in Several Variables*, vol. 34 of *Progress in Nonlinear Differential Equations and Their Applications*, Birkhäuser Boston, Boston, Mass, USA, 1998.
- [2] J. Aczél and J. Dhombres, *Functional Equations in Several Variables*, vol. 31 of *Encyclopedia of Mathematics and Its Applications*, Cambridge University Press, Cambridge, UK, 1989.
- [3] M.-L. Dubreil-Jacotin, “Propriétés algébriques des transformations de Reynolds,” *Comptes Rendus de l’Académie des Sciences*, vol. 236, pp. 1950–1951, 1953.
- [4] Y. Matras, “Sur l’équation fonctionnelle $f[x \cdot f(y)] = f(x) \cdot f(y)$,” *Académie Royale de Belgique. Bulletin de la Classe des Sciences. 5e Série*, vol. 55, pp. 731–751, 1969.
- [5] R. Ger and P. Šemrl, “The stability of the exponential equation,” *Proceedings of the American Mathematical Society*, vol. 124, no. 3, pp. 779–787, 1996.
- [6] J. A. Baker, J. Lawrence, and F. Zorzitto, “The stability of the equation $f(x + y) = f(x)f(y)$,” *Proceedings of the American Mathematical Society*, vol. 74, no. 2, pp. 242–246, 1979.
- [7] J. A. Baker, “The stability of the cosine equation,” *Proceedings of the American Mathematical Society*, vol. 80, no. 3, pp. 411–416, 1980.

Adam Najdecki: Institute of Mathematics, University of Rzeszów, Rejtana 16A,
35-310 Rzeszów, Poland
Email address: najdecki@univ.rzeszow.pl