

Research Article

Regularity Criterion for Weak Solutions to the Navier-Stokes Equations in Terms of the Gradient of the Pressure

Jishan Fan¹ and Tohru Ozawa²

¹ Department of Applied Mathematics, Nanjing Forestry University, Nanjing 210037, China

² Department of Applied Physics, Waseda University, Tokyo 169-8555, Japan

Correspondence should be addressed to Tohru Ozawa, txozdwa@waseda.jp

Received 26 June 2008; Accepted 14 October 2008

Recommended by Michel Chipot

We prove a regularity criterion $\nabla\pi \in L^{2/3}(0, T; \text{BMO})$ for weak solutions to the Navier-Stokes equations in three-space dimensions. This improves the available result with $L^{2/3}(0, T; L^\infty)$.

Copyright © 2008 J. Fan and T. Ozawa. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

We study the regularity condition of weak solutions to the Navier-Stokes equations

$$u_t - \Delta u + (u \cdot \nabla)u + \nabla\pi = 0, \quad (1.1)$$

$$\operatorname{div} u = 0, \quad \text{in } (0, T) \times \mathbb{R}^3, \quad (1.2)$$

$$u|_{t=0} = u_0(x), \quad x \in \mathbb{R}^3. \quad (1.3)$$

Here, u is the unknown velocity vector and π is the unknown scalar pressure.

For $u_0 \in L^2(\mathbb{R}^3)$ with $\operatorname{div} u_0 = 0$ in \mathbb{R}^3 , Leray [1] constructed global weak solutions. The smoothness of Leray's weak solutions is unknown. While the existence of regular solutions is still an open problem, there are many interesting sufficient conditions which guarantee that a given weak solution is smooth. A well-known condition states that if

$$u \in L^r(0, T; L^s(\mathbb{R}^3)) \quad \text{with} \quad \frac{2}{r} + \frac{3}{s} = 1, \quad 3 \leq s \leq \infty, \quad (1.4)$$

then the solution u is actually regular [2–8]. A similar condition

$$\omega := \operatorname{curl} u \in L^r(0, T; L^s(\mathbb{R}^3)), \quad \text{with } \frac{2}{r} + \frac{3}{s} = 2, \quad \frac{3}{2} \leq s \leq \infty, \quad (1.5)$$

also implies the regularity as shown by Beirão da Veiga [9].

As regards (1.4) and (1.5) for $s = \infty$, Kozono et al. made an improvement to the following condition:

$$u \in L^2(0, T; \dot{B}_{\infty, \infty}^0(\mathbb{R}^3)), \quad (1.6)$$

or

$$\omega \in L^1(0, T; \dot{B}_{\infty, \infty}^0(\mathbb{R}^3)), \quad (1.7)$$

where $\dot{B}_{\infty, \infty}^0$ is the homogeneous Besov space. On the other hand, Chae and Lee [10] proposed another regularity criterion in terms of the pressure. They showed that if the pressure π satisfies

$$\pi \in L^r(0, T; L^s(\mathbb{R}^3)) \quad \text{with } \frac{2}{r} + \frac{3}{s} < 2, \quad \frac{3}{2} < s \leq \infty, \quad (1.8)$$

then u is smooth. Berselli and Galdi [11] have extended the range of r and s to $2/r + 3/s = 2$ and $3/2 < s \leq \infty$. When $s = \infty$, Chen and Zhang [12] (also see Fan et al. [13]) refined it to the following condition:

$$\pi \in L^1(0, T; \dot{B}_{\infty, \infty}^0(\mathbb{R}^3)). \quad (1.9)$$

Zhou [14] (see also Struwe [15]) proposed the following criterion in terms of the gradient of the pressure:

$$\nabla \pi \in L^r(0, T; L^s(\mathbb{R}^3)) \quad \text{with } \frac{2}{r} + \frac{3}{s} = 3, \quad 1 < s \leq \infty. \quad (1.10)$$

The aim of this paper is to refine (1.10) when $s = \infty$. We will use the following interpolation inequality:

$$\|u\|_{L^{2p}(\mathbb{R}^n)}^2 \leq C \|u\|_{L^p(\mathbb{R}^n)} \|u\|_{\text{BMO}}, \quad 1 \leq p < \infty, \quad (1.11)$$

which follows from the bilinear estimates

$$\|fg\|_{L^p} \leq C(\|f\|_{L^p} \|g\|_{\text{BMO}} + \|g\|_{L^p} \|f\|_{\text{BMO}}), \quad 1 \leq p < \infty, \quad (1.12)$$

due to Kozono and Taniuchi [16]. Here, BMO is the space of functions of bounded mean oscillations.

Definition 1.1. Let $u_0 \in L^2(\mathbb{R}^3)$ with $\operatorname{div} u_0 = 0$ in \mathbb{R}^3 . The function u is called a Leray weak solution of (1.1)–(1.3) in $(0, T)$ if u satisfies the following properties.

- (1) $u \in L^\infty(0, T; L^2) \cap L^2(0, T; H^1)$.
- (2) Equation (1.1) and (1.2) hold in the distributional sense, and

$$u(t) \rightharpoonup u_0 \quad \text{weakly in } L^2 \text{ as } t \rightarrow 0. \quad (1.13)$$

- (3) The energy inequality is

$$\|u(t)\|_{L^2}^2 + 2 \int_0^t \|\nabla u(s)\|_{L^2}^2 ds \leq \|u_0\|_{L^2}^2, \quad \text{for any } t \in [0, T]. \quad (1.14)$$

Our main result reads as follows.

Theorem 1.2. Let $u_0 \in (L^2 \cap L^4)(\mathbb{R}^3)$ with $\operatorname{div} u_0 = 0$ in \mathbb{R}^3 . Suppose that u is a Leray weak solution of (1.1)–(1.3) in $(0, T)$. If the gradient of the pressure satisfies the condition

$$\nabla \pi \in L^{2/3}(0, T; \text{BMO}), \quad (1.15)$$

then u is smooth in $(0, T]$.

Remark 1.3. If the interpolation inequality

$$\|u\|_{L^{2p}(\mathbb{R}^n)}^2 \leq C \|u\|_{L^p} \|u\|_{\dot{B}_{\infty, \infty}^0} \quad (1.16)$$

is true, then as in the argument below, (1.15) may be improved to the following condition:

$$\nabla \pi \in L^{2/3}(0, T; \dot{B}_{\infty, \infty}^0). \quad (1.17)$$

Remark 1.4. Inequality (1.11) plays an important role in our proof. Chen and Zhu [17] extended (1.11) to the following inequality:

$$\|u\|_{L^q} \leq C \|u\|_{L^r}^{r/q} \|u\|_{\text{BMO}}^{1-r/q}, \quad 1 \leq r < q < \infty, \quad (1.18)$$

and used (1.18) to obtain (1.12). Kozono and Wadade [18] give another proof of (1.18). Here, we give an elementary and short proof of (1.18) by (1.11).

For given $1 \leq r < q < \infty$, there exists a positive integer n and $\theta \in (0, 1)$ such that $r < q < 2^n r$ and $1/q = \theta \cdot (1/r) + (1 - \theta) \cdot (1/2^n r) = (\theta + (1 - \theta)/2^n) \cdot (1/r)$. By the Hölder inequality, we have

$$\|u\|_{L^q} \leq \|u\|_{L^r}^\theta \|u\|_{L^{2^n r}}^{1-\theta}. \quad (1.19)$$

Using (1.11) for $p = 2^{n-1}r, 2^{n-2}r, \dots, r$, n times and plugging them into (1.6), we find that

$$\begin{aligned} \|u\|_{L^q} &\leq C \|u\|_{L^r}^{\theta+(1-\theta)/2^n} \|u\|_{\text{BMO}}^{(1-\theta)(1/2+1/2^2+\dots+1/2^n)} \\ &= C \|u\|_{L^r}^{\theta+(1-\theta)/2^n} \|u\|_{\text{BMO}}^{(1-1/2^n)(1-\theta)} \\ &= C \|u\|_{L^r}^{r/q} \|u\|_{\text{BMO}}^{1-r/q}, \end{aligned} \quad (1.20)$$

which proves (1.18).

Remark 1.5. From Remark 1.4, we know that if (1.16) holds true, then we have

$$\|u\|_{L^q} \leq C \|u\|_{L^r}^{r/q} \|u\|_{\dot{B}_{\infty,\infty}^0}^{1-r/q}, \quad 1 \leq r < q < \infty. \quad (1.21)$$

2. Proof of Theorem 1.2

This section is devoted to the proof of Theorem 1.2. First, we recall the following result according to Giga [5].

Proposition 2.1 (see [5]). *Suppose $u_0 \in L^s(\mathbb{R}^3)$, $s \geq 3$; then there exists T and a unique classical solution $u \in (L^\infty \cap C)([0, T); L^s)$. Moreover, let $(0, T^*)$ be the maximal interval such that u solves (1.1)–(1.3) in $C((0, T^*); L^s)$, $s > 3$. Then, for any $t \in (0, T^*)$,*

$$\|u(t)\|_{L^s} \geq \frac{C}{(T^* - t)^{(s-3)/2s}} \quad (2.1)$$

with the constant C independent of T^* and s .

Next, we derive a priori estimates for smooth solutions of (1.1)–(1.3). To this end, multiplying (1.1) by $|u|^2 u$, integrating by parts, and using (1.2), (1.11) for $p = 2$, we see that

$$\begin{aligned} \frac{1}{4} \frac{d}{dt} \|u\|_{L^4}^4 + \int |\nabla u|^2 |u|^2 dx + \frac{1}{4} \int |\nabla |u|^2|^2 dx &= - \int \nabla \pi \cdot |u|^2 u dx \leq \|\nabla \pi\|_{L^4} \|u\|_{L^4}^3 \\ &\leq C \|\nabla \pi\|_{L^2}^{1/2} \|\nabla \pi\|_{\text{BMO}}^{1/2} \|u\|_{L^4}^3 \\ &\leq C \|u \nabla u\|_{L^2}^{1/2} \|\nabla \pi\|_{\text{BMO}}^{1/2} \|u\|_{L^4}^3 \\ &\leq \frac{1}{2} \int |u \cdot \nabla u|_{L^2}^2 + C \|\nabla \pi\|_{\text{BMO}}^{2/3} \|u\|_{L^4}^4, \end{aligned} \quad (2.2)$$

which yields

$$\|u\|_{L^4} \leq \|u_0\|_{L^4} \exp \left(C \int_0^T \|\nabla \pi\|_{\text{BMO}}^{2/3} dt \right), \quad (2.3)$$

by Gronwall's inequality. Here, we have used the estimate

$$\|\nabla\pi\|_{L^2} \leq C\|u \cdot \nabla u\|_{L^2}. \quad (2.4)$$

Now, we are in a position to complete the proof of Theorem 1.2. From Proposition 2.1, it follows that there exists $T^* > 0$ and the smooth solution v of (1.1)–(1.3) satisfies

$$v(t) \in (L^\infty \cap C)([0, T^*]; L^4), \quad v(0) = u_0. \quad (2.5)$$

Since the weak solution u satisfies the energy inequality, we may apply Serrin's uniqueness criterion [19] to conclude that

$$u \equiv v \quad \text{on } [0, T^*]. \quad (2.6)$$

Thus, it is sufficient to show that $T^* = T$. Suppose that $T^* < T$. Without loss of generality, we may assume that T^* is the maximal existence time for $v(t)$. By Proposition 2.1 again, we find that

$$\|u(t)\|_{L^4} \geq \frac{C}{(T^* - t)^{1/8}} \quad \text{for any } t \in (0, T^*). \quad (2.7)$$

On the other hand, from (2.3), we know that

$$\sup_{0 \leq t \leq T^*} \|u(t)\|_{L^4} \leq \|u_0\|_{L^4} \exp\left(C \int_0^{T^*} \|\nabla\pi\|_{\text{BMO}}^{2/3} dt\right), \quad (2.8)$$

which contradicts with (2.7). Thus, $T^* = T$.

This completes the proof.

Acknowledgment

This work is supported by NSFC (Grant no. 10301014).

References

- [1] J. Leray, "Sur le mouvement d'un liquide visqueux emplissant l'espace," *Acta Mathematica*, vol. 63, no. 1, pp. 193–248, 1934.
- [2] J. Serrin, "On the interior regularity of weak solutions of the Navier-Stokes equations," *Archive for Rational Mechanics and Analysis*, vol. 9, no. 1, pp. 187–195, 1962.
- [3] T. Ohyaama, "Interior regularity of weak solutions of the time-dependent Navier-Stokes equation," *Proceedings of the Japan Academy*, vol. 36, no. 5, pp. 273–277, 1960.
- [4] E. B. Fabes, B. F. Jones, and N. M. Riviere, "The initial value problem for the Navier-Stokes equations with data in L^p ," *Archive for Rational Mechanics and Analysis*, vol. 45, no. 3, pp. 222–240, 1972.
- [5] Y. Giga, "Solutions for semilinear parabolic equations in L^p and regularity of weak solutions of the Navier-Stokes system," *Journal of Differential Equations*, vol. 62, no. 2, pp. 186–212, 1986.
- [6] M. Struwe, "On partial regularity results for the Navier-Stokes equations," *Communications on Pure and Applied Mathematics*, vol. 41, no. 4, pp. 437–458, 1988.

- [7] H. Sohr and W. von Wahl, "On the regularity of the pressure of weak solutions of Navier-Stokes equations," *Archiv der Mathematik*, vol. 46, no. 5, pp. 428–439, 1986.
- [8] L. Iskauriaza, G. A. Serëgin, and V. Shverak, " $L^{3,\infty}$ -solutions of Navier-Stokes equations and backward uniqueness," *Russian Mathematical Surveys*, vol. 58, no. 2, pp. 211–250, 2003.
- [9] H. Beirão da Veiga, "A new regularity class for the Navier-Stokes equations in \mathbb{R}^n ," *Chinese Annals of Mathematics. Series B*, vol. 16, no. 4, pp. 407–412, 1995.
- [10] D. Chae and J. Lee, "Regularity criterion in terms of pressure for the Navier-Stokes equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 46, no. 5, pp. 727–735, 2001.
- [11] L. C. Berselli and G. P. Galdi, "Regularity criteria involving the pressure for the weak solutions to the Navier-Stokes equations," *Proceedings of the American Mathematical Society*, vol. 130, no. 12, pp. 3585–3595, 2002.
- [12] Q. Chen and Z. Zhang, "Regularity criterion via the pressure on weak solutions to the 3D Navier-Stokes equations," *Proceedings of the American Mathematical Society*, vol. 135, no. 6, pp. 1829–1837, 2007.
- [13] J. Fan, S. Jiang, and G. Ni, "On regularity criteria for the n -dimensional Navier-Stokes equations in terms of the pressure," *Journal of Differential Equations*, vol. 244, no. 11, pp. 2963–2979, 2008.
- [14] Y. Zhou, "On regularity criteria in terms of pressure for the Navier-Stokes equations in \mathbb{R}^3 ," *Proceedings of the American Mathematical Society*, vol. 134, no. 1, pp. 149–156, 2006.
- [15] M. Struwe, "On a Serrin-type regularity criterion for the Navier-Stokes equations in terms of the pressure," *Journal of Mathematical Fluid Mechanics*, vol. 9, no. 2, pp. 235–242, 2007.
- [16] H. Kozono and Y. Taniuchi, "Bilinear estimates in BMO and the Navier-Stokes equations," *Mathematische Zeitschrift*, vol. 235, no. 1, pp. 173–194, 2000.
- [17] J. Chen and X. Zhu, "A note on BMO and its application," *Journal of Mathematical Analysis and Applications*, vol. 303, no. 2, pp. 696–698, 2005.
- [18] H. Kozono and H. Wadade, "Remarks on Gagliardo-Nirenberg type inequality with critical Sobolev space and BMO ," *Mathematische Zeitschrift*, vol. 259, no. 4, pp. 935–950, 2008.
- [19] J. Serrin, "The initial value problem for the Navier-Stokes equations," in *Nonlinear Problems*, R. E. Langer, Ed., pp. 69–98, University of Wisconsin Press, Madison, Wis, USA, 1963.