

## Research Article

# Ostrowski Type Inequalities for Higher-Order Derivatives

Mingjin Wang<sup>1</sup> and Xilai Zhao<sup>2</sup>

<sup>1</sup> Department of Applied Mathematics, Jiangsu Polytechnic University, Changzhou 213164, Jiangsu, China

<sup>2</sup> Department of Mechanical and Electrical Engineering, Hebi College of Vocation and Technology, Hebi, Henan 458030, China

Correspondence should be addressed to Mingjin Wang, wang197913@126.com

Received 12 February 2009; Revised 16 May 2009; Accepted 14 July 2009

Recommended by Patricia J. Y. Wong

This paper has shown some new Ostrowski type inequalities involving higher-order derivatives. The results generalized the Ostrowski type inequalities. Applications of the inequalities are also given.

Copyright © 2009 M. Wang and X. Zhao. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Main Result and Introduction

The following inequality is well known in literature as Ostrowski's integral inequality.

Let  $f : [a, b] \rightarrow R$  be continuous on  $[a, b]$  and differentiable on  $(a, b)$  whose derivative  $f' : (a, b) \rightarrow R$  is bounded on  $(a, b)$ , that is,  $\|f'\|_\infty = \sup_{t \in (a, b)} |f'(t)| < \infty$ . Then

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left\{ \frac{1}{4} + \frac{(x - (a+b)/2)^2}{(b-a)^2} \right\} (b-a) \|f'\|_\infty. \quad (1.1)$$

Moreover the constant  $1/4$  is the best possible. Because Ostrowski's integral inequality is useful in some fields, many generalizations, extensions, and variants of this inequality have appeared in the literature; see [1–9] and the references given therein. The main aim of this paper is to establish some new Ostrowski type inequalities involving higher-order derivatives. The analysis used in the proof is elementary. The main result of this paper is the following inequality.

**Theorem 1.1.** *Suppose*

- (1)  $f : [a, b] \rightarrow \mathbb{R}$  to be continuous on  $[a, b]$ ;
- (2)  $f : [a, b] \rightarrow \mathbb{R}$  to be  $n$ th order differentiable on  $(a, b)$  whose  $n$ th order derivative  $f^{(n)} : (a, b) \rightarrow \mathbb{R}$  is bounded on  $(a, b)$ , that is,  $\|f^{(n)}\|_\infty = \sup_{t \in (a, b)} |f^{(n)}(t)| < \infty$ ;
- (3) there exists  $x_0 \in (a, b)$  such that  $f^{(k)}(x_0) = 0$ ,  $k = 1, 2, \dots, n-1$ .

Then for any  $x \in [a, b]$ , we have

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{\|f^{(n)}\|_\infty}{n!} \left\{ \left( \left| x - \frac{a+b}{2} \right| + \frac{b-a}{2} \right)^n + \frac{(b-a)^n}{n+1} \right\}. \quad (1.2)$$

As applications of the inequality (1.2), we give more Ostrowski type inequalities.

## 2. The Proof of Theorem 1.1

In this section, we use the Taylor expansion to prove Theorem 1.1. Before the proof, we need the following lemmas.

**Lemma 2.1.** *Suppose  $a \leq x \leq b$  and  $a < t < b$ , then we have*

$$(x-t)^2 \leq \left( \left| x - \frac{a+b}{2} \right| + \frac{b-a}{2} \right)^2 \quad (2.1)$$

*Proof.* When  $a \leq x \leq (a+b)/2$ , then

$$(x-t)^2 \leq (x-b)^2 = \left( \left| x - \frac{a+b}{2} \right| + \frac{b-a}{2} \right)^2. \quad (2.2)$$

When  $(a+b)/2 \leq x \leq b$ , then

$$(x-t)^2 \leq (x-a)^2 = \left( \left| x - \frac{a+b}{2} \right| + \frac{b-a}{2} \right)^2. \quad (2.3)$$

From (2.2) and (2.3), we know that (2.1) holds.  $\square$

**Lemma 2.2.** *Suppose  $a \leq t \leq b$ , then for  $n \geq 1$  we have*

$$(b-t)^n + (t-a)^n \leq (b-a)^n. \quad (2.4)$$

*Proof.* It is obvious that (2.4) is true for  $n = 1$ . When  $n \geq 2$ , let

$$g(t) = (b-t)^n + (t-a)^n, \quad a \leq t \leq b, \quad (2.5)$$

then

$$g'(t) = n[(t-a)^{n-1} - (b-t)^{n-1}]. \quad (2.6)$$

The only real root of  $g'(t) = 0$  is  $t = (a+b)/2$ . Notice

$$g\left(\frac{a+b}{2}\right) = \frac{(b-a)^n}{2^{n-1}} \leq (b-a)^n = g(a) = g(b). \quad (2.7)$$

Therefore we get the inequality (2.4). □

Now, we give the proof of Theorem 1.1.

*Proof.* Using the Taylor expansion of  $f(x)$  at  $x_0$  gives

$$f(x) = f(x_0) + \frac{f^{(n)}(x_0 + \theta(x-x_0))}{n!}(x-x_0)^n, \quad 0 \leq \theta \leq 1. \quad (2.8)$$

Taking the integral on both sides of (2.8) with respect to variable  $x$  over  $[a, b]$ , we have

$$\frac{1}{b-a} \int_a^b f(x) dx = f(x_0) + \frac{1}{n!(b-a)} \int_a^b f^{(n)}(x_0 + \theta(x-x_0))(x-x_0)^n dx, \quad (2.9)$$

where the parameter  $\theta$  is not a constant but depends on  $x$ . From (2.8) and (2.9) one gets

$$\begin{aligned} f(x) - \frac{1}{b-a} \int_a^b f(x) dx &= \frac{f^{(n)}(x_0 + \theta(x-x_0))}{n!}(x-x_0)^n \\ &\quad - \frac{1}{n!(b-a)} \int_a^b f^{(n)}(x_0 + \theta(x-x_0))(x-x_0)^n dx. \end{aligned} \quad (2.10)$$

So we have

$$\begin{aligned}
 & \left| f(x) - \frac{1}{b-a} \int_a^b f(x) dx \right| \\
 &= \left| \frac{f^{(n)}(x_0 + \theta(x-x_0))}{n!} (x-x_0)^n \right. \\
 &\quad \left. - \frac{1}{n!(b-a)} \int_a^b f^{(n)}(x_0 + \theta(x-x_0)) (x-x_0)^n dx \right| \\
 &\leq \left| \frac{f^{(n)}(x_0 + \theta(x-x_0))}{n!} (x-x_0)^n \right| \\
 &\quad + \left| \frac{1}{n!(b-a)} \int_a^b f^{(n)}(x_0 + \theta(x-x_0)) (x-x_0)^n dx \right| \\
 &\leq \frac{\|f^{(n)}\|_\infty}{n!} \left\{ \left( |x-x_0|^n + \frac{1}{b-a} \right) \int_a^b |x-x_0|^n dx \right\} \\
 &= \frac{\|f^{(n)}\|_\infty}{n!} \left\{ \left( |x-x_0|^n + \frac{1}{(n+1)(b-a)} \right) [(b-x_0)^{n+1} + (x_0-a)^{n+1}] \right\}.
 \end{aligned} \tag{2.11}$$

Using Lemmas 2.1 and 2.2 gives (1.2). Thus, we complete the proof.  $\square$

### 3. Some Applications

In this section, we show some applications of the inequality (1.2). In fact, we can use (1.2) to derive some new Ostrowski type inequalities.

**Theorem 3.1.** *Suppose*

- (1)  $f : [a, b] \rightarrow R$  to be continuous on  $[a, b]$ ;
- (2)  $f : [a, b] \rightarrow R$  to be second order differentiable on  $(a, b)$  whose second derivative  $f'' : (a, b) \rightarrow R$  is bounded on  $(a, b)$ , that is,  $\|f''\|_\infty = \sup_{t \in (a, b)} |f''(t)| < \infty$ ;
- (3)  $f(a) = f(b)$ .

Then for any  $x \in [a, b]$ , we have

$$\begin{aligned}
 & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\
 &\leq \frac{1}{2} \|f''\|_\infty (b-a)^2 \left\{ \frac{(x - (a+b)/2)^2}{(b-a)^2} + \frac{|x - (a+b)/2|}{b-a} + \frac{7}{12} \right\}.
 \end{aligned} \tag{3.1}$$

*Proof.* From Rolle's mean value theorem, we know that there exists  $x_0 \in (a, b)$  such that  $f'(x_0) = 0$ . Let  $n = 2$  in the inequality (1.2), then we have (3.1).  $\square$

**Corollary 3.2.** *With the assumptions in Theorem 3.1, we have*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{2} \|f''\|_{\infty} (b-a)^2 \left\{ \frac{(x - (a+b)/2)^2}{(b-a)^2} + \frac{13}{12} \right\}. \quad (3.2)$$

*Proof.* For any  $x \in [a, b]$ , we have

$$\left| x - \frac{a+b}{2} \right| \leq \frac{b-a}{2}. \quad (3.3)$$

Consequently, (3.1) gives

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{2} \|f''\|_{\infty} (b-a)^2 \left\{ \frac{(x - (a+b)/2)^2}{(b-a)^2} + \frac{|x - (a+b)/2|}{b-a} + \frac{7}{12} \right\} \\ & \leq \frac{1}{2} \|f''\|_{\infty} (b-a)^2 \left\{ \frac{(x - (a+b)/2)^2}{(b-a)^2} + \frac{b-a}{2(b-a)} + \frac{7}{12} \right\} \\ & = \frac{1}{2} \|f''\|_{\infty} (b-a)^2 \left\{ \frac{(x - (a+b)/2)^2}{(b-a)^2} + \frac{13}{12} \right\}. \end{aligned} \quad (3.4)$$

□

**Corollary 3.3.** *With the assumptions in Theorem 3.1, we have*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{2} \|f''\|_{\infty} (b-a)^2 \left\{ \frac{|x - (a+b)/2|}{b-a} + \frac{5}{6} \right\}. \quad (3.5)$$

*Proof.* For any  $x \in [a, b]$ , we have

$$\left( x - \frac{a+b}{2} \right)^2 \leq \frac{(b-a)^2}{4}. \quad (3.6)$$

Substituting (3.6) into (3.1) gives (3.5). □

**Theorem 3.4.** *Suppose*

- (1)  $f : [a, b] \rightarrow R$  to be continuous on  $[a, b]$ ;
- (2)  $f : [a, b] \rightarrow R$  to be  $n$ th order differentiable on  $(a, b)$  whose  $n$ th order derivative  $f^{(n)} : (a, b) \rightarrow R$  is bounded on  $(a, b)$ , that is,  $\|f^{(n)}\|_{\infty} = \sup_{t \in (a, b)} |f^{(n)}(t)| < \infty$ .

Then for any  $x_0 \in (a, b)$  and  $x \in [a, b]$ , we have

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{\|f^{(n)}\|_\infty}{n!} \left\{ \left( \left| x - \frac{a+b}{2} \right| + \frac{b-a}{2} \right)^n + \frac{(b-a)^n}{n+1} \right\} \\ & \quad + (b-a) \left( \frac{1}{4} + \frac{(x - (a+b)/2)^2}{(b-a)^2} \right) \sum_{k=1}^{n-1} \frac{|f^{(k)}(x_0)|}{(k-1)!} \left( \left| x_0 - \frac{a+b}{2} \right| + \frac{b-a}{2} \right)^{k-1}. \end{aligned} \quad (3.7)$$

*Proof.* Let

$$p(x) = \sum_{k=1}^{n-1} \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k, \quad (3.8)$$

$$F(x) = f(x) - p(x).$$

Then we have

$$F^{(k)}(x) = 0, \quad k = 1, 2, \dots, n-1, \quad (3.9)$$

$$F^{(n)}(x) = f^{(n)}(x).$$

Using inequality (1.2) to  $F(x)$  gives

$$\begin{aligned} & \left| F(x) - \frac{1}{b-a} \int_a^b F(t) dt \right| \\ & = \left| f(x) - p(x) - \frac{1}{b-a} \int_a^b (f(t) - p(t)) dt \right| \\ & \leq \frac{\|f^{(n)}\|_\infty}{n!} \left\{ \left( \left| x - \frac{a+b}{2} \right| + \frac{b-a}{2} \right)^n + \frac{(b-a)^n}{n+1} \right\}. \end{aligned} \quad (3.10)$$

Since

$$\begin{aligned} & \left| f(x) - p(x) - \frac{1}{b-a} \int_a^b (f(t) - p(t)) dt \right| \\ & = \left| \left( f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right) - \left( p(x) - \frac{1}{b-a} \int_a^b p(t) dt \right) \right| \\ & \geq \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| - \left| p(x) - \frac{1}{b-a} \int_a^b p(t) dt \right|, \end{aligned} \quad (3.11)$$

we have

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{\|f^{(n)}\|_\infty}{n!} \left\{ \left( \left| x - \frac{a+b}{2} \right| + \frac{b-a}{2} \right)^n + \frac{(b-a)^n}{n+1} \right\} \\ & \quad + \left| p(x) - \frac{1}{b-a} \int_a^b p(t) dt \right|. \end{aligned} \quad (3.12)$$

Using Ostrowski's integral inequality (1.1) one gets

$$\left| p(x) - \frac{1}{b-a} \int_a^b p(t) dt \right| \leq \left\{ \frac{1}{4} + \frac{(x - (a+b)/2)^2}{(b-a)^2} \right\} (b-a) \|p'\|_\infty. \quad (3.13)$$

Notice

$$\begin{aligned} \|p'\|_\infty &= \sup_{x \in (a,b)} |p'(x)| \\ &= \sup_{x \in (a,b)} \left| \sum_{k=1}^{n-1} \frac{f^{(k)}(x_0)}{(k-1)!} (x-x_0)^{k-1} \right| \\ &\leq \sum_{k=1}^{n-1} \frac{|f^{(k)}(x_0)|}{(k-1)!} \left( \left| x_0 - \frac{a+b}{2} \right| + \frac{b-a}{2} \right)^{k-1}. \end{aligned} \quad (3.14)$$

Substituting (3.13) and (3.14) into (3.12) gives (3.7).  $\square$

It is easy to see that (3.7) is the generalization of (1.2). If we let  $x_0 = (a+b)/2$  in (3.7) and use (3.6), we get the following inequality.

**Corollary 3.5.** *With the assumptions in Theorem 3.4, we have*

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{\|f^{(n)}\|_\infty}{n!} \left\{ \left( \left| x - \frac{a+b}{2} \right| + \frac{b-a}{2} \right)^n + \frac{(b-a)^n}{n+1} \right\} \\ & \quad + \sum_{k=1}^{n-1} \frac{|f^{(k)}((a+b)/2)|}{(k-1)!} \left( \frac{b-a}{2} \right)^k. \end{aligned} \quad (3.15)$$

## Acknowledgments

The author would like to express deep appreciation to the referees for the helpful suggestions. Mingjin Wang was supported by STF of Jiangsu Polytechnic University.

## References

- [1] A. Ostrowski, "Über die absolutabweichung einer differentienbaren Funktionen von ihren integralmittelwert," *Commentarii Mathematici Helvetici*, vol. 10, pp. 226–227, 1938.
- [2] G. A. Anastassiou, "Multivariate Ostrowski type inequalities," *Acta Mathematica Hungarica*, vol. 76, no. 4, pp. 267–278, 1997.
- [3] N. S. Barnett and S. S. Dragomir, "An Ostrowski type inequality for double integrals and applications for cubature formulae," *RGMI Research Report Collection*, vol. 1, no. 1, pp. 13–22, 1998.
- [4] S. S. Dragomir, N. S. Barnett, and P. Cerone, "An n-dimensional version of Ostrowski's inequality for mappings of the Hölder type," *RGMI Research Report Collection*, vol. 2, no. 2, pp. 169–180, 1999.
- [5] S. S. Dragomir, R. P. Agarwal, and P. Cerone, "On Simpson's inequality and applications," *Journal of Inequalities and Applications*, vol. 5, no. 6, pp. 533–579, 2000.
- [6] S. S. Dragomir, "Ostrowski type inequalities for isotonic linear functionals," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 3, no. 5, article 68, pp. 1–13, 2002.
- [7] A. Florea and C. P. Niculescu, "A note on Ostrowski's inequality," *Journal of Inequalities and Applications*, no. 5, pp. 459–468, 2005.
- [8] B. G. Pachpatte, "On an inequality of Ostrowski type in three independent variables," *Journal of Mathematical Analysis and Applications*, vol. 249, no. 2, pp. 583–591, 2000.
- [9] B. G. Pachpatte, "On a new Ostrowski type inequality in two independent variables," *Tamkang Journal of Mathematics*, vol. 32, no. 1, pp. 45–49, 2001.