

Research Article

A Recent Note on Quasi-Power Increasing Sequence for Generalized Absolute Summability

E. Savaş¹ and H. Şevli²

¹ Department of Mathematics, İstanbul Ticaret University, Üsküdar, 34672-İstanbul, Turkey

² Department of Mathematics, Faculty of Arts & Sciences, Yüzüncü Yıl University, 65080-Van, Turkey

Correspondence should be addressed to E. Savaş, ekremsavas@yahoo.com

Received 15 May 2009; Accepted 30 July 2009

Recommended by Ramm Mohapatra

We prove two theorems on $|A, \delta|_k$, $k \geq 1, 0 \leq \delta < 1/k$, summability factors for an infinite series by using quasi-power increasing sequences. We obtain sufficient conditions for $\sum a_n \lambda_n$ to be summable $|A, \delta|_k$, $k \geq 1, 0 \leq \delta < 1/k$, by using quasi- f -increasing sequences.

Copyright © 2009 E. Savaş and H. Şevli. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Quite recently, Savaş [1] obtained sufficient conditions for $\sum a_n \lambda_n$ to be summable $|A, \delta|_k$, $k \geq 1, 0 \leq \delta < 1/k$. The purpose of this paper is to obtain the corresponding result for quasi- f -increasing sequence. Our result includes and moderates the conditions of his theorem with the special case $\mu = 0$.

A sequence $\{\lambda_n\}$ is said to be of bounded variation (bv) if $\sum_n |\Delta \lambda_n| < \infty$. Let $bv_0 = bv \cap c_0$, where c_0 denotes the set of all null sequences.

The concept of absolute summability of order $k \geq 1$ was defined by Flett [2] as follows. Let $\sum a_n$ denote a series with partial sums $\{s_n\}$, and A a lower triangular matrix. Then $\sum a_n$ is said to be absolutely A -summable of order $k \geq 1$, written that $\sum a_n$ is summable $|A|_k$, $k \geq 1$, if

$$\sum_{n=1}^{\infty} n^{k-1} |T_{n-1} - T_n|^k < \infty, \quad (1.1)$$

where

$$T_n = \sum_{v=0}^n a_{nv} s_v. \quad (1.2)$$

In [3], Flett considered further extension of absolute summability in which he introduced a further parameter δ . The series $\sum a_n$ is said to be summable $|A, \delta|_k$, $k \geq 1$, $\delta \geq 0$, if

$$\sum_{n=1}^{\infty} n^{\delta k + k - 1} |T_{n-1} - T_n|^k < \infty. \quad (1.3)$$

A positive sequence $\{b_n\}$ is said to be an almost increasing sequence if there exist an increasing sequence $\{c_n\}$ and positive constants A and B such that $Ac_n \leq b_n \leq Bc_n$ (see [4]). Obviously, every increasing sequence is almost increasing. However, the converse need not be true as can be seen by taking the example, say $b_n = e^{(-1)^n} n$.

A positive sequence $\gamma := \{\gamma_n\}$ is said to be a quasi- β -power increasing sequence if there exists a constant $K = K(\beta, \gamma) \geq 1$ such that

$$Kn^\beta \gamma_n \geq m^\beta \gamma_m \quad (1.4)$$

holds for all $n \geq m \geq 1$. It should be noted that every almost increasing sequence is a quasi- β -power increasing sequence for any nonnegative β , but the converse need not be true as can be seen by taking an example, say $\gamma_n = n^{-\beta}$ for $\beta > 0$ (see [5]). If (1.4) stays with $\beta = 0$, then γ is simply called a quasi-increasing sequence. It is clear that if $\{\gamma_n\}$ is quasi- β -power increasing, then $\{n^\beta \gamma_n\}$ is quasi-increasing.

A positive sequence $\gamma = \{\gamma_n\}$ is said to be a quasi- f -power increasing sequence, if there exists a constant $K = K(\gamma, f) \geq 1$ such that $Kf_n \gamma_n \geq f_m \gamma_m$ holds for all $n \geq m \geq 1$, [6].

We may associate A two lower triangular matrices \bar{A} and \hat{A} as follows:

$$\begin{aligned} \bar{a}_{nv} &= \sum_{r=v}^n a_{nr}, \quad n, v = 0, 1, \dots, \\ \hat{a}_{nv} &= \bar{a}_{nv} - \bar{a}_{n-1, v}, \quad n = 1, 2, \dots, \end{aligned} \quad (1.5)$$

where

$$\hat{a}_{00} = \bar{a}_{00} = a_{00}. \quad (1.6)$$

Given any sequence $\{x_n\}$, the notation $x_n \asymp O(1)$ means $x_n = O(1)$ and $1/x_n = O(1)$. For any matrix entry a_{nv} , $\Delta_v a_{nv} := a_{nv} - a_{n, v+1}$.

Quite recently, Savaş [1] obtained sufficient conditions for $\sum a_n \lambda_n$ to be summable $|A, \delta|_k$, $k \geq 1$, $0 \leq \delta < 1/k$ as follows.

Theorem 1.1. *Let A be a lower triangular matrix with nonnegative entries satisfying*

$$a_{n-1,v} \geq a_{nv} \quad \text{for } n \geq v+1, \quad (1.7)$$

$$\bar{a}_{n0} = 1, \quad n = 0, 1, \dots, \quad (1.8)$$

$$na_{nn} \asymp O(1), \quad n \rightarrow \infty, \quad (1.9)$$

$$\sum_{v=1}^{n-1} a_{vv} \hat{a}_{n,v+1} = O(a_{nn}), \quad (1.10)$$

$$\sum_{n=v+1}^{m+1} n^{\delta k} |\Delta_v \hat{a}_{nv}| = O(v^{\delta k} a_{vv}), \quad (1.11)$$

$$\sum_{n=v+1}^{m+1} n^{\delta k} \hat{a}_{n,v+1} = O(v^{\delta k}), \quad (1.12)$$

and let $\{\beta_n\}$ and $\{\lambda_n\}$ be sequences such that

$$|\Delta \lambda_n| \leq \beta_n, \quad (1.13)$$

$$\beta_n \rightarrow 0, \quad n \rightarrow \infty. \quad (1.14)$$

If $\{X_n\}$ is a quasi- β -power increasing sequence for some $0 < \beta < 1$ such that

$$|\lambda_n| X_n = O(1), \quad n \rightarrow \infty, \quad (1.15)$$

$$\sum_{n=1}^{\infty} n X_n |\Delta \beta_n| < \infty, \quad (1.16)$$

$$\sum_{n=1}^m n^{\delta k-1} |s_n|^k = O(X_m), \quad m \rightarrow \infty, \quad (1.17)$$

then the series $\sum a_n \lambda_n$ is summable $|A, \delta|_k$, $k \geq 1$, $0 \leq \delta < 1/k$.

Theorem 1.1 enhanced a theorem of Savas [7] by replacing an almost increasing sequence with a quasi- β -power increasing sequence for some $0 < \beta < 1$. It should be noted that if $\{X_n\}$ is an almost increasing sequence, then (1.15) implies that the sequence $\{\lambda_n\}$ is bounded. However, when $\{X_n\}$ is a quasi- β -power increasing sequence or a quasi- f -increasing sequence, (1.15) does not imply $|\lambda_m| = O(1)$, $m \rightarrow \infty$. For example, since $X_m = m^{-\beta}$ is a quasi- β -power increasing sequence for $0 < \beta < 1$ and if we take $\lambda_m = m^\delta$, $0 < \delta < \beta < 1$, then $|\lambda_m| X_m = m^{\delta-\beta} = O(1)$, $m \rightarrow \infty$ holds but $|\lambda_m| = m^\delta \neq O(1)$ (see [8]). Therefore, we remark that condition $\{\lambda_n\} \in bv_0$ should be added to the statement of Theorem 1.1.

The goal of this paper is to prove the following theorem by using quasi- f -increasing sequences. Our main result includes the moderated version of Theorem 1.1. We will show that the crucial condition of our proof, $\{\lambda_n\} \in bv_0$, can be deduced from another condition of the theorem. Also, we shall eliminate condition (1.15) in our theorem; however we shall deduce this condition from the conditions of our theorem.

2. The Main Results

We now shall prove the following theorems.

Theorem 2.1. *Let A satisfy conditions (1.7)–(1.12), and let $\{\beta_n\}$ and $\{\lambda_n\}$ be sequences satisfying conditions (1.13) and (1.14) of Theorem 1.1 and*

$$\sum_{n=1}^m \lambda_n = o(m), \quad m \rightarrow \infty. \quad (2.1)$$

If $\{X_n\}$ is a quasi- f -increasing sequence and conditions (1.17) and

$$\sum_{n=1}^{\infty} nX_n(\beta, \mu) |\Delta\beta_n| < \infty \quad (2.2)$$

are satisfied, then the series $\sum a_n \lambda_n$ is summable $|A, \delta|_k, k \geq 1, 0 \leq \delta < 1/k$, where $\{f_n\} := \{n^\beta (\log n)^\mu\}, \mu \geq 0, 0 \leq \beta < 1$, and $X_n(\beta, \mu) := (n^\beta (\log n)^\mu X_n)$.

Theorem 2.1 includes the following theorem with the special case $\mu = 0$. Theorem 2.2 moderates the hypotheses of Theorem 1.1.

Theorem 2.2. *Let A satisfy conditions (1.7)–(1.12), and let $\{\beta_n\}$ and $\{\lambda_n\}$ be sequences satisfying conditions (1.13), (1.14), and (2.1). If $\{X_n\}$ is a quasi- β -power increasing sequence for some $0 \leq \beta < 1$ and conditions (1.17) and*

$$\sum_{n=1}^{\infty} nX_n(\beta) |\Delta\beta_n| < \infty \quad (2.3)$$

are satisfied, where $X_n(\beta) := (n^\beta X_n)$, then the series $\sum a_n \lambda_n$ is summable $|A, \delta|_k, k \geq 1, 0 \leq \delta < 1/k$.

Remark 2.3. The crucial condition, $\{\lambda_n\} \in bv_0$, and condition (1.15) do not appear among the conditions of Theorems 2.1 and 2.2. By Lemma 3.3, under the conditions on $\{X_n\}, \{\beta_n\}$, and $\{\lambda_n\}$ as taken in the statement of Theorem 2.1, also in the statement of Theorem 2.2 with the special case $\mu = 0$, conditions $\{\lambda_n\} \in bv_0$ and (1.15) hold.

3. Lemmas

We shall need the following lemmas for the proof of our main Theorem 2.1.

Lemma 3.1 (see [9]). *Let $\{\varphi_n\}$ be a sequence of real numbers and denote*

$$\Phi_n := \sum_{k=1}^n \varphi_k, \quad \Psi_n := \sum_{k=n}^{\infty} |\Delta\varphi_k|. \quad (3.1)$$

If $\Phi_n = o(n)$, then there exists a natural number \mathbb{N} such that

$$|\varphi_n| \leq 2\Psi_n \quad (3.2)$$

for all $n \geq \mathbb{N}$.

Lemma 3.2 (see [8]). *If $\{X_n\}$ is a quasi- f -increasing sequence, where $\{f_n\} = \{n^\beta(\log n)^\mu\}$, $\mu \geq 0$, $0 \leq \beta < 1$, then conditions (2.1) of Theorem 2.1,*

$$\sum_{n=1}^m |\Delta\lambda_n| = o(m), \quad m \rightarrow \infty, \quad (3.3)$$

$$\sum_{n=1}^{\infty} nX_n(\beta, \mu) |\Delta|\Delta\lambda_n|| < \infty, \quad (3.4)$$

where $X_n(\beta, \mu) = (n^\beta(\log n)^\mu X_n)$, imply conditions (1.15) and

$$\lambda_n \rightarrow 0, \quad n \rightarrow \infty. \quad (3.5)$$

Lemma 3.3. *If $\{X_n\}$ is a quasi- f -increasing sequence, where $\{f_n\} = \{n^\beta(\log n)^\mu\}$, $\mu \geq 0$, $0 \leq \beta < 1$, then, under conditions (1.13), (1.14), (2.1), and (2.2), conditions (1.15) and (3.5) are satisfied.*

Proof. It is clear that (1.13) and (1.14) \Rightarrow (3.3). Also, (1.13) and (2.2) \Rightarrow (3.4). By Lemma 3.2, under conditions (1.13)-(1.14) and (2.1)-(2.2), we have (1.15) and (3.5). \square

Lemma 3.4. *Let $\{X_n\}$ be a quasi- f -increasing sequence, where $\{f_n\} = \{n^\beta(\log n)^\mu\}$, $\mu \geq 0$, $0 \leq \beta < 1$. If conditions (1.13), (1.14), and (2.2) are satisfied, then*

$$n\beta_n X_n = O(1), \quad (3.6)$$

$$\sum_{n=1}^{\infty} \beta_n X_n < \infty. \quad (3.7)$$

Proof. It is clear that if $\{X_n\}$ is quasi- f -increasing, then $\{n^\beta(\log n)^\mu X_n\}$ is quasi-increasing. Since $\beta_n \rightarrow 0$, $n \rightarrow \infty$, from the fact that $\{n^{1-\beta}(\log n)^{-\mu}\}$ is increasing and (2.2), we have

$$\begin{aligned} n\beta_n X_n &= nX_n \sum_{k=n}^{\infty} |\Delta\beta_k| \\ &= O(1)n^{1-\beta}(\log n)^{-\mu} \sum_{k=n}^{\infty} k^\beta(\log k)^\mu X_k |\Delta\beta_k| \\ &= O(1) \sum_{k=n}^{\infty} kX_k |\Delta\beta_k| = O(1). \end{aligned} \quad (3.8)$$

Again using (2.2),

$$\begin{aligned}
 \sum_{n=1}^{\infty} \beta_n X_n &= O(1) \sum_{n=1}^{\infty} X_n \sum_{k=n}^{\infty} |\Delta \beta_k| \\
 &= O(1) \sum_{k=1}^{\infty} |\Delta \beta_k| \sum_{n=1}^k n^\beta (\log n)^\mu X_n n^{-\beta} (\log n)^{-\mu} \\
 &= O(1) \sum_{k=1}^{\infty} k^\beta (\log k)^\mu X_k |\Delta \beta_k| \sum_{n=1}^k n^{-\beta} (\log n)^{-\mu} \\
 &= O(1) \sum_{k=1}^{\infty} k X_k(\beta, \mu) |\Delta \beta_k| = O(1).
 \end{aligned} \tag{3.9}$$

□

4. Proof of Theorem 2.1

Let y_n denote the n th term of the A -transform of the series $\sum a_n \lambda_n$. Then, by definition, we have

$$y_n = \sum_{i=0}^n a_{ni} s_i = \sum_{v=0}^n \bar{a}_{nv} \lambda_v a_v. \tag{4.1}$$

Then, for $n \geq 1$, we have

$$Y_n := y_n - y_{n-1} = \sum_{v=0}^n \hat{a}_{nv} \lambda_v a_v. \tag{4.2}$$

Applying Abel's transformation, we may write

$$Y_n = \sum_{v=1}^{n-1} \Delta_v (\hat{a}_{nv} \lambda_v) \sum_{r=1}^v a_r + \hat{a}_{nn} \lambda_n \sum_{v=1}^n a_v. \tag{4.3}$$

Since

$$\Delta_v (\hat{a}_{nv} \lambda_v) = \lambda_v \Delta_v \hat{a}_{nv} + \Delta \lambda_v \hat{a}_{n,v+1}, \tag{4.4}$$

we have

$$\begin{aligned}
 Y_n &= a_{nn} \lambda_n s_n + \sum_{v=1}^{n-1} \Delta_v \hat{a}_{nv} \lambda_v s_v + \sum_{v=1}^{n-1} \hat{a}_{n,v+1} \Delta \lambda_v s_v \\
 &= Y_{n,1} + Y_{n,2} + Y_{n,3}, \text{ say.}
 \end{aligned} \tag{4.5}$$

Since

$$|Y_{n,1} + Y_{n,2} + Y_{n,3}|^k \leq 3^k (|Y_{n,1}|^k + |Y_{n,2}|^k + |Y_{n,3}|^k), \quad (4.6)$$

to complete the proof, it is sufficient to show that

$$\sum_{n=1}^{\infty} n^{\delta k+k-1} |Y_{n,r}|^k < \infty, \quad \text{for } r = 1, 2, 3. \quad (4.7)$$

Since $\{\lambda_n\}$ is bounded by Lemma 3.3, using (1.9), we have

$$\begin{aligned} I_1 &= \sum_{n=1}^m n^{\delta k+k-1} |Y_{n,1}|^k = \sum_{n=1}^m n^{\delta k+k-1} |a_{nn} \lambda_n s_n|^k \\ &\leq \sum_{n=1}^m n^{\delta k} (n a_{nn})^{k-1} a_{nn} |\lambda_n|^{k-1} |\lambda_n| |s_n|^k \\ &= O(1) \sum_{n=1}^m n^{\delta k} a_{nn} |\lambda_n| |s_n|^k. \end{aligned} \quad (4.8)$$

Using properties (1.15), in view of Lemma 3.3, and (3.7), from (1.9), (1.13), and (1.17),

$$\begin{aligned} I_1 &= O(1) \sum_{n=1}^{m-1} |\Delta \lambda_n| \sum_{v=1}^n v^{\delta k} a_{vv} |s_v|^k + O(1) |\lambda_m| \sum_{v=1}^m v^{\delta k} a_{vv} |s_v|^k \\ &= O(1) \sum_{n=1}^{m-1} |\Delta \lambda_n| \sum_{v=1}^n v^{\delta k-1} |s_v|^k + O(1) |\lambda_m| \sum_{v=1}^m v^{\delta k-1} |s_v|^k \\ &= O(1) \sum_{n=1}^{m-1} \beta_n X_n + O(1) |\lambda_m| X_m = O(1) \quad \text{as } m \rightarrow \infty. \end{aligned} \quad (4.9)$$

Applying Hölder's inequality,

$$\begin{aligned} I_2 &= \sum_{n=2}^{m+1} n^{\delta k+k-1} |Y_{n,2}|^k = O(1) \sum_{n=2}^{m+1} n^{\delta k+k-1} \left(\sum_{v=1}^{n-1} |\Delta_v \hat{a}_{nv}| |\lambda_v| |s_v| \right)^k \\ &= O(1) \sum_{n=2}^{m+1} n^{\delta k+k-1} \sum_{v=1}^{n-1} |\Delta_v \hat{a}_{nv}| |\lambda_v|^k |s_v|^k \left(\sum_{v=1}^{n-1} |\Delta_v \hat{a}_{nv}| \right)^{k-1}. \end{aligned} \quad (4.10)$$

Using (1.9) and (1.11) and boundedness of $\{\lambda_n\}$,

$$\begin{aligned}
 I_2 &= O(1) \sum_{n=2}^{m+1} n^{\delta k} (na_{nm})^{k-1} \sum_{v=1}^{n-1} |\Delta_v \hat{a}_{nv}| |s_v|^k |\lambda_v|^{k-1} |\lambda_v| \\
 &= O(1) \sum_{v=1}^m |\lambda_v| |s_v|^k \sum_{n=v+1}^{m+1} n^{\delta k} |\Delta_v \hat{a}_{nv}| \\
 &= O(1) \sum_{v=1}^m v^{\delta k} a_{vv} |\lambda_v| |s_v|^k = O(1), \quad \text{as } m \rightarrow \infty,
 \end{aligned} \tag{4.11}$$

as in the proof of I_1 .

Finally, again using Hölder's inequality, from (1.9), (1.10), and (1.12),

$$\begin{aligned}
 I_3 &= \sum_{n=2}^{m+1} n^{\delta k+k-1} |Y_{n,3}|^k = O(1) \sum_{n=2}^{m+1} n^{\delta k+k-1} \left(\sum_{v=1}^{n-1} \hat{a}_{n,v+1} |\Delta \lambda_v| |s_v| \right)^k \\
 &= O(1) \sum_{n=2}^{m+1} n^{\delta k+k-1} \sum_{v=1}^{n-1} \hat{a}_{n,v+1} |\Delta \lambda_v|^k |s_v|^k a_{vv}^{1-k} \left(\sum_{v=1}^{n-1} a_{vv} \hat{a}_{n,v+1} \right)^{k-1} \\
 &= O(1) \sum_{n=2}^{m+1} n^{\delta k} \sum_{v=1}^{n-1} \hat{a}_{n,v+1} |\Delta \lambda_v|^k |s_v|^k a_{vv}^{1-k} \\
 &= O(1) \sum_{v=1}^m |\Delta \lambda_v|^k |s_v|^k a_{vv}^{1-k} \sum_{n=v+1}^{m+1} n^{\delta k} \hat{a}_{n,v+1} \\
 &= O(1) \sum_{v=1}^m (v |\Delta \lambda_v|)^k v^{\delta k} a_{vv} |s_v|^k.
 \end{aligned} \tag{4.12}$$

By Lemma 3.1, condition (3.3), in view of Lemma 3.3, implies that

$$n |\Delta \lambda_n| \leq 2n \sum_{k=n}^{\infty} |\Delta \lambda_k| \leq 2 \sum_{k=n}^{\infty} k |\Delta \lambda_k| \tag{4.13}$$

holds. Thus, by Lemma 3.3, (3.4) implies that $\{n |\Delta \lambda_n|\}$ is bounded. Therefore, from (1.9) and (1.13),

$$\begin{aligned}
 I_3 &= O(1) \sum_{v=1}^m (v |\Delta \lambda_v|)^{k-1} v |\Delta \lambda_v| v^{\delta k} a_{vv} |s_v|^k \\
 &= O(1) \sum_{v=1}^m v \beta_v v^{\delta k-1} |s_v|^k.
 \end{aligned} \tag{4.14}$$

Using Abel transformation and (1.17),

$$\begin{aligned} I_3 &= O(1) \sum_{v=1}^{m-1} |\Delta(v\beta_v)| \left(\sum_{r=1}^v r^{\delta k-1} |s_r|^k \right) + O(1) m \beta_m \sum_{v=1}^m v^{\delta k-1} |s_v|^k \\ &= O(1) \sum_{v=1}^{m-1} |\Delta(v\beta_v)| X_v + O(1) m \beta_m X_m. \end{aligned} \quad (4.15)$$

Since

$$\Delta(v\beta_v) = v\beta_v - (v+1)\beta_{v+1} = v\Delta\beta_v - \beta_{v+1}, \quad (4.16)$$

we have

$$\begin{aligned} I_3 &= O(1) \sum_{v=1}^{m-1} v X_v |\Delta\beta_v| + O(1) \sum_{v=1}^{m-1} X_{v+1} \beta_{v+1} + O(1) m X_m \beta_m \\ &= O(1), \quad \text{as } m \rightarrow \infty, \end{aligned} \quad (4.17)$$

by virtue of (2.2) and properties (3.6) and (3.7) of Lemma 3.4.

So we obtain (4.7). This completes the proof.

5. Corollaries and Applications to Weighted Means

Setting $\delta = 0$ in Theorems 2.1 and 2.2 yields the following two corollaries, respectively.

Corollary 5.1. *Let A satisfy conditions (1.7)–(1.10), and let $\{\beta_n\}$ and $\{\lambda_n\}$ be sequences satisfying conditions (1.13), (1.14), and (2.1). If $\{X_n\}$ is a quasi- f -increasing sequence, where $\{f_n\} := \{n^\beta (\log n)^\mu\}$, $\mu \geq 0$, $0 \leq \beta < 1$, and conditions (2.2) and*

$$\sum_{n=1}^m \frac{1}{n} |s_n|^k = O(X_m), \quad m \rightarrow \infty, \quad (5.1)$$

are satisfied, then the series $\sum a_n \lambda_n$ is summable $|A|_k$, $k \geq 1$.

Proof. If we take $\delta = 0$ in Theorem 2.1, then condition (1.17) reduces condition (5.1). In this case conditions (1.11) and (1.12) are obtained by conditions (1.7)–(1.10). \square

Corollary 5.2. *Let A satisfy conditions (1.7)–(1.10), and let $\{\beta_n\}$ and $\{\lambda_n\}$ be sequences satisfying conditions (1.13), (1.14), and (2.1). If $\{X_n\}$ is a quasi- β -power increasing sequence for some $0 \leq \beta < 1$ and conditions (2.3) and (5.1) are satisfied, then the series $\sum a_n \lambda_n$ is summable $|A|_k$, $k \geq 1$.*

A weighted mean matrix, denoted by (\overline{N}, p_n) , is a lower triangular matrix with entries $a_{nv} = p_v / P_n$, where $\{p_n\}$ is nonnegative sequence with $p_0 > 0$ and $P_n := \sum_{v=0}^n p_v \rightarrow \infty$, as $n \rightarrow \infty$.

Corollary 5.3. Let $\{p_n\}$ be a positive sequence satisfying

$$np_n \asymp O(P_n), \quad \text{as } n \rightarrow \infty, \quad (5.2)$$

$$\sum_{n=v+1}^{m+1} n^{\delta k} \frac{p_n}{P_n P_{n-1}} = O\left(\frac{v^{\delta k}}{P_v}\right), \quad (5.3)$$

and let $\{\beta_n\}$ and $\{\lambda_n\}$ be sequences satisfying conditions (1.13), (1.14), and (2.1). If $\{X_n\}$ is a quasi- f -increasing sequence, where $\{f_n\} := \{n^\beta (\log n)^\mu\}$, $\mu \geq 0$, $0 \leq \beta < 1$, and conditions (1.17) and (2.2) are satisfied, then the series, $\sum a_n \lambda_n$ is summable $|\overline{N}, p_n, \delta|_k$ for $k \geq 1$ and $0 \leq \delta < 1/k$.

Proof. In Theorem 2.1 set $A = (\overline{N}, p_n)$. It is clear that conditions (1.7), (1.8), and (1.10) are automatically satisfied. Condition (1.9) becomes condition (5.2), and conditions (1.11) and (1.12) become condition (5.3) for weighted mean method. \square

Corollary 5.3 includes the following result with the special case $\mu = 0$.

Corollary 5.4. Let $\{p_n\}$ be a positive sequence satisfying (5.2) and (5.3), and let $\{X_n\}$ be a quasi- β -power increasing sequence for some $0 \leq \beta < 1$. Then under conditions (1.13), (1.14), (1.17), (2.1), and (2.3), $\sum a_n \lambda_n$ is summable $|\overline{N}, p_n, \delta|_k$, $k \geq 1$, $0 \leq \delta < 1/k$.

References

- [1] E. Savaş, "Quasi-power increasing sequence for generalized absolute summability," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 68, no. 1, pp. 170–176, 2008.
- [2] T. M. Flett, "On an extension of absolute summability and some theorems of Littlewood and Paley," *Proceedings of the London Mathematical Society*, vol. 7, pp. 113–141, 1957.
- [3] T. M. Flett, "Some more theorems concerning the absolute summability of Fourier series and power series," *Proceedings of the London Mathematical Society*, vol. 8, pp. 357–387, 1958.
- [4] S. Alijancic and D. Arendelovic, "O-regularly varying functions," *Publications de l'Institut Mathématique*, vol. 22, no. 36, pp. 5–22, 1977.
- [5] L. Leindler, "A new application of quasi power increasing sequences," *Publicationes Mathematicae Debrecen*, vol. 58, no. 4, pp. 791–796, 2001.
- [6] W. T. Sulaiman, "Extension on absolute summability factors of infinite series," *Journal of Mathematical Analysis and Applications*, vol. 322, no. 2, pp. 1224–1230, 2006.
- [7] E. Savaş, "On almost increasing sequences for generalized absolute summability," *Mathematical Inequalities & Applications*, vol. 9, no. 4, pp. 717–723, 2006.
- [8] H. Şevli and L. Leindler, "On the absolute summability factors of infinite series involving quasi-power-increasing sequences," *Computers & Mathematics with Applications*, vol. 57, no. 5, pp. 702–709, 2009.
- [9] L. Leindler, "A note on the absolute Riesz summability factors," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 6, no. 4, article 96, 5 pages, 2005.