

## Research Article

# A Note on $(C_p, \alpha)$ -Hyponormal Operators

**Xiaohuan Wang<sup>1</sup> and Zongsheng Gao<sup>2</sup>**

<sup>1</sup> LMIB and School of Mathematics and Systems Science, Beihang University, Beijing 100191, China

<sup>2</sup> LMIB and Department of Mathematics, Beihang University, Beijing 100191, China

Correspondence should be addressed to Xiaohuan Wang, xiaohuan@smss.buaa.edu.cn

Received 20 January 2010; Accepted 22 April 2010

Academic Editor: Sin Takahasi

Copyright © 2010 X. Wang and Z. Gao. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study  $(C_p, \alpha)$ -normal operators and  $(C_p, \alpha)$ -hyponormal operators. We show the inclusion relation between these classes under various hypotheses for  $p$  and  $\alpha$ . We also obtain some sufficient conditions for Aluthge transform  $\tilde{T}_{s,t} = |T|^s U |T|^t$  and  $T^2$  of  $(C_p, \alpha)$ -hyponormal operators still to be  $(C_p, \alpha)$ -hyponormal.

## 1. Introduction

Let  $\mathcal{H}$  be a separable, infinite dimensional, complex Hilbert space, and denote by  $\mathcal{L}(\mathcal{H})$  the algebra of all bounded linear operators on  $\mathcal{H}$ . Recently, Lauric in [1] introduced  $(C_p, \alpha)$ -hyponormal operators. For  $\alpha > 0$  and  $T \in \mathcal{L}(\mathcal{H})$ , denote by  $D_T^\alpha = (T^*T)^\alpha - (TT^*)^\alpha$ . We denote that  $C_p(\mathcal{H})$ ,  $1 \leq p < \infty$ , the ideal of operators in the Schatten  $p$ -class [2]. Although, for  $0 < p < 1$ , the usual definition of  $\|\cdot\|_p$  does not satisfy the triangle inequality, nevertheless  $(C_p, \|\cdot\|_p)$  is closed and  $\|TK\|_p \leq \|T\| \cdot \|K\|_p$ , when  $T \in \mathcal{L}(\mathcal{H})$  and  $K \in C_p(\mathcal{H})$ . An operator  $T$  in  $\mathcal{L}(\mathcal{H})$  is  $(C_p, \alpha)$ -normal if  $D_T^\alpha \in C_p(\mathcal{H})$ , and denote the class of  $(C_p, \alpha)$ -normal operators by  $\mathcal{N}_p^\alpha(\mathcal{H})$ . An operator  $T$  in  $\mathcal{L}(\mathcal{H})$  will be called  $(C_p, \alpha)$ -hyponormal if  $D_T^\alpha = P + K$ , where  $P$  is a positive semidefinite operator ( $P \geq 0$ ) and  $K \in C_p(\mathcal{H})$ . The class of  $(C_p, \alpha)$ -hyponormal operators will be denoted by  $\mathcal{H}_p^\alpha(\mathcal{H})$ . In particular, an operator  $T$  in  $\mathcal{H}_1^1(\mathcal{H})$  will be called almost hyponormal. Furthermore, an operator  $T \in \mathcal{L}(\mathcal{H})$  whose  $D_T^\alpha$  is positive semidefinite is called  $\alpha$ -hyponormal (notation:  $T \in \mathcal{H}_0^\alpha(\mathcal{H})$ ).

In this paper, we first study the inclusion relation between these classes under various hypotheses for  $p$  and  $\alpha$  in Section 2. Then we study the Aluthge transform  $\tilde{T}_{s,t} = |T|^s U |T|^t$  and  $T^2$  of  $(C_p, \alpha)$ -hyponormal operators in Section 3.

Before proceeding, we will make use of the following inequality.

**Theorem F** (See Furuta inequality in [3]). If  $A \geq B \geq 0$ , then, for each  $r \geq 0$ ,

$$\begin{aligned} (B^{r/2} A^p B^{r/2})^{1/q} &\geq (B^{r/2} B^p B^{r/2})^{1/q}, \\ (A^{r/2} A^p A^{r/2})^{1/q} &\geq (A^{r/2} B^p A^{r/2})^{1/q}, \end{aligned} \quad (1.1)$$

as long as real numbers  $p, r, q$  satisfy

$$p \geq 0, q \geq 1 \text{ with } (1+r)q \geq p+r. \quad (1.2)$$

**Lemma 1.1** (see [1]). Let  $A \in \mathcal{L}(\mathcal{H})$ ,  $A \geq 0$ ,  $\alpha \in (0, 1]$ ,  $p \geq \alpha$ , and  $K \in \mathcal{C}_p(\mathcal{H})$ , such that  $A + K \geq 0$ . Then  $(A + K)^\alpha = A^\alpha + K_1$ , where  $K_1 \in \mathcal{C}_{p/\alpha}(\mathcal{H})$ . If in addition  $K \geq 0$ , then  $K_1 \geq 0$ .

**Lemma 1.2** (see [1]). Let  $A \in \mathcal{L}(\mathcal{H})$ ,  $A \geq 0$ ,  $p \geq 1$ , and  $K \in \mathcal{C}_p(\mathcal{H})$ , such that  $A + K \geq 0$ , and let  $\alpha \in [1, +\infty)$ . Then  $(A + K)^\alpha = A^\alpha + K_1$ , where  $K_1 \in \mathcal{C}_p(\mathcal{H})$ .

## 2. Some Inclusions

According to Löwner-Heinz (L-H) inequality [4, 5] that  $A \geq B \geq 0$  ensures that  $A^\alpha \geq B^\alpha$  for each  $\alpha \in [0, 1]$ , we obtain  $\mathcal{L}_0^\alpha(\mathcal{H}) \supseteq \mathcal{L}_0^\beta(\mathcal{H})$  when  $\alpha \leq \beta$ . However, the similar inclusions for the classes  $\mathcal{N}_p^\alpha(\mathcal{H})$  and  $\mathcal{L}_p^\alpha(\mathcal{H})$  are less obvious. In this section, we will examine various inclusions between these classes of operators. (1) of Theorem 2.1 has been already shown in [1]. But we will give a proof for the readers' convenience.

**Theorem 2.1.** Let  $\alpha > 0$ ,  $p \geq 1$ , and let  $T$  be in  $\mathcal{N}_p^\alpha(\mathcal{H})$ .

- (1) If  $\beta \geq \alpha$ , then  $T$  belongs to  $\mathcal{N}_p^\beta(\mathcal{H})$ , and therefore  $\mathcal{N}_p^\alpha(\mathcal{H}) \subseteq \mathcal{N}_p^\beta(\mathcal{H})$ .
- (2) If  $0 < \beta \leq \alpha$ , then  $T$  belongs to  $\mathcal{N}_{\alpha p/\beta}^\beta(\mathcal{H})$ , and therefore  $\mathcal{N}_p^\alpha(\mathcal{H}) \subseteq \mathcal{N}_{\alpha p/\beta}^\beta(\mathcal{H})$ .

*Proof.* Let  $\alpha$ ,  $p$ , and  $T$  be as in the hypotheses and let  $T = U|T|$  be the polar decomposition of  $T$ .

For  $T \in \mathcal{N}_p^\alpha(\mathcal{H})$ , we have

$$D_T^\alpha = (T^*T)^\alpha - (TT^*)^\alpha = |T|^{2\alpha} - |T^*|^{2\alpha} = K, \quad (2.1)$$

with  $K \in \mathcal{C}_p(\mathcal{H})$ . Then we obtain

$$|T|^{2\alpha} = |T^*|^{2\alpha} + K \geq 0. \quad (2.2)$$

- (1) First we consider the case  $\beta \geq \alpha$ . According to Lemma 1.2, we obtain

$$|T|^{2\beta} = \left(|T^*|^{2\alpha} + K\right)^{\beta/\alpha} = |T^*|^{2\beta} + K_1, \quad (2.3)$$

with  $K_1 \in \mathcal{C}_p(\mathcal{H})$ . Then  $T \in \mathcal{N}_p^\beta(\mathcal{H})$ .

(2) Next we consider the case  $0 < \beta \leq \alpha$ . According to Lemma 1.1, we obtain

$$|T|^{2\beta} = \left(|T^*|^{2\alpha} + K\right)^{\beta/\alpha} = |T^*|^{2\beta} + K_1, \quad (2.4)$$

with  $K_1 \in \mathcal{C}_{\alpha p/\beta}(\mathcal{H})$ . Then  $T \in \mathcal{N}_{\alpha p/\beta}^\beta(\mathcal{H})$ .  $\square$

The following corollary is a consequence of Theorem 2.1.

**Corollary 2.2.** *Let  $\alpha > 0$ ,  $p \geq 1$ , then, for  $0 < \beta \leq \alpha$ ,*

$$\mathcal{N}_p^\beta(\mathcal{H}) \subseteq \mathcal{N}_p^\alpha(\mathcal{H}) \subseteq \mathcal{N}_{\alpha p/\beta}^\beta(\mathcal{H}) \subseteq \mathcal{N}_{\alpha p/\beta}^\alpha(\mathcal{H}). \quad (2.5)$$

**Theorem 2.3.** *Let  $\alpha > 0$ ,  $p \geq 1$ , and let  $T$  be in  $\mathcal{H}_p^\alpha(\mathcal{H})$ . If  $0 < \beta \leq \alpha$ , then  $T$  belongs to  $\mathcal{H}_{\alpha p/\beta}^\beta(\mathcal{H})$ , and therefore  $\mathcal{H}_p^\alpha(\mathcal{H}) \subseteq \mathcal{H}_{\alpha p/\beta}^\beta(\mathcal{H})$ .*

*Proof.* Let  $\alpha$ ,  $p$ , and  $T$  be as in the hypotheses and let  $T = U|T|$  be the polar decomposition of  $T$ .

For  $T \in \mathcal{H}_p^\alpha(\mathcal{H})$ , we have

$$D_T^\alpha = (T^*T)^\alpha - (TT^*)^\alpha = |T|^{2\alpha} - |T^*|^{2\alpha} = P + K, \quad (2.6)$$

with  $P \geq 0$ ,  $K \in \mathcal{C}_p(\mathcal{H})$ . Then we obtain

$$|T|^{2\alpha} = |T^*|^{2\alpha} + P + K \geq 0. \quad (2.7)$$

For  $0 < \beta \leq \alpha$ , according to Lemma 1.1 and L-H inequality, we obtain

$$\begin{aligned} |T|^{2\beta} &= \left(|T^*|^{2\alpha} + P + K\right)^{\beta/\alpha} \\ &= \left(|T^*|^{2\alpha} + P\right)^{\beta/\alpha} + K_1 \\ &\geq |T^*|^{2\beta} + K_1, \end{aligned} \quad (2.8)$$

with  $K_1 \in \mathcal{C}_{\alpha p/\beta}(\mathcal{H})$ . Then we obtain  $T \in \mathcal{H}_{\alpha p/\beta}^\beta(\mathcal{H})$ .  $\square$

### 3. Some Properties of $(\mathcal{C}_p, \alpha)$ -Hyponormal Operators

Let  $T = U|T|$  be the polar decomposition of an operator  $T$  on a Hilbert space  $\mathcal{H}$ , where  $U$  is a partial isometry operator. Recently, Lauric [1] shows some theorems on the Aluthge transform  $\tilde{T} = |T|^{1/2}U|T|^{1/2}$  of  $(\mathcal{C}_p, \alpha)$ -hyponormal operators. In this section, we will show some sufficient conditions for the generalized Aluthge transform  $\tilde{T}_{s,t} = |T|^s U |T|^t$  ( $s, t > 0$ ) and

$T^2$  of  $(\mathcal{C}_p, \alpha)$ -hyponormal operators to be  $(\mathcal{C}_p, \alpha)$ -hyponormal. Aluthge transform  $\tilde{T}_{s,t}$  arose in the study of  $p$ -hyponormal operators [6, 7] and has since been studied in many different contexts [8–15].

Let  $T$  belong to  $\mathcal{H}_p^\alpha(\mathcal{H})$ , for some  $\alpha > 0, p > 0$ , such that  $D_T^\alpha = P + K$  with  $P \geq 0, K \in \mathcal{C}_p(\mathcal{H})$ . Since  $K = K^* = K_+ - K_-$  and  $K_+, K_- \geq 0$  are  $\mathcal{C}_p$ -class operators, in what follows we will assume that  $D_T^\alpha = P_1 - K_1$  with  $P_1 \geq 0$  and  $K_1 \geq 0, K_1 \in \mathcal{C}_p(\mathcal{H})$ .

**Theorem 3.1.** *Let  $p \geq 1, \alpha \geq \max\{s, t\}$ , and  $T \in \mathcal{H}_p^\alpha(\mathcal{H})$  such that  $D_T^\alpha = P - K$  with  $P, K \geq 0, K \in \mathcal{C}_p(\mathcal{H})$ , and let  $\varepsilon \in (0, 1/2]$  such that  $\alpha + \varepsilon \leq 1$ . Then  $\tilde{T}_{s,t} \in \mathcal{H}_{2\alpha p/(\alpha+\varepsilon)s}^{(\alpha+\varepsilon)}(\mathcal{H})$ .*

*Proof.* We may assume that  $T = U|T|$  with  $U$  being unitary. The equality  $D_T^\alpha = P - K$  with  $P, K \geq 0$  implies that  $|T|^{2\alpha} + K \geq U|T|^{2\alpha}U^*$ . Multiplying this inequality by  $U^*$  to the left and by  $U$  to the right, we obtain

$$A = U^*|T|^{2\alpha}U + U^*KU \geq |T|^{2\alpha} = B. \quad (3.1)$$

According to Lemma 1.1,

$$A^{s/\alpha} = \left\{ U^* \left( |T|^{2\alpha} + K \right) U \right\}^{s/\alpha} = U^* \left( |T|^{2\alpha} + K \right)^{s/\alpha} U = U^* \left( |T|^{2s} + K_1 \right) U, \quad (3.2)$$

with  $K_1 \in \mathcal{C}_{\alpha p/s}(\mathcal{H})$ . Setting  $K_2 = |T|^t U^* K_1 U |T|^t$ , by Theorem F we have

$$\begin{aligned} \left( \tilde{T}_{s,t}^* \tilde{T}_{s,t} + K_2 \right)^{\alpha+\varepsilon} &= \left\{ |T|^t \left[ U^* \left( |T|^{2s} + K_1 \right) U \right] |T|^t \right\}^{\alpha+\varepsilon} \\ &= \left\{ |T|^t \left[ U^* \left( |T|^{2\alpha} + K \right) U \right]^{s/\alpha} |T|^t \right\}^{\alpha+\varepsilon} \\ &= \left( B^{t/2\alpha} A^{s/\alpha} B^{t/2\alpha} \right)^{\alpha+\varepsilon} \\ &\geq B^{(s+t)(\alpha+\varepsilon)/\alpha} \\ &= |T|^{2(s+t)(\alpha+\varepsilon)}. \end{aligned} \quad (3.3)$$

On the other hand, according to Lemma 1.1,

$$\left( \tilde{T}_{s,t}^* \tilde{T}_{s,t} + K_2 \right)^{\alpha+\varepsilon} = \left( \tilde{T}_{s,t}^* \tilde{T}_{s,t} \right)^{\alpha+\varepsilon} + K_3, \quad (3.4)$$

with  $K_3 \in \mathcal{C}_{\alpha p/(\alpha+\varepsilon)s}(\mathcal{H})$ . Then we have

$$\left( \tilde{T}_{s,t}^* \tilde{T}_{s,t} \right)^{\alpha+\varepsilon} + K_3 \geq |T|^{2(s+t)(\alpha+\varepsilon)}. \quad (3.5)$$

According to the following inequality

$$C = |T|^{2\alpha} + K \geq U|T|^{2\alpha}U^* = D, \quad (3.6)$$




Figure 1: Domain of Furuta inequality.

by Theorem F, we have

$$\left(C^{s/2\alpha} D^{t/\alpha} C^{s/2\alpha}\right)^{\alpha+\varepsilon} \leq C^{(s+t)(\alpha+\varepsilon)/\alpha}. \quad (3.7)$$

Again, according to Lemma 1.1,

$$C^{s/2\alpha} = \left(|T|^{2\alpha} + K\right)^{s/2\alpha} = |T|^s + K_4, \quad (3.8)$$

with  $K_4 \in \mathcal{C}_{2\alpha p/s}(\mathcal{H})$ .

Next, obviously,

$$D^{t/\alpha} = \left(U|T|^{2\alpha}U^*\right)^{t/\alpha} = U|T|^{2t}U^*. \quad (3.9)$$

Then we have

$$\begin{aligned} \left(C^{s/2\alpha} D^{t/\alpha} C^{s/2\alpha}\right)^{\alpha+\varepsilon} &= \left\{ (|T|^s + K_4) \left( U|T|^{2t}U^* \right) (|T|^s + K_4) \right\}^{\alpha+\varepsilon} \\ &= \left( |T|^s U|T|^{2t}U^* |T|^s + K_5 \right)^{\alpha+\varepsilon} \\ &= \left( \tilde{T}_{s,t} \tilde{T}_{s,t}^* + K_5 \right)^{\alpha+\varepsilon} \\ &= \left( \tilde{T}_{s,t} \tilde{T}_{s,t}^* \right)^{\alpha+\varepsilon} + K_6, \end{aligned} \quad (3.10)$$

with  $K_5 \in \mathcal{C}_{2\alpha p/s}(\mathcal{H})$ ,  $K_6 \in \mathcal{C}_{2\alpha p/(\alpha+\varepsilon)s}(\mathcal{H})$ .

(1) First we consider the case  $0 \leq ((s+t)/\alpha) \leq 1$ . According to Lemma 1.1, we have

$$\begin{aligned} \left(C^{s+t/\alpha}\right)^{\alpha+\varepsilon} &= \left\{\left(|T|^{2\alpha} + K\right)^{s+t/\alpha}\right\}^{\alpha+\varepsilon} \\ &= \left(|T|^{2(s+t)} + K_7\right)^{\alpha+\varepsilon} \\ &= |T|^{2(s+t)(\alpha+\varepsilon)} + K_8, \end{aligned} \quad (3.11)$$

with  $K_7 \in \mathcal{C}_{ap/s+t}(\mathcal{A})$  and  $K_8 \in \mathcal{C}_{ap/(\alpha+\varepsilon)(s+t)}(\mathcal{A})$ .

Then by (3.7) and (3.10), set  $K_9 = K_6 - K_8 \in \mathcal{C}_{2ap/(\alpha+\varepsilon)s}(\mathcal{A})$ , and

$$|T|^{2(s+t)(\alpha+\varepsilon)} \geq \left(\tilde{T}_{s,t} \tilde{T}_{s,t}^*\right)^{\alpha+\varepsilon} + K_9. \quad (3.12)$$

Combining (3.5) and (3.12), we obtain

$$\left(\tilde{T}_{s,t}^* \tilde{T}_{s,t}\right)^{\alpha+\varepsilon} - \left(\tilde{T}_{s,t} \tilde{T}_{s,t}^*\right)^{\alpha+\varepsilon} \geq K_{10}, \quad (3.13)$$

where  $K_{10} = K_9 - K_3 \in \mathcal{C}_{2ap/(\alpha+\varepsilon)s}(\mathcal{A})$ .

(2) Next we consider the case  $(s+t/\alpha) > 1$ . According to Lemmas 1.1 and 1.2,

$$\begin{aligned} \left(C^{s+t/\alpha}\right)^{\alpha+\varepsilon} &= \left\{\left(|T|^{2\alpha} + K\right)^{s+t/\alpha}\right\}^{\alpha+\varepsilon} \\ &= \left(|T|^{2(s+t)} + K'_7\right)^{\alpha+\varepsilon} \\ &= |T|^{2(s+t)(\alpha+\varepsilon)} + K'_8, \end{aligned} \quad (3.14)$$

with  $K'_7 \in \mathcal{C}_p(\mathcal{A})$  and  $K'_8 \in \mathcal{C}_{p/\alpha+\varepsilon}(\mathcal{A})$ . and

Then by (3.7) and (3.10), set  $K'_9 = K_6 - K'_8 \in \mathcal{C}_{2ap/(\alpha+\varepsilon)s}(\mathcal{A})$ ,

$$|T|^{2(s+t)(\alpha+\varepsilon)} \geq \left(\tilde{T}_{s,t} \tilde{T}_{s,t}^*\right)^{\alpha+\varepsilon} + K'_9. \quad (3.15)$$

Combining (3.5) and (3.15), we obtain

$$\left(\tilde{T}_{s,t}^* \tilde{T}_{s,t}\right)^{\alpha+\varepsilon} - \left(\tilde{T}_{s,t} \tilde{T}_{s,t}^*\right)^{\alpha+\varepsilon} \geq K'_{10}, \quad (3.16)$$

where  $K'_{10} = K'_9 - K_3 \in \mathcal{C}_{2ap/(\alpha+\varepsilon)s}(\mathcal{A})$ .

By (3.13) and (3.16), we obtain  $\tilde{T}_{s,t} \in \mathcal{A}_{2ap/(\alpha+\varepsilon)s}^{(\alpha+\varepsilon)}(\mathcal{A})$ . □

**Remark 3.2.** The main theorem of [1] was considered in the case  $\alpha \in [1/2, 1]$ . Apparently, Theorem 3.1 implies (Theorems 13 in [1]) when  $s = t = 1/2$ . And we also obtain the following theorem.

**Theorem 3.3.** Let  $p \geq 1$ ,  $0 < \alpha \leq \min\{s, t\}$ , and  $T \in \mathcal{H}_p^\alpha(\mathcal{H})$  such that  $D_T^\alpha = P - K$  with  $P, K \geq 0$ ,  $K \in \mathcal{C}_p(\mathcal{H})$ , and let  $\varepsilon \geq 0$  such that  $\alpha + \varepsilon \leq 2\alpha/(s + t)$ .

- (1) If  $s \geq 2\alpha$ , then  $\tilde{T}_{s,t} \in \mathcal{H}_{p/(\alpha+\varepsilon)}^{(\alpha+\varepsilon)}(\mathcal{H})$ .
- (2) If  $0 < s < 2\alpha$ , then  $\tilde{T}_{s,t} \in \mathcal{H}_{2\alpha p/(\alpha+\varepsilon)s}^{(\alpha+\varepsilon)}(\mathcal{H})$ .

*Proof.* The proof of Theorem 3.3 is similar to the proof of Theorem 3.1.  $\square$

**Corollary 3.4.** Let  $p \geq 1$ ,  $T \in \mathcal{H}_p^\alpha(\mathcal{H})$  such that  $D_T^\alpha = P - K$  with  $P, K \geq 0$ ,  $K \in \mathcal{C}_p(\mathcal{H})$ , and let  $\varepsilon \in (0, 1/2]$ .

- (1) If  $\alpha \in (0, 1/4]$ , then  $\tilde{T} \in \mathcal{H}_{p/(\alpha+\varepsilon)}^{(\alpha+\varepsilon)}(\mathcal{H})$ .
- (2) If  $\alpha \in (1/4, 1/2]$ , then  $\tilde{T} \in \mathcal{H}_{4\alpha p/(\alpha+\varepsilon)}^{(\alpha+\varepsilon)}(\mathcal{H})$ .

*Proof.* Put  $s = t = 1/2$  in Theorem 3.3.

- (1) When  $\alpha \in (0, 1/4]$ , we have  $s \geq 2\alpha$ . According to (1) of Theorem 3.3, then  $\tilde{T} \in \mathcal{H}_{p/(\alpha+\varepsilon)}^{(\alpha+\varepsilon)}(\mathcal{H})$ .
- (2) When  $\alpha \in (1/4, 1/2]$ , we have  $0 < s < 2\alpha$ . According to (2) of Theorem 3.3, then  $\tilde{T} \in \mathcal{H}_{4\alpha p/(\alpha+\varepsilon)}^{(\alpha+\varepsilon)}(\mathcal{H})$ .  $\square$

Next, we will study  $T^2$  of  $(\mathcal{C}_p, \alpha)$ -hyponormal operators. And first we will prove the following lemma.

**Lemma 3.5.** Let  $p \geq 1$ ,  $\alpha \in (0, 1]$ , and  $T \in \mathcal{H}_p^\alpha(\mathcal{H})$  such that  $D_T^\alpha = P + K$  with  $P \geq 0$ ,  $K \in \mathcal{C}_p(\mathcal{H})$ , and  $D_T^\alpha = P_1 - K_1$  with  $P_1 \geq 0$ ,  $K_1 \geq 0$ ,  $K_1 \in \mathcal{C}_p(\mathcal{H})$ . Then if  $|T|^{2\alpha} - P \geq 0$ , one has the following inequalities

- (1) There exists  $K' \in \mathcal{C}_{2p/\alpha}(\mathcal{H})$  such that  $(|T||T^*|^2|T|)^{\alpha/2} + K' \leq |T|^{2\alpha}$ .
- (2) There exists  $K'' \in \mathcal{C}_{2p/\alpha}(\mathcal{H})$  such that  $(|T^*||T|^2|T^*|)^{\alpha/2} + K'' \geq |T^*|^{2\alpha}$ .

*Proof.* Let  $\alpha, p$ , and  $T$  be as in the hypotheses and let  $T = U|T|$  be the polar decomposition of  $T$ . Then we have

$$D_T^\alpha = (T^*T)^\alpha - (TT^*)^\alpha = |T|^{2\alpha} - |T^*|^{2\alpha} = P + K, \quad (3.17)$$

with  $P \geq 0, K \in \mathcal{C}_p(\mathcal{H})$ .

$$D_T^\alpha = |T|^{2\alpha} - |T^*|^{2\alpha} = P_1 - K_1, \quad (3.18)$$

with and  $P_1 \geq 0, K_1 \geq 0$ , and  $K_1 \in \mathcal{C}_p(\mathcal{H})$ .

By (3.17), we have

$$A_1 = |T|^{2\alpha} \geq |T^*|^{2\alpha} + K = B_1 \geq 0. \quad (3.19)$$

And according to Lemma 1.2,

$$B_1^{1/\alpha} = \left(|T^*|^{2\alpha} + K\right)^{1/\alpha} = |T^*|^2 + K_2, \quad (3.20)$$

with  $K_2 \in \mathcal{C}_p(\mathcal{A})$ . Setting  $K_3 = |T|K_2|T|$ , by Theorem F we have

$$\begin{aligned} \left(|T||T^*|^2|T| + K_3\right)^{\alpha/2} &= \left\{|T|\left(|T^*|^2 + K_2\right)|T|\right\}^{\alpha/2} \\ &= \left(A_1^{1/2\alpha} B_1^{1/\alpha} A_1^{1/2\alpha}\right)^{\alpha/2} \\ &\leq A_1 \\ &= |T|^{2\alpha}. \end{aligned} \quad (3.21)$$

By (3.18), we have

$$A_2 = |T|^{2\alpha} + K_1 \geq |T^*|^{2\alpha} = B_2. \quad (3.22)$$

And according to Lemma 1.2,

$$A_2^{1/\alpha} = \left(|T|^{2\alpha} + K_1\right)^{1/\alpha} = |T|^2 + K_4, \quad (3.23)$$

with  $K_4 \in \mathcal{C}_p(\mathcal{A})$ . Setting  $K_5 = |T^*|K_4|T^*|$ , by Theorem F we have

$$\begin{aligned} \left(|T^*||T|^2|T^*| + K_5\right)^{\alpha/2} &= \left\{|T^*|\left(|T|^2 + K_4\right)|T^*|\right\}^{\alpha/2} \\ &= \left(B_2^{1/2\alpha} A_2^{1/\alpha} B_2^{1/2\alpha}\right)^{\alpha/2} \\ &\geq B_2 \\ &= |T^*|^{2\alpha}. \end{aligned} \quad (3.24)$$

On the other hand, by Lemma 1.1,

$$\begin{aligned} \left(|T||T^*|^2|T| + K_3\right)^{\alpha/2} &= \left(|T||T^*|^2|T|\right)^{\alpha/2} + K', \\ \left(|T^*||T|^2|T^*| + K_5\right)^{\alpha/2} &= \left(|T^*||T|^2|T^*|\right)^{\alpha/2} + K'', \end{aligned} \quad (3.25)$$

with  $K', K'' \in \mathcal{C}_{2p/\alpha}(\mathcal{A})$ .



Then by (3.21) and (3.24), we obtain

$$\left(|T||T^*|^2|T|\right)^{\alpha/2} + K' \leq |T|^{2\alpha}, \quad \left(|T^*||T|^2|T^*|\right)^{\alpha/2} + K'' \geq |T^*|^{2\alpha}, \quad (3.26)$$

with  $K', K'' \in \mathcal{C}_{2p/\alpha}(\mathcal{H})$ .  $\square$

**Theorem 3.6.** Let  $p \geq 1, \alpha \in (0, 1]$ , and  $T \in \mathcal{H}_p^\alpha(\mathcal{H})$  such that  $D_T^\alpha = P + K$  with  $P \geq 0, K \in \mathcal{C}_p(\mathcal{H})$ , and  $D_T^\alpha = P_1 - K_1$  with  $P_1 \geq 0, K_1 \geq 0$ , and  $K_1 \in \mathcal{C}_p(\mathcal{H})$ . Then if  $|T|^{2\alpha} - P \geq 0$ , one has  $T^2 \in \mathcal{H}_{2p/\alpha}^{\alpha/2}(\mathcal{H})$ .

*Proof.* Let  $\alpha, p$ , and  $T$  be as in the hypotheses. We may assume that  $T = U|T|$  with  $U$  being unitary. Then obviously,

$$\left\{T^2(T^2)^*\right\}^{\alpha/2} = U\left(|T||T^*|^2|T|\right)^{\alpha/2}U^*, \quad (3.27)$$

$$\left\{(T^2)^*T^2\right\}^{\alpha/2} = \left(|T|U^*|T|^2U|T|\right)^{\alpha/2} = U^*\left(|T^*||T|^2|T^*|\right)^{\alpha/2}U. \quad (3.28)$$

By Lemma 3.5, there exists  $K', K'' \in \mathcal{C}_{2p/\alpha}(\mathcal{H})$  such that

$$\left(|T||T^*|^2|T|\right)^{\alpha/2} + K' \leq |T|^{2\alpha}, \quad (3.29)$$

$$\left(|T^*||T|^2|T^*|\right)^{\alpha/2} + K'' \geq |T^*|^{2\alpha}. \quad (3.30)$$

Multiplying (3.29) by  $U$  to the left and by  $U^*$  to the right, we obtain

$$U\left(|T||T^*|^2|T|\right)^{\alpha/2}U^* + UK'U^* \leq U|T|^{2\alpha}U^* = |T^*|^{2\alpha}. \quad (3.31)$$

Multiplying (3.30) by  $U^*$  to the left and by  $U$  to the right, we obtain

$$U^*\left(|T^*||T|^2|T^*|\right)^{\alpha/2}U + U^*K''U \geq U^*|T^*|^{2\alpha}U = |T|^{2\alpha}. \quad (3.32)$$

By (3.27) and (3.31), we have

$$\left\{T^2(T^2)^*\right\}^{\alpha/2} + UK'U^* \leq |T^*|^{2\alpha}. \quad (3.33)$$

By (3.28) and (3.32), we have

$$\left\{(T^2)^*T^2\right\}^{\alpha/2} + U^*K''U \geq |T|^{2\alpha}. \quad (3.34)$$

Setting  $K_2 = UK'U^* - U^*K''U$ ,  $K_2 \in \mathcal{C}_{2p/\alpha}(\mathcal{H})$ , we have

$$\left\{ \left( T^2 \right)^* T^2 \right\}^{\alpha/2} - \left\{ T^2 \left( T^2 \right)^* \right\}^{\alpha/2} \geq |T|^{2\alpha} - |T^*|^{2\alpha} + K_2. \quad (3.35)$$

Therefore, for  $K_3 = K + K_2$ ,  $K_3 \in \mathcal{C}_{2p/\alpha}(\mathcal{H})$ , we have

$$\left\{ \left( T^2 \right)^* T^2 \right\}^{\alpha/2} - \left\{ T^2 \left( T^2 \right)^* \right\}^{\alpha/2} \geq P + K_3. \quad (3.36)$$

Then the proof of Theorem 3.6 is finished.  $\square$

## Acknowledgment

The authors would like to express their cordial gratitude to the referee for his kind comments.

## References

- [1] V. Lauric, “ $(C_p, \alpha)$ -hyponormal operators and trace-class self-commutators with trace zero,” *Proceedings of the American Mathematical Society*, vol. 137, no. 3, pp. 945–953, 2009.
- [2] R. Schatten, *Norm Ideals of Completely Continuous Operators*, Ergebnisse der Mathematik und ihrer Grenzgebiete. N. F., Heft 27, Springer, Berlin, Germany, 1960.
- [3] T. Furuta, “ $A \geq B \geq 0$  assures  $(B^r A^p B^r)^{1/q} \geq B^{(p+2r)/q}$  for  $r \geq 0$ ,  $p \geq 0$ ,  $q \geq 1$  with  $(1+2r)q \geq p+2r$ ,” *Proceedings of the American Mathematical Society*, vol. 101, no. 1, pp. 85–88, 1987.
- [4] E. Heinz, “Beiträge zur Störungstheorie der Spektralzerlegung,” *Mathematische Annalen*, vol. 123, pp. 415–438, 1951 (German).
- [5] K. Löwner, “Über monotone Matrixfunktionen,” *Mathematische Zeitschrift*, vol. 38, no. 1, pp. 177–216, 1934 (German).
- [6] A. Aluthge, “On  $p$ -hyponormal operators for  $0 < p < 1$ ,” *Integral Equations and Operator Theory*, vol. 13, no. 3, pp. 307–315, 1990.
- [7] T. Furuta and M. Yanagida, “Further extensions of Aluthge transformation on  $p$ -hyponormal operators,” *Integral Equations and Operator Theory*, vol. 29, no. 1, pp. 122–125, 1997.
- [8] T. Ando, “Aluthge transforms and the convex hull of the eigenvalues of a matrix,” *Linear and Multilinear Algebra*, vol. 52, no. 3-4, pp. 281–292, 2004.
- [9] J. Antezana, P. Massey, and D. Stojanoff, “ $\lambda$ -Aluthge transforms and Schatten ideals,” *Linear Algebra and Its Applications*, vol. 405, pp. 177–199, 2005.
- [10] T. Ando and T. Yamazaki, “The iterated Aluthge transforms of a 2-by-2 matrix converge,” *Linear Algebra and Its Applications*, vol. 375, pp. 299–309, 2003.
- [11] T. Furuta and M. Yanagida, “On powers of  $p$ -hyponormal and log-hyponormal operators,” *Journal of Inequalities and Applications*, vol. 5, no. 4, pp. 367–380, 2000.
- [12] M. Ito, “Generalizations of the results on powers of  $p$ -hyponormal operators,” *Journal of Inequalities and Applications*, vol. 6, no. 1, pp. 1–15, 2001.
- [13] I. B. Jung, E. Ko, and C. Pearcy, “Aluthge transforms of operators,” *Integral Equations and Operator Theory*, vol. 37, no. 4, pp. 437–448, 2000.
- [14] K. Tanahashi, “On log-hyponormal operators,” *Integral Equations and Operator Theory*, vol. 34, no. 3, pp. 364–372, 1999.
- [15] J. Yuan and Z. Gao, “Spectrum of class  $wF(p, r, q)$  operators,” *Journal of Inequalities and Applications*, vol. 2007, Article ID 27195, 10 pages, 2007.