

Research Article

A New Method for Solving Monotone Generalized Variational Inequalities

Pham Ngoc Anh and Jong Kyu Kim

Department of Mathematics, Kyungnam University, Masan, Kyungnam 631-701, Republic of Korea

Correspondence should be addressed to Jong Kyu Kim, jongkyuk@kyungnam.ac.kr

Received 11 May 2010; Revised 27 August 2010; Accepted 4 October 2010

Academic Editor: Siegfried Carl

Copyright © 2010 P. N. Anh and J. K. Kim. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We suggest new dual algorithms and iterative methods for solving monotone generalized variational inequalities. Instead of working on the primal space, this method performs a dual step on the dual space by using the dual gap function. Under the suitable conditions, we prove the convergence of the proposed algorithms and estimate their complexity to reach an ε -solution. Some preliminary computational results are reported.

1. Introduction

Let C be a convex subset of the real Euclidean space \mathbb{R}^n , F be a continuous mapping from C into \mathbb{R}^n , and φ be a lower semicontinuous convex function from C into \mathbb{R} . We say that a point x^* is a solution of the following generalized variational inequality if it satisfies

$$\langle F(x^*), x - x^* \rangle + \varphi(x) - \varphi(x^*) \geq 0, \quad \forall x \in C, \quad (\text{GVI})$$

where $\langle \cdot, \cdot \rangle$ denotes the standard dot product in \mathbb{R}^n .

Associated with the problem (GVI), the dual form of this is expressed as following which is to find $y^* \in C$ such that

$$\langle F(x), x - y^* \rangle + \varphi(x) - \varphi(y^*) \geq 0, \quad \forall x \in C. \quad (\text{DGVI})$$

In recent years, this generalized variational inequalities become an attractive field for many researchers and have many important applications in electricity markets, transportations, economics, and nonlinear analysis (see [1–9]).

It is well known that the interior quadratic and dual technique are powerful tools for analyzing and solving the optimization problems (see [10–16]). Recently these techniques have been used to develop proximal iterative algorithm for variational inequalities (see [17–22]).

In addition Nesterov [23] introduced a dual extrapolation method for solving variational inequalities. Instead of working on the primal space, this method performs a dual step on the dual space.

In this paper we extend results in [23] to the generalized variational inequality problem (GVI) in the dual space. In the first approach, a gap function $g(x)$ is constructed such that $g(x) \geq 0$, for all $x^* \in C$ and $g(x^*) = 0$ if and only if x^* solves (GVI). Namely, we first develop a convergent algorithm for (GVI) with F being monotone function satisfying a certain Lipschitz type condition on C . Next, in order to avoid the Lipschitz condition we will show how to find a regularization parameter at every iteration k such that the sequence x^k converges to a solution of (GVI).

The remaining part of the paper is organized as follows. In Section 2, we present two convergent algorithms for monotone and generalized variational inequality problems with Lipschitzian condition and without Lipschitzian condition. Section 3 deals with some preliminary results of the proposed methods.

2. Preliminaries

First, let us recall the well-known concepts of monotonicity that will be used in the sequel (see [24]).

Definition 2.1. Let C be a convex set in \mathbb{R}^n , and $F : C \rightarrow \mathbb{R}^n$. The function F is said to be

(i) pseudomonotone on C if

$$\langle F(y), x - y \rangle \geq 0 \implies \langle F(x), x - y \rangle \geq 0, \quad \forall x, y \in C, \quad (2.1)$$

(ii) monotone on C if for each $x, y \in C$,

$$\langle F(x) - F(y), x - y \rangle \geq 0, \quad (2.2)$$

(iii) strongly monotone on C with constant $\beta > 0$ if for each $x, y \in C$,

$$\langle F(x) - F(y), x - y \rangle \geq \beta \|x - y\|^2, \quad (2.3)$$

(iv) Lipschitz with constant $L > 0$ on C (shortly L -Lipschitz), if

$$\|F(x) - F(y)\| \leq L \|x - y\|, \quad \forall x, y \in C. \quad (2.4)$$

Note that when φ is differentiable on some open set containing C , then, since φ is lower semicontinuous proper convex, the generalized variational inequality (GVI) is equivalent to the following variational inequalities (see [25, 26]):

Find $x^* \in C$ such that

$$\langle F(x^*) + \nabla\varphi(x^*), x - x^* \rangle \geq 0, \quad \forall x \in C. \quad (2.5)$$

Throughout this paper, we assume that:

- (A₁) the interior set of C , $\text{int } C$ is nonempty,
- (A₂) the set C is bounded,
- (A₃) F is upper semicontinuous on C , and φ is proper, closed convex and subdifferentiable on C ,
- (A₄) F is monotone on C .

In special case $\varphi = 0$, problem (GVI) can be written by the following.

Find $x^* \in C$ such that

$$\langle F(x^*), x - x^* \rangle \geq 0, \quad \forall x \in C. \quad (\text{VI})$$

It is well known that the problem (VI) can be formulated as finding the zero points of the operator $T(x) = F(x) + N_C(x)$, where

$$N_C(x) = \begin{cases} \{y \in C : \langle y, z - x \rangle \leq 0, \forall z \in C\}, & \text{if } x \in C, \\ \emptyset, & \text{otherwise.} \end{cases} \quad (2.6)$$

The dual gap function of problem (GVI) is defined as follows:

$$g(x) := \sup\{\langle F(y), x - y \rangle + \varphi(x) - \varphi(y) \mid y \in C\}. \quad (2.7)$$

The following lemma gives two basic properties of the dual gap function (2.7) whose proof can be found, for instance, in [6].

Lemma 2.2. *The function g is a gap function of (GVI), that is,*

- (i) $g(x) \geq 0$ for all $x \in C$,
- (ii) $x^* \in C$ and $g(x^*) = 0$ if and only if x^* is a solution to (DGVI). Moreover, if F is pseudomonotone then x^* is a solution to (DGVI) if and only if it is a solution to (GVI).

The problem $\sup\{\langle F(y), x - y \rangle + \varphi(x) - \varphi(y) \mid y \in C\}$ may not be solvable and the dual gap function g may not be well-defined. Instead of using gap function g , we consider a truncated dual gap function g_R . Suppose that $\bar{x} \in \text{int } C$ fixed and $R > 0$. The truncated dual gap function is defined as follows:

$$g_R(x) := \max\{\langle F(y), x - y \rangle + \varphi(x) - \varphi(y) \mid y \in C, \|y - \bar{x}\| \leq R\}. \quad (2.8)$$

For the following consideration, we define $B_R(\bar{x}) := \{y \in \mathbb{R}^n \mid \|y - \bar{x}\| \leq R\}$ as a closed ball in \mathbb{R}^n centered at \bar{x} and radius R , and $C_R := C \cap B_R(\bar{x})$. The following lemma gives some properties for g_R .

Lemma 2.3. *Under assumptions (A_1) – (A_4) , the following properties hold.*

- (i) *The function $g_R(\cdot)$ is well-defined and convex on C .*
- (ii) *If a point $x^* \in C \cap B_R(\bar{x})$ is a solution to (DGVI) then $g_R(x^*) = 0$.*
- (iii) *If there exists $x^0 \in C$ such that $g_R(x^0) = 0$ and $\|x^0 - \bar{x}\| < R$, and F is pseudomonotone, then x^0 is a solution to (DGVI) (and also (GVI)).*

Proof. (i) Note that $\langle F(y), x - y \rangle + \varphi(x) - \varphi(y)$ is upper semicontinuous on C for $x \in C$ and $B_R(\bar{x})$ is bounded. Therefore, the supremum exists which means that g_R is well-defined. Moreover, since φ is convex on C and g is the supremum of a parametric family of convex functions (which depends on the parameter x), then g_R is convex on C .

(ii) By definition, it is easy to see that $g_R(x) \geq 0$ for all $x \in C \cap B_R(\bar{x})$. Let x^* be a solution of (DGVI) and $x^* \in B_R(\bar{x})$. Then we have

$$\langle F(y), x^* - y \rangle + \varphi(x^*) - \varphi(y) \leq 0 \quad \forall y \in C. \quad (2.9)$$

In particular, we have

$$\langle F(y), x^* - y \rangle + \varphi(y) - \varphi(x^*) \leq 0 \quad (2.10)$$

for all $y \in C \cap B_R(\bar{x})$. Thus

$$g_R(x^*) = \sup\{\langle F(y), x^* - y \rangle + \varphi(x^*) - \varphi(y) \mid y \in C \cap B_R(\bar{x})\} \leq 0, \quad (2.11)$$

this implies $g_R(x^*) = 0$.

(iii) For some $x^0 \in C \cap \text{int } B_R(\bar{x})$, $g_R(x^0) = 0$ means that x is a solution to (DGVI) restricted to $C \cap \text{int } B_R(\bar{x})$. Since F is pseudomonotone, x^0 is also a solution to (GVI) restricted to $C \cap B_R(\bar{x})$. Since $x^0 \in \text{int } B_R(\bar{x})$, for any $y \in C$, we can choose $\lambda > 0$ sufficiently small such that

$$y_\lambda := x^0 + \lambda(y - x^0) \in C \cap B_R(\bar{x}), \quad (2.12)$$

$$\begin{aligned} 0 &\leq \langle F(x^0), y_\lambda - x^0 \rangle + \varphi(y_\lambda) - \varphi(x^0) \\ &= \langle F(x^0), x^0 + \lambda(y - x^0) - x^0 \rangle + \varphi(x^0 + \lambda(y - x^0)) - \varphi(x^0) \\ &\leq \lambda \langle F(x^0), y - x^0 \rangle + \lambda \varphi(y) + (1 - \lambda) \varphi(x^0) - \varphi(x^0) \\ &= \lambda (\langle F(x^0), y - x^0 \rangle + \varphi(y) - \varphi(x^0)), \end{aligned} \quad (2.13)$$

where (2.13) follows from the convexity of $\varphi(\cdot)$. Since $\lambda > 0$, dividing this inequality by λ , we obtain that x^0 is a solution to (GVI) on C . Since F is pseudomonotone, x^0 is also a solution to (DGVI). \square

Let $C \subseteq \mathbb{R}^n$ be a nonempty, closed convex set and $x \in \mathbb{R}^n$. Let us denote $d_C(x)$ the Euclidean distance from x to C and $Pr_C(x)$ the point attained this distance, that is,

$$d_C(x) := \min_{y \in C} \|y - x\|, \quad Pr_C(x) := \arg \min_{y \in C} \|y - x\|. \quad (2.14)$$

As usual, Pr_C is referred to the Euclidean projection onto the convex set C . It is well-known that Pr_C is a nonexpansive and co-coercive operator on C (see [27, 28]).

The following lemma gives a tool for the next discussion.

Lemma 2.4. *For any $x, y, z \in \mathbb{R}^n$ and for any $\beta > 0$, the function d_C and the mapping Pr_C defined by (2.14) satisfy*

$$\langle Pr_C(x) - x, y - Pr_C(x) \rangle \geq 0, \quad \forall y \in C, \quad (2.15)$$

$$d_C^2(x + y) \geq d_C^2(x) + d_C^2(Pr_C(x) + y) - 2\langle y, Pr_C(x) - x \rangle, \quad (2.16)$$

$$\left\| x - Pr_C\left(x + \frac{1}{\beta}y\right) \right\|^2 \leq \frac{1}{\beta^2} \|y\|^2 - d_C^2\left(x + \frac{1}{\beta}y\right), \quad \forall x \in C. \quad (2.17)$$

Proof. Inequality (2.15) is obvious from the property of the projection Pr_C (see [27]). Now, we prove the inequality (2.16). For any $v \in C$, applying (2.15) we have

$$\begin{aligned} \|v - (x + y)\|^2 &= \|v - (Pr_C(x) + y) + (Pr_C(x) - x)\|^2 \\ &= \|v - (Pr_C(x) + y)\|^2 + 2\langle v - (Pr_C(x) + y), Pr_C(x) - x \rangle + \|Pr_C(x) - x\|^2 \\ &= \|v - (Pr_C(x) + y)\|^2 + 2\langle Pr_C(x) - x, v - Pr_C(x) \rangle \\ &\quad - 2\langle y, Pr_C(x) - x \rangle + \|Pr_C(x) - x\|^2 \\ &\geq \|v - (Pr_C(x) + y)\|^2 - 2\langle y, Pr_C(x) - x \rangle + \|Pr_C(x) - x\|^2. \end{aligned} \quad (2.18)$$

Using the definition of $d_C(\cdot)$ and noting that $d_C^2(x) = \|Pr_C(x) - x\|^2$ and taking minimum with respect to $v \in C$ in (2.18), then we have

$$d_C^2(x + y) \geq d_C^2(Pr_C(x) + y) + d_C^2(x) - 2\langle y, Pr_C(x) - x \rangle, \quad (2.19)$$

which proves (2.16).

From the definition of d_C , we have

$$\begin{aligned}
 d_C^2\left(x + \frac{1}{\beta}y\right) &= \left\|Pr_C\left(x + \frac{1}{\beta}y\right) - x - \frac{1}{\beta}y\right\|^2 \\
 &= \frac{1}{\beta^2}\|y^2\| - \left\|x + \frac{1}{\beta}y - Pr_C\left(x + \frac{1}{\beta}y\right) - \left(x - Pr_C\left(x + \frac{1}{\beta}y\right)\right)\right\|^2 \\
 &\quad + \left\|Pr_C\left(x + \frac{1}{\beta}y\right) - x - \frac{1}{\beta}y\right\|^2 \\
 &= \frac{1}{\beta^2}\|y^2\| - \left\|x - Pr_C\left(x + \frac{1}{\beta}y\right)\right\|^2 \\
 &\quad + 2\left\langle x + \frac{1}{\beta}y - Pr_C\left(x + \frac{1}{\beta}y\right), x - Pr_C\left(x + \frac{1}{\beta}y\right)\right\rangle.
 \end{aligned} \tag{2.20}$$

Since $x \in C$, applying (2.15) with $Pr_C(x + (1/\beta)y)$ instead of $Pr_C(x)$ and $y = x$ for (2.20), we obtain the last inequality in Lemma 2.4. \square

For a given integer number $m \geq 0$, we consider a finite sequence of arbitrary points $\{x^k\}_{k=0}^m \subset C$, a finite sequence of arbitrary points $\{w^k\}_{k=0}^m \subset \mathbb{R}^n$ and a finite positive sequence $\{\lambda_k\}_{k=0}^m \subseteq (0, +\infty)$. Let us define

$$\bar{w}^m = \sum_{k=0}^m \lambda_k w^k, \quad \bar{\lambda}_m = \sum_{k=0}^m \lambda_k, \quad \bar{x}^m = \frac{1}{\bar{\lambda}_m} \sum_{k=0}^m \lambda_k x^k. \tag{2.21}$$

Then upper bound of the dual gap function g_R is estimated in the following lemma.

Lemma 2.5. *Suppose that Assumptions (A₁)–(A₄) are satisfied and*

$$w^k \in -F(x^k) - \partial\varphi(x^k). \tag{2.22}$$

Then, for any $\beta > 0$,

- (i) $\max\{\langle w, y - \bar{x} \rangle \mid y \in C_R\} \leq (1/2\beta)\|w\|^2 - (\beta/2)d_C^2(\bar{x} + (1/\beta)w) + \beta R^2/2$, for all $\bar{x} \in C$, $w \in \mathbb{R}^n$.
- (ii) $g_R(\bar{x}^m) \leq (1/\bar{\lambda}_m)(\sum_{k=0}^m \lambda_k \langle w^k, \bar{x} - x^k \rangle + (1/2\beta)\|\bar{w}^m\|^2 - (\beta/2)d_C^2(\bar{x} + (1/\beta)\bar{w}^m) + \beta R^2/2)$.

Proof. (i) We define $L(x, \rho) = \langle w, y - \bar{x} \rangle + (\rho/2)(R^2 - \|y - \bar{x}\|^2)$ as the Lagrange function of the maximizing problem $\max\{\langle w, y - \bar{x} \rangle \mid y \in C_R\}$. Using duality theory in convex optimization, then we have

$$\begin{aligned}
 \max\{\langle w, y - \bar{x} \rangle \mid y \in C_R\} &= \max\{\langle w, y - \bar{x} \rangle \mid y \in C, \|y - \bar{x}\|^2 \leq R^2\} \\
 &= \max_{y \in C} \min_{\rho \geq 0} \left\{ \langle w, y - \bar{x} \rangle + \rho \left(R^2 - \|y - \bar{x}\|^2 \right) \right\} \\
 &= \min_{\rho \geq 0} \left\{ \max_{y \in C} \left\{ \langle w, y - \bar{x} \rangle - \frac{\rho}{2} \|y - \bar{x}\|^2 \right\} + \frac{\rho}{2} R^2 \right\} \\
 &= \min_{\rho \geq 0} \left\{ \frac{1}{2\rho} \max_{y \in C} \left\{ \|w\|^2 - \rho^2 \left\| y - \bar{x} - \frac{1}{\rho} w \right\|^2 \right\} + \frac{\rho}{2} R^2 \right\} \quad (2.23) \\
 &\leq \frac{1}{2\beta} \left[\|w\|^2 - \beta^2 \min_{y \in C} \left\| y - \bar{x} - \frac{1}{\beta} w \right\|^2 \right] + \frac{\beta R^2}{2} \\
 &= \frac{1}{2\beta} \|w\|^2 - \frac{\beta}{2} d_C^2 \left(\bar{x} + \frac{1}{\beta} w \right) + \frac{\beta R^2}{2}.
 \end{aligned}$$

(ii) From the monotonicity of F and (2.22), we have

$$\begin{aligned}
 \sum_{k=0}^m \lambda_k \left(\langle F(y), x^k - y \rangle + \varphi(x^k) - \varphi(y) \right) &\leq - \sum_{k=0}^m \lambda_k \left(\langle F(x^k), y - x^k \rangle + \varphi(y) - \varphi(x^k) \right) \\
 &\leq \sum_{k=0}^m \lambda_k \langle w^k, y - x^k \rangle \\
 &\leq \sum_{k=0}^m \lambda_k \langle w^k, y - \bar{x} \rangle + \sum_{k=0}^m \lambda_k \langle w^k, \bar{x} - x^k \rangle \\
 &= \langle \bar{w}^m, y - \bar{x} \rangle + \sum_{k=0}^m \lambda_k \langle w^k, \bar{x} - x^k \rangle.
 \end{aligned} \quad (2.24)$$

Combining (2.24), Lemma 2.5(i) and

$$\begin{aligned}
 g_R(\bar{x}^m) &= \max\{\langle F(y), \bar{x}^m - y \rangle + \varphi(\bar{x}^m) - \varphi(y) \mid y \in C_R\} \\
 &= \max\left\{ \left\langle F(y), \frac{1}{\lambda_m} \sum_{k=0}^m \lambda_k x^k - y \right\rangle + \varphi\left(\frac{1}{\lambda_m} \sum_{k=0}^m \lambda_k x^k \right) - \varphi(y) \mid y \in C_R \right\} \\
 &\leq \max\left\{ \frac{1}{\lambda_m} \sum_{k=0}^m \lambda_k \left(\langle F(y), x^k - y \rangle + \varphi(x^k) - \varphi(y) \right) \mid y \in C_R \right\} \\
 &= \frac{1}{\lambda_m} \max\left\{ \sum_{k=0}^m \lambda_k \left(\langle F(y), x^k - y \rangle + \varphi(x^k) - \varphi(y) \right) \mid y \in C_R \right\},
 \end{aligned} \quad (2.25)$$

we get

$$\begin{aligned} g_R(\bar{x}^m) &\leq \frac{1}{\lambda_m} \max\{\langle \bar{w}^m, y - \bar{x} \rangle \mid y \in C_R\} + \sum_{k=0}^m \lambda_k \langle w^k, \bar{x} - x^k \rangle \\ &\leq \frac{1}{\lambda_m} \left(\frac{1}{2\beta} \|\bar{w}^m\|^2 - \frac{\beta}{2} d_C^2 \left(\bar{x} + \frac{1}{\beta} \bar{w}^m \right) + \frac{\beta R^2}{2} + \sum_{k=0}^m \lambda_k \langle w^k, \bar{x} - x^k \rangle \right). \end{aligned} \quad (2.26)$$

□

3. Dual Algorithms

Now, we are going to build the dual interior proximal step for solving (GVI). The main idea is to construct a sequence $\{\bar{x}^k\}$ such that the sequence $g_R(\bar{x}^k)$ tends to 0 as $k \rightarrow \infty$. By virtue of Lemma 2.5, we can check whether \bar{x}^k is an ε -solution to (GVI) or not.

The dual interior proximal step $(u^k, x^k, \bar{w}^k, w^k)$ at the iteration $k \geq 0$ is generated by using the following scheme:

$$\begin{aligned} u^k &:= \text{Pr}_C \left(\bar{x} + \frac{1}{\beta} \bar{w}^{k-1} \right), \\ x^k &:= \arg \min \left\{ \langle F(u^k), y - u^k \rangle + \varphi(y) - \varphi(u^k) + \frac{\beta \rho_k}{2} \|y - u^k\|^2 \mid y \in C \right\}, \\ \bar{w}^k &:= \bar{w}^{k-1} + \frac{1}{\rho_k} w^k, \end{aligned} \quad (3.1)$$

where $\rho_k > 0$ and $\beta > 0$ are given parameters, $w^k \in \mathbb{R}^n$ is the solution to (2.22).

The following lemma shows an important property of the sequence (u^k, x^k, s^k, w^k) .

Lemma 3.1. *The sequence $(u^k, x^k, \bar{w}^k, w^k)$ generated by scheme (3.1) satisfies*

$$\begin{aligned} d_C^2 \left(\bar{x} + \frac{1}{\beta} \bar{w}^k \right) &\geq d_C^2 \left(\bar{x} + \frac{1}{\beta} \bar{w}^{k-1} \right) + \|x^k - u^k\|^2 + \|\pi_C^k - x^k\|^2 \\ &\quad - \frac{2}{\beta \rho_k} \langle \pi_C^k - x^k, \xi^k + w^k \rangle \\ &\quad + \frac{1}{\beta^2 \rho_k^2} \|w^k\|^2 + \frac{2}{\beta \rho_k} \left\langle w^k, \bar{x} - x^k + \frac{1}{\beta} \bar{w}^{k-1} \right\rangle, \end{aligned} \quad (3.2)$$

where $\eta^k \in \partial\varphi(x^k)$, $\xi^k = \eta^k + F(u^k)$ and $\pi_C^k = \text{Pr}_C(x^k + (1/\beta\rho_k)(\xi^k + w^k))$. As a consequence, we have

$$\begin{aligned} d_C^2 \left(\bar{x} + \frac{1}{\beta} \bar{w}^k \right) - d_C^2 \left(\bar{x} + \frac{1}{\beta} \bar{w}^{k-1} \right) &\geq \frac{2}{\beta \rho_k} \langle w^k, \bar{x} - x^k \rangle + \frac{1}{\beta^2} \|\bar{w}^k\|^2 - \frac{1}{\beta^2} \|\bar{w}^{k-1}\|^2 \\ &\quad - \frac{1}{\beta^2 \rho_k^2} \|\xi^k + w^k\|^2. \end{aligned} \quad (3.3)$$

Proof. We replace x by $x + (1/\beta)y$ and y by $(1/\beta)z$ into (2.16) to obtain

$$\begin{aligned} d_C^2\left(x + \frac{1}{\beta}(y+z)\right) &\geq d_C^2\left(x + \frac{1}{\beta}y\right) + d_C^2\left(\text{Pr}_C\left(x + \frac{1}{\beta}y\right) + \frac{1}{\beta}z\right) \\ &\quad - \frac{2}{\beta}\left\langle z, \text{Pr}_C\left(x + \frac{1}{\beta}y\right) - \left(x + \frac{1}{\beta}y\right) \right\rangle. \end{aligned} \quad (3.4)$$

Using the inequality (3.4) with $x = \bar{x}$, $y = \bar{w}^{k-1}$, $z = (1/\rho_k)w^k$ and noting that $u^k = \text{Pr}_C(\bar{x} + (1/\beta)\bar{w}^{k-1})$, we get

$$\begin{aligned} d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1} + \frac{1}{\beta\rho_k}w^k\right) &\geq d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) + d_C^2\left(\text{Pr}_C\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) + \frac{1}{\beta\rho_k}w^k\right) \\ &\quad - \frac{2}{\beta\rho_k}\left\langle w^k, \text{Pr}_C\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) - \bar{x} - \frac{1}{\beta}\bar{w}^{k-1} \right\rangle. \end{aligned} \quad (3.5)$$

This implies that

$$d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^k\right) \geq d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) + d_C^2\left(u^k + \frac{1}{\beta\rho_k}w^k\right) - \frac{2}{\beta\rho_k}\left\langle w^k, u^k - \bar{x} - \frac{1}{\beta}\bar{w}^{k-1} \right\rangle. \quad (3.6)$$

From the subdifferentiability of the convex function φ to scheme (3.1), using the first-order necessary optimality condition, we have

$$\left\langle F(u^k) + \eta^k + \beta\rho_k(x^k - u^k), v - x^k \right\rangle \geq 0, \quad \forall v \in C, \quad (3.7)$$

for all $\eta^k \in \partial\varphi(x^k)$. This inequality implies that

$$x^k = \text{Pr}_C\left(u^k - \frac{1}{\beta\rho_k}\xi^k\right), \quad (3.8)$$

where $\xi^k = \eta^k + F(u^k)$.

We apply inequality (3.4) with $x = u^k$, $y = -(1/\rho_k)\xi^k$ and $z = (1/\rho_k)(\xi^k + w^k)$ and using (3.8) to obtain

$$\begin{aligned}
 d_C^2\left(u^k + \frac{1}{\beta\rho_k}w^k\right) &\geq d_C^2\left(u^k - \frac{1}{\beta\rho_k}\xi^k\right) + d_C^2\left(x^k + \frac{1}{\beta\rho_k}(\xi^k + w^k)\right) \\
 &\quad - \frac{2}{\beta\rho_k}\left\langle \xi^k + w^k, x^k - u^k + \frac{1}{\beta\rho_k}\xi^k \right\rangle \\
 &= \left\| Pr_C\left(u^k - \frac{1}{\beta\rho_k}\xi^k\right) - u^k + \frac{1}{\beta\rho_k}\xi^k \right\|^2 \\
 &\quad + d_C^2\left(x^k + \frac{1}{\beta\rho_k}(\xi^k + w^k)\right) - \frac{2}{\beta\rho_k}\left\langle \xi^k + w^k, x^k - u^k + \frac{1}{\beta\rho_k}\xi^k \right\rangle \\
 &= \left\| x^k - u^k + \frac{1}{\beta\rho_k}\xi^k \right\|^2 + d_C^2\left(x^k + \frac{1}{\beta\rho_k}(\xi^k + w^k)\right) \\
 &\quad + \frac{2}{\beta\rho_k}\left\langle \xi^k + w^k, u^k - \frac{1}{\beta\rho_k}\xi^k - x^k \right\rangle \\
 &= \left\| x^k - u^k \right\|^2 + \frac{1}{\beta^2\rho_k^2}\left\| \xi^k \right\|^2 + \frac{2}{\beta\rho_k}\left\langle \xi^k, x^k - u^k \right\rangle \\
 &\quad + d_C^2\left(x^k + \frac{1}{\beta\rho_k}(\xi^k + w^k)\right) + \frac{2}{\beta\rho_k}\left\langle \xi^k + w^k, u^k - \frac{1}{\beta\rho_k}\xi^k - x^k \right\rangle.
 \end{aligned} \tag{3.9}$$

Combine this inequality and (3.6), we get

$$\begin{aligned}
 d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^k\right) &\geq d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) - \frac{2}{\beta\rho_k}\left\langle w^k, u^k - \bar{x} - \frac{1}{\beta}\bar{w}^{k-1} \right\rangle \\
 &\quad + \left\| x^k - u^k \right\|^2 + \frac{1}{\beta^2\rho_k^2}\left\| \xi^k \right\|^2 + \frac{2}{\beta\rho_k}\left\langle \xi^k, x^k - u^k \right\rangle \\
 &\quad + d_C^2\left(x^k + \frac{1}{\beta\rho_k}(\xi^k + w^k)\right) + \frac{2}{\beta\rho_k}\left\langle \xi^k + w^k, u^k - \frac{1}{\beta\rho_k}\xi^k - x^k \right\rangle.
 \end{aligned} \tag{3.10}$$

On the other hand, if we denote $\pi_C^k = Pr_C(x^k + (1/\beta\rho_k)(\xi^k + w^k))$, then it follows that

$$\begin{aligned}
 d_C^2\left(x^k + \frac{1}{\beta\rho_k}(\xi^k + w^k)\right) &= \left\| \pi_C^k - x^k - \frac{1}{\beta\rho_k}(\xi^k + w^k) \right\|^2 \\
 &= \left\| \pi_C^k - x^k \right\|^2 - \frac{2}{\beta\rho_k}\left\langle \pi_C^k - x^k, \xi^k + w^k \right\rangle + \frac{1}{\beta^2\rho_k^2}\left\| \xi^k + w^k \right\|^2.
 \end{aligned} \tag{3.11}$$

Combine (3.10) and (3.11), we get

$$\begin{aligned}
 d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^k\right) &\geq d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) + \|x^k - u^k\|^2 + \|\pi_C^k - x^k\|^2 \\
 &\quad - \frac{2}{\beta\rho_k} \langle \pi_C^k - x^k, \xi^k + w^k \rangle \\
 &\quad + \frac{1}{\beta^2\rho_k^2} \|w^k\|^2 + \frac{2}{\beta\rho_k} \left\langle w^k, \bar{x} - x^k + \frac{1}{\beta}\bar{w}^{k-1} \right\rangle,
 \end{aligned}
 \tag{3.12}$$

which proves (3.2).

On the other hand, from (3.9) we have

$$\begin{aligned}
 d_C^2\left(u^k + \frac{1}{\beta\rho_k}w^k\right) &\geq \|x^k - u^k\|^2 + \frac{1}{\beta^2\rho_k^2} \|\xi^k\|^2 + \frac{2}{\beta\rho_k} \langle \xi^k, x^k - u^k \rangle \\
 &\quad + \frac{2}{\beta\rho_k} \left\langle \xi^k + w^k, u^k - \frac{1}{\beta\rho_k}\xi^k - x^k \right\rangle.
 \end{aligned}
 \tag{3.13}$$

Then the inequality (3.3) is deduced from this inequality and (3.6). □

The dual algorithm is an iterative method which generates a sequence $(u^k, x^k, \bar{w}^k, w^k)$ based on scheme (3.1). The algorithm is presented in detail as follows:

Algorithm 3.2. One has the following.

Initialization:

Given a tolerance $\varepsilon > 0$, fix an arbitrary point $\bar{x} \in \text{int } C$ and choose $\beta \geq L, R = \max\{\|x\| \mid x \in C\}$. Take $\bar{w}^{-1} := 0$ and $k := -1$.

Iterations:

For each $k = 0, 1, 2, \dots, k_\varepsilon$, execute four steps below.

Step 1. Compute a projection point u^k by taking

$$u^k := Pr_C\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right).
 \tag{3.14}$$

Step 2. Solve the strongly convex programming problem

$$\min \left\{ \langle F(u^k), y - u^k \rangle + \varphi(y) + \frac{\beta}{2} \|y - u^k\|^2 \mid y \in C \right\}
 \tag{3.15}$$

to get the unique solution x^k .

Step 3. Find $w^k \in \mathbb{R}^n$ such that

$$w^k \in -F(x^k) - \partial\varphi(x^k). \quad (3.16)$$

Set $\bar{w}^k := \bar{w}^{k-1} + w^k$.

Step 4. Compute

$$r_k := \sum_{i=0}^k \langle w^i, \bar{x} - x^i \rangle + \max \left\{ \langle \bar{w}^k, y - \bar{x} \rangle \mid y \in C_R \right\}. \quad (3.17)$$

If $r_k \leq (k+1)\varepsilon$, where $\varepsilon > 0$ is a given tolerance, then stop.

Otherwise, increase k by 1 and go back to Step 1.

Output:

Compute the final output \bar{x}^k as:

$$\bar{x}_k := \frac{1}{k+1} \sum_{i=0}^k x^i. \quad (3.18)$$

Now, we prove the convergence of Algorithm 3.2 and estimate its complexity.

Theorem 3.3. *Suppose that assumptions (A₁)–(A₃) are satisfied and F is L -Lipschitz continuous on C . Then, one has*

$$g_R(\bar{x}^k) \leq \frac{\beta R^2}{2(k+1)}, \quad (3.19)$$

where \bar{x}^k is the final output defined by the sequence $(u^k, x^k, \bar{w}^k, w^k)_{k \geq 0}$ in Algorithm 3.2. As a consequence, the sequence $\{g_R(\bar{x}^k)\}$ converges to 0 and the number of iterations to reach an ε -solution is $k_\varepsilon := \lceil \beta R^2 / 2\varepsilon \rceil$, where $\lceil x \rceil$ denotes the largest integer such that $\lceil x \rceil \leq x$.

Proof. From $\xi^k = \eta^k + F(u^k)$, where $\eta_k \in \partial\varphi(x^k)$ and $\pi_C^k \in C$, we get

$$\begin{aligned} \langle \xi^k + w^k, \pi_C^k - x^k \rangle &= \langle F(x^k) - F(u^k), x^k - \pi_C^k \rangle \\ &\leq \frac{L}{2} \left(\|x^k - u^k\|^2 + \|x^k - \pi_C^k\|^2 \right). \end{aligned} \quad (3.20)$$

Substituting (3.20) into (3.2), we obtain

$$\begin{aligned} d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^k\right) &\geq d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) + \left(1 - \frac{L}{\beta\rho_k}\right)\left(\|x^k - u^k\|^2 + \|\pi_C^k - x^k\|^2\right) \\ &\quad + \frac{1}{\beta^2\rho_k^2}\|w^k\|^2 + \frac{2}{\beta\rho_k}\left\langle w^k, \bar{x} - x^k + \frac{1}{\beta}\bar{w}^{k-1} \right\rangle. \end{aligned} \quad (3.21)$$

Using this inequality with $\rho_i = 1$ for all $i \geq 0$ and $\beta \geq L$, we obtain

$$\begin{aligned} d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^k\right) &\geq d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) + \left(1 - \frac{L}{\beta}\right)\left(\|x^k - u^k\|^2 + \|\pi_C^k - x^k\|^2\right) \\ &\quad + \frac{1}{\beta^2}\|w^k\|^2 + \frac{2}{\beta}\left\langle w^k, \bar{x} - x^k + \frac{1}{\beta}\bar{w}^{k-1} \right\rangle \\ &\geq d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) + \frac{1}{\beta^2}\|w^k\|^2 + \frac{2}{\beta}\left\langle w^k, \bar{x} - x^k + \frac{1}{\beta}\bar{w}^{k-1} \right\rangle. \end{aligned} \quad (3.22)$$

If we choose $\lambda_i = 1$ for all $i \geq 0$ in (2.21), then we have

$$\bar{w}^k = \sum_{i=0}^k w^i, \quad \bar{\lambda}_k = k + 1, \quad \bar{x}^k = \frac{1}{k+1} \sum_{i=0}^k x^i. \quad (3.23)$$

Hence, from Lemma 2.5(ii), we have

$$(k+1)g_R(\bar{x}^k) \leq \sum_{i=0}^k \langle w^i, \bar{x} - x^i \rangle + \frac{1}{2\beta} \|\bar{w}^k\|^2 - \frac{\beta}{2} d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^k\right) + \frac{\beta R^2}{2}. \quad (3.24)$$

Using inequality (3.22) and $\|\bar{w}^k\|^2 = \|w^k + \bar{w}^{k-1}\|^2$, it implies that

$$\begin{aligned} a_k &:= \sum_{i=0}^k \langle w^i, \bar{x} - x^i \rangle + \frac{1}{2\beta} \|\bar{w}^k\|^2 - \frac{\beta}{2} d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^k\right) + \frac{\beta R^2}{2} \\ &= \sum_{i=0}^{k-1} \langle w^i, \bar{x} - x^i \rangle + \langle w^k, \bar{x} - x^k \rangle + \frac{1}{2\beta} \|\bar{w}^k\|^2 - \frac{\beta}{2} d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^k\right) + \frac{\beta R^2}{2} \\ &\leq \sum_{i=0}^{k-1} \langle w^i, \bar{x} - x^i \rangle + \langle w^k, \bar{x} - x^k \rangle + \frac{1}{2\beta} \|\bar{w}^k\|^2 + \frac{\beta R^2}{2} \\ &\quad - \frac{\beta}{2} \left(d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) + \frac{1}{\beta^2} \|w^k\|^2 + \frac{2}{\beta} \left\langle w^k, \bar{x} - x^k + \frac{1}{\beta}\bar{w}^{k-1} \right\rangle \right) \\ &= \sum_{i=0}^{k-1} \langle w^i, \bar{x} - x^i \rangle + \frac{1}{2\beta} \|\bar{w}^{k-1}\|^2 - \frac{\beta}{2} d_C^2\left(\bar{x} + \frac{1}{\beta}\bar{w}^{k-1}\right) + \frac{\beta R^2}{2} \\ &= a_{k-1}. \end{aligned} \quad (3.25)$$

Note that $a_{-1} = \beta R^2/2$. It follows from the inequalities (3.24) and (3.25) that

$$(k+1)g_R(\bar{x}^k) \leq \frac{\beta R^2}{2}, \quad (3.26)$$

which implies that $g_R(\bar{x}^k) \leq \beta R^2/2(k+1)$. The termination criterion at Step 4, $r_k \leq (k+1)\epsilon$, using inequality (2.26) we obtain $g_R(\bar{x}^k) \leq \epsilon$ and the number of iterations to reach an ϵ -solution is $k_\epsilon := \lceil \beta R^2/2\epsilon \rceil$. \square

If there is no the guarantee for the Lipschitz condition, but the sequences (w^k) and (ξ^k) are uniformly bounded, we suppose that

$$M = \sup_k \|F(x^k) - F(u^k)\| = \sup_k \|w^k + \xi^k\|, \quad (3.27)$$

then the algorithm can be modified to ensure that it still converges. The variant of Algorithm 3.2 is presented as Algorithm 3.4 below.

Algorithm 3.4. One has the following.

Initialization:

Fix an arbitrary point $\bar{x} \in \text{int} C$ and set $R = \max\{\|x\| \mid x \in C\}$. Take $\bar{w}^{-1} := 0$ and $k := -1$. Choose $\beta_k = M/R$ for all $k \geq 0$.

Iterations:

For each $k = 0, 1, 2, \dots$ execute the following steps.

Step 1. Compute the projection point u_k by taking

$$u^k := \text{Pr}_C\left(\bar{x} + \frac{1}{\beta_k} \bar{w}^{k-1}\right). \quad (3.28)$$

Step 2. Solve the strong convex programming problem

$$\min \left\{ \langle F(u^k), y - u^k \rangle + \varphi(y) + \frac{\beta_k}{2} \|y - u^k\|^2 \mid y \in C \right\} \quad (3.29)$$

to get the unique solution x^k .

Step 3. Find $w^k \in \mathbb{R}^n$ such that

$$w^k \in -F(x^k) - \partial\varphi(x^k). \quad (3.30)$$

Set $\bar{w}^k := \bar{w}^{k-1} + w^k$.

Step 4. Compute

$$r_k := \sum_{i=0}^k \langle w^i, \bar{x} - x^i \rangle + \max \left\{ \langle \bar{w}^k, y - \bar{x} \rangle \mid y \in C_R \right\}. \quad (3.31)$$

If $r_k \leq (k+1)\varepsilon$, where $\varepsilon > 0$ is a given tolerance, then stop.

Otherwise, increase k by 1, update $\beta_k := (M/R)\sqrt{k+1}$ and go back to Step 1.

Output:

Compute the final output \bar{x}^k as

$$\bar{x}_k := \frac{1}{k+1} \sum_{i=0}^k x^i. \quad (3.32)$$

The next theorem shows the convergence of Algorithm 3.4.

Theorem 3.5. *Let assumptions (A_1) – (A_3) be satisfied and the sequence $(u^k, x^k, \bar{w}^k, w^k)$ be generated by Algorithm 3.4. Suppose that the sequences $(F(x^k))$ and $(F(u^k))$ are uniformly bounded by (3.27). Then, we have*

$$g_R(\bar{x}^k) \leq \frac{MR}{\sqrt{k+1}}. \quad (3.33)$$

As a consequence, the sequence $\{g_R(\bar{x}^k)\}$ converges to 0 and the number of iterations to reach an ε -solution is $k_\varepsilon := \lceil M^2 R^2 / \varepsilon^2 \rceil$.

Proof. If we choose $\lambda_k = 1$ for all $k \geq 0$ in (2.21), then we have $\bar{\lambda}_k = k+1$. Since $\bar{w}^{-1} = 0$, it follows from Step 3 of Algorithm 3.4 that

$$\bar{w}^k = \sum_{i=0}^k w^i. \quad (3.34)$$

From (3.34) and Lemma 2.5(ii), for all $\beta_k \geq 1$ we have

$$(k+1)g_R(\bar{x}^k) \leq \sum_{i=0}^k \langle w^i, \bar{x} - x^i \rangle + \frac{1}{2\beta_k} \|\bar{w}^k\|^2 - \frac{\beta_k}{2} d_C^2 \left(\bar{x} + \frac{1}{\beta_k} \bar{w}^k \right) + \frac{\beta_k R^2}{2}. \quad (3.35)$$

We define $b_k := \sum_{i=0}^k \langle w^i, \bar{x} - x^i \rangle + (1/2\beta_k) \|\bar{w}^k\|^2 - (\beta_k/2) d_C^2(\bar{x} + (1/\beta)_k \bar{w}^k)$. Then, we have

$$\begin{aligned} b_k - b_{k-1} &= \langle w^k, \bar{x} - x^k \rangle + \frac{1}{2\beta_k} \|\bar{w}^k\|^2 - \frac{\beta_k}{2} d_C^2\left(\bar{x} + \frac{1}{\beta_k} \bar{w}^k\right) - \frac{1}{2\beta_{k-1}} \|\bar{w}^{k-1}\|^2 \\ &\quad + \frac{\beta_{k-1}}{2} d_C^2\left(\bar{x} + \frac{1}{\beta_{k-1}} \bar{w}^{k-1}\right). \end{aligned} \quad (3.36)$$

We consider, for all $y \in \mathbb{R}^n$

$$\begin{aligned} q(\beta) &:= \frac{1}{2\beta} \|y\|^2 - \frac{\beta}{2} d_C^2\left(x + \frac{1}{\beta} w\right) \\ &= \frac{1}{2\beta} \|y\|^2 - \frac{\beta}{2} \min_{v \in C} \left\| v - x - \frac{1}{\beta} w \right\|^2. \end{aligned} \quad (3.37)$$

Then derivative of q is given by

$$q'(\beta) = - \left\| \text{Pr}_C\left(x + \frac{1}{\beta} y\right) - x \right\|^2 \leq 0. \quad (3.38)$$

Thus q is nonincreasing. Combining this with (3.36) and $0 < \beta_{k-1} < \beta_k$, we have

$$\begin{aligned} b_k - b_{k-1} &\leq \langle w^k, \bar{x} - x^k \rangle + \frac{1}{2\beta_k} \|\bar{w}^k\|^2 - \frac{\beta_k}{2} d_C^2\left(\bar{x} + \frac{1}{\beta_k} \bar{w}^k\right) - \frac{1}{2\beta_k} \|\bar{w}^{k-1}\|^2 \\ &\quad + \frac{\beta_k}{2} d_C^2\left(\bar{x} + \frac{1}{\beta_k} \bar{w}^{k-1}\right). \end{aligned} \quad (3.39)$$

From Lemma 3.1, $\beta = \beta_k$ and $\rho_k = 1$, we have

$$\begin{aligned} d_C^2\left(\bar{x} + \frac{1}{\beta_k} \bar{w}^k\right) - d_C^2\left(\bar{x} + \frac{1}{\beta_k} \bar{w}^{k-1}\right) &\geq \frac{2}{\beta_k} \langle w^k, \bar{x} - x^k \rangle + \frac{1}{\beta_k^2} \|\bar{w}^k\|^2 - \frac{1}{\beta_k^2} \|\bar{w}^{k-1}\|^2 \\ &\quad - \frac{1}{\beta_k^2} \|\xi^k + w^k\|^2. \end{aligned} \quad (3.40)$$

Combining (3.39) and this inequality, we have

$$b_k - b_{k-1} \leq \frac{\|\xi^k + w^k\|^2}{2\beta_k} = \frac{\|F(x^k) - F(u^k)\|^2}{2\beta_k} \leq \frac{MR}{2\sqrt{k+1}}. \quad (3.41)$$

By induction on k , it follows from (3.41) and $\beta_0 := (M_x + M_u)/R$ that

$$b_k \leq \frac{MR}{2} \sum_{i=0}^k \frac{1}{\sqrt{i+1}} \leq \frac{MR}{2} \sqrt{k+1} \equiv \frac{\beta_k R^2}{2}. \quad (3.42)$$

From (3.35) and (3.42), we obtain

$$(k+1)g_R(\bar{x}^k) \leq \beta_k R^2 = MR\sqrt{k+1}, \quad (3.43)$$

which implies that $g_R(x^k) \leq MR/\sqrt{k+1}$. The remainder of the theorem is trivially follows from (3.33). \square

4. Illustrative Example and Numerical Results

In this section, we illustrate the proposed algorithms on a class of generalized variational inequalities (GVI), where C is a polyhedral convex set given by

$$C := \{x \in \mathbb{R}^n \mid Ax \leq b\}, \quad (4.1)$$

where $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$. The cost function $F : C \rightarrow \mathbb{R}$ is defined by

$$F(x) = D(x) - Mx + q, \quad (4.2)$$

where $D : C \rightarrow \mathbb{R}^n$, $M \in \mathbb{R}^{n \times n}$ is a symmetric positive semidefinite matrix and $q \in \mathbb{R}^n$. The function φ is defined by

$$\varphi(x) := \sum_{i=1}^n (x_i^2 + |x_i - i|). \quad (4.3)$$

Then φ is subdifferentiable, but it is not differentiable on \mathbb{R}^n .

For this class of problem (GVI) we have the following results.

Lemma 4.1. *Let $D : C \rightarrow \mathbb{R}^n$. Then*

- (i) *if D is τ -strongly monotone on C , then F is monotone on C whenever $\tau = \|M\|$.*
- (ii) *if D is τ -strongly monotone on C , then F is $(\tau - \|M\|)$ -strongly monotone on C whenever $\tau > \|M\|$.*
- (iii) *if D is L -Lipschitz on C , then F is $(L + \|M\|)$ -Lipschitz on C .*

Proof. Since D is τ -strongly monotone on C , that is

$$\begin{aligned} \langle D(x) - D(y), x - y \rangle &\geq \tau \|x - y\|^2, \quad \forall x, y \in C, \\ \langle M(x - y), x - y \rangle &\leq \|M\| \|x - y\|^2, \quad \forall x, y \in C, \end{aligned} \quad (4.4)$$

we have

$$\begin{aligned} \langle F(x) - F(y), x - y \rangle &= \langle D(x) - D(y), x - y \rangle - \langle M(x - y), x - y \rangle \\ &\geq (\tau - \|M\|) \|x - y\|^2, \quad \forall x, y \in C. \end{aligned} \quad (4.5)$$

Then (i) and (ii) easily follow.

Using the Lipschitz condition, it is not difficult to obtain (iii). \square

To illustrate our algorithms, we consider the following data.

$$n = 10, \quad D(x) := \tau x, \quad q = (1, -1, 2, -3, 1, -4, 5, 6, -2, 7)^T,$$

$$C := \left\{ x \in \mathbb{R}^{10} \mid \sum_{i=1}^{10} x_i \geq -2, -1 \leq x_i \leq 1 \right\},$$

$$M = \begin{bmatrix} 1 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 2.2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 4 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 4.5 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & 1.5 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 2.5 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 3.5 \end{bmatrix}, \quad (4.6)$$

$$\bar{x} = (0, 0, 0, 0, 0, 0, 0, 0, 0, 0) \in \text{int } C, \quad \epsilon = 10^{-6}, \quad R = \sqrt{10},$$

with $\tau = \|M\| = 2.2071$, $L = \tau + \|M\| = 4.4142$, $\beta = L/2 = 2.2071$. From Lemma 4.1, we have F is monotone on C . The subproblems in Algorithm 3.2 can be solved efficiently, for example, by using MATLAB Optimization Toolbox R2008a. We obtain the approximate solution

$$x^{10} = (0.0510, 0.6234, -0.2779, 1.0000, 0.0449, 1.0000, -1.0000, 1.0000, 0.7927, -1.0000)^T. \quad (4.7)$$

Now we use Algorithm 3.4 on the same variational inequalities except that

$$F(x) := \tau x + D(x) - Mx + q, \quad (4.8)$$

where the n components of the $D(x)$ are defined by: $D_j(x) = d_j \arctan(x_j)$, with d_j randomly chosen in $(0, 1)$ and the n components of q are randomly chosen in $(-1, 3)$. The function D is given by Bnouhachem [19]. Under these assumptions, it can be proved that F is continuous and monotone on C .

Table 1: Numerical results: Algorithm 3.4 with $n = 10$.

P	x_1^k	x_2^k	x_3^k	x_4^k	x_5^k	x_6^k	x_7^k	x_8^k	x_9^k	x_{10}^k
1	-0.278	0.001	-0.006	-0.377	0.272	-0.007	-0.462	-0.227	0.395	-0.364
2	-0.054	0.133	-0.245	-0.435	-0.348	0.080	0.493	-0.223	-0.146	0.307
3	-0.417	0.320	-0.027	-0.270	0.463	-0.375	-0.381	0.255	-0.087	-0.403
4	0.197	0.161	0.434	-0.090	0.505	-0.001	0.451	-0.358	-0.320	0.278
5	0.291	0.071	-0.383	-0.290	0.453	-0.035	-0.393	-0.536	0.238	0.166
6	-0.021	0.246	0.211	-0.036	0.044	-0.241	0.466	-0.186	0.486	-0.072
7	-0.429	0.220	0.134	0.321	-0.312	0.364	-0.278	0.551	0.421	-0.118
8	-0.349	-0.448	0.365	-0.467	-0.137	0.387	0.217	-0.049	-0.443	-0.453
9	-0.115	0.562	-0.371	-0.536	-0.198	-0.248	-0.233	0.124	-0.149	0.319
10	0.071	0.134	-0.268	-0.340	0.307	0.010	0.052	-0.168	-0.206	-0.244

With $\bar{x} = (0, 0, 0, 0, 0, 0, 0, 0, 0, 0) \in \text{int } C$ and the tolerance $\epsilon = 10^{-6}$, we obtained the computational results (see, the Table 1).

Acknowledgments

The authors would like to thank the referees for their useful comments, remarks and suggestions. This work was completed while the first author was staying at Kyungnam University for the NRF Postdoctoral Fellowship for Foreign Researchers. And the second author was supported by Kyungnam University Research Fund, 2010.

References

- [1] P. N. Anh, L. D. Muu, and J.-J. Strodiot, "Generalized projection method for non-Lipschitz multivalued monotone variational inequalities," *Acta Mathematica Vietnamica*, vol. 34, no. 1, pp. 67–79, 2009.
- [2] P. N. Anh, L. D. Muu, V. H. Nguyen, and J. J. Strodiot, "Using the Banach contraction principle to implement the proximal point method for multivalued monotone variational inequalities," *Journal of Optimization Theory and Applications*, vol. 124, no. 2, pp. 285–306, 2005.
- [3] J. Y. Bello Cruz and A. N. Iusem, "Convergence of direct methods for paramontone variational inequalities," *Computational Optimization and Applications*, vol. 46, no. 2, pp. 247–263, 2010.
- [4] F. Facchinei and J. S. Pang, *Finite-Dimensional Variational Inequalities and Complementary Problems*, Springer, New York, NY, USA, 2003.
- [5] M. Fukushima, "Equivalent differentiable optimization problems and descent methods for asymmetric variational inequality problems," *Mathematical Programming*, vol. 53, no. 1, pp. 99–110, 1992.
- [6] I. V. Konnov, *Combined Relaxation Methods for Variational Inequalities*, Springer, Berlin, Germany, 2000.
- [7] J. Mashreghi and M. Nasri, "Forcing strong convergence of Korpelevich's method in Banach spaces with its applications in game theory," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 3-4, pp. 2086–2099, 2010.
- [8] M. A. Noor, "Iterative schemes for quasimonotone mixed variational inequalities," *Optimization*, vol. 50, no. 1-2, pp. 29–44, 2001.
- [9] D. L. Zhu and P. Marcotte, "Co-coercivity and its role in the convergence of iterative schemes for solving variational inequalities," *SIAM Journal on Optimization*, vol. 6, no. 3, pp. 714–726, 1996.
- [10] P. Daniele, F. Giannessi, and A. Maugeri, *Equilibrium Problems and Variational Models*, vol. 68 of *Nonconvex Optimization and Its Applications*, Kluwer Academic Publishers, Norwell, Mass, USA, 2003.
- [11] S. C. Fang and E. L. Peterson, "Generalized variational inequalities," *Journal of Optimization Theory and Applications*, vol. 38, no. 3, pp. 363–383, 1982.

- [12] C. J. Goh and X. Q. Yang, *Duality in Optimization and Variational Inequalities*, vol. 2 of *Optimization Theory and Applications*, Taylor & Francis, London, UK, 2002.
- [13] A. N. Iusem and M. Nasri, "Inexact proximal point methods for equilibrium problems in Banach spaces," *Numerical Functional Analysis and Optimization*, vol. 28, no. 11-12, pp. 1279–1308, 2007.
- [14] J. K. Kim and K. S. Kim, "New systems of generalized mixed variational inequalities with nonlinear mappings in Hilbert spaces," *Journal of Computational Analysis and Applications*, vol. 12, no. 3, pp. 601–612, 2010.
- [15] J. K. Kim and K. S. Kim, "A new system of generalized nonlinear mixed quasivariational inequalities and iterative algorithms in Hilbert spaces," *Journal of the Korean Mathematical Society*, vol. 44, no. 4, pp. 823–834, 2007.
- [16] R. A. Waltz, J. L. Morales, J. Nocedal, and D. Orban, "An interior algorithm for nonlinear optimization that combines line search and trust region steps," *Mathematical Programming*, vol. 107, no. 3, pp. 391–408, 2006.
- [17] P. N. Anh, "An interior proximal method for solving monotone generalized variational inequalities," *East-West Journal of Mathematics*, vol. 10, no. 1, pp. 81–100, 2008.
- [18] A. Auslender and M. Teboulle, "Interior projection-like methods for monotone variational inequalities," *Mathematical Programming*, vol. 104, no. 1, pp. 39–68, 2005.
- [19] A. Bnouhachem, "An LQP method for pseudomonotone variational inequalities," *Journal of Global Optimization*, vol. 36, no. 3, pp. 351–363, 2006.
- [20] A. N. Iusem and M. Nasri, "Augmented Lagrangian methods for variational inequality problems," *RAIRO Operations Research*, vol. 44, no. 1, pp. 5–25, 2010.
- [21] J. K. Kim, S. Y. Cho, and X. Qin, "Hybrid projection algorithms for generalized equilibrium problems and strictly pseudocontractive mappings," *Journal of Inequalities and Applications*, vol. 2010, Article ID 312062, 17 pages, 2010.
- [22] J. K. Kim and N. Buong, "Regularization inertial proximal point algorithm for monotone hemicontinuous mapping and inverse strongly monotone mappings in Hilbert spaces," *Journal of Inequalities and Applications*, vol. 2010, Article ID 451916, 10 pages, 2010.
- [23] Y. Nesterov, "Dual extrapolation and its applications to solving variational inequalities and related problems," *Mathematical Programming*, vol. 109, no. 2-3, pp. 319–344, 2007.
- [24] J.-P. Aubin and I. Ekeland, *Applied Nonlinear Analysis*, Pure and Applied Mathematics, John Wiley & Sons, New York, NY, USA, 1984.
- [25] P. N. Anh and L. D. Muu, "Coupling the Banach contraction mapping principle and the proximal point algorithm for solving monotone variational inequalities," *Acta Mathematica Vietnamica*, vol. 29, no. 2, pp. 119–133, 2004.
- [26] G. Cohen, "Auxiliary problem principle extended to variational inequalities," *Journal of Optimization Theory and Applications*, vol. 59, no. 2, pp. 325–333, 1988.
- [27] O. L. Mangasarian and M. V. Solodov, "A linearly convergent derivative-free descent method for strongly monotone complementarity problems," *Computational Optimization and Applications*, vol. 14, no. 1, pp. 5–16, 1999.
- [28] R. T. Rockafellar, "Monotone operators and the proximal point algorithm," *SIAM Journal on Control and Optimization*, vol. 14, no. 5, pp. 877–898, 1976.