

## Research Article

# An Iterative Algorithm of Solution for Quadratic Minimization Problem in Hilbert Spaces

Li Liu,<sup>1</sup> Guanghui Gu,<sup>1</sup> and Yongfu Su<sup>2</sup>

<sup>1</sup> Department of Mathematics, Cangzhou Normal University, Hebei, Cangzhou 061001, China

<sup>2</sup> Department of Mathematics, Tianjin Polytechnic University, Tianjin 300160, China

Correspondence should be addressed to Yongfu Su, suyongfu@tjpu.edu.cn

Received 24 October 2009; Accepted 28 December 2009

Academic Editor: Jong Kim

Copyright © 2010 Li Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The purpose of this paper is to introduce an iterative algorithm for finding a solution of quadratic minimization problem in the set of fixed points of a nonexpansive mapping and to prove a strong convergence theorem of the solution for quadratic minimization problem. The result of this article improved and extended the result of G. Marino and H. K. Xu and some others.

## 1. Introduction and Preliminaries

Iterative methods for nonexpansive mappings have recently been applied to solve convex minimization problems; see, for example, [1, 2] and the references therein. A typical problem is to minimize a quadratic function over the set of fixed points of a nonexpansive mapping on a real Hilbert space  $H$ :

$$\min_{x \in C} \frac{1}{2} \langle Ax, x \rangle - \langle x, u \rangle, \quad (1.1)$$

where  $C$  is the fixed point set of a nonexpansive mapping  $T$  defined on  $H$ , and  $u$  is a given point in  $H$ . Let  $A$  be a strongly positive operator defined on  $H$ , that is, there is a constant  $\gamma > 0$  with the property

$$\langle Ax, x \rangle \geq \gamma \|x\|^2, \quad \forall x \in H. \quad (1.2)$$

Then minimization (1.1) has a unique solution  $x^* \in C$  which satisfies the optimality condition

$$\langle Ax^* - u, x - x^* \rangle \geq 0, \quad \forall x \in C. \quad (1.3)$$

In [1, 2] it is proved that the sequence  $\{x_n\}$  generated by the following algorithm

$$x_{n+1} = (I - \alpha_n A)Tx_n + \alpha_n u, \quad n \geq 0 \quad (1.4)$$

converges in norm to the solution  $x^*$  of (1.1) provided that the sequence  $\{\alpha_n\}$  in  $(0, 1)$  satisfies conditions

$$\lim_{n \rightarrow \infty} \alpha_n = 0, \quad (C_1)$$

$$\sum_{n=1}^{\infty} \alpha_n = \infty, \quad (C_2)$$

and additionally, either the condition

$$\sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty, \quad (C_3)$$

or the condition

$$\lim_{n \rightarrow \infty} \frac{|\alpha_{n+1} - \alpha_n|}{\alpha_{n+1}} = 0. \quad (C_4)$$

The purpose of this paper is to introduce the following iterative algorithm:

$$\begin{aligned} x_{n+1} &= (I - \alpha_n A)y_n + \alpha_n u, \\ y_n &= \beta_n x_n + (1 - \beta_n)Tx_n, \end{aligned} \quad (1.5)$$

and to prove that the iterative sequence  $\{x_n\}$  defined by (1.5) converges strongly to the solution  $x^*$  of (1.1) under the conditions  $(C_1)$ ,  $(C_2)$  and  $0 < a \leq \beta_n \leq b < 1$  for some constants  $a, b$ .

**Lemma 1.1** (see [3, 4]). *Let  $\{x_n\}$  and  $\{y_n\}$  be bounded sequences in a Banach space  $X$  such that*

$$x_{n+1} = \lambda_n x_n + (1 - \lambda_n)y_n, \quad n \geq 0, \quad (1.6)$$

where  $\{\lambda_n\}$  is a sequence in  $[0, 1]$  such that

$$0 < \liminf_{n \rightarrow \infty} \lambda_n \leq \limsup_{n \rightarrow \infty} \lambda_n < 1. \quad (1.7)$$

Assume that

$$\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (1.8)$$

Then  $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$ .

**Lemma 1.2** (see [1]). Assume that  $A$  is a strongly positive linear bounded operator on a real Hilbert space  $H$  with coefficient  $\gamma > 0$  and  $0 < \alpha \leq \|A\|^{-1}$ . Then  $\|I - \alpha A\| \leq (1 - \alpha\gamma)$ .

**Lemma 1.3** (see [5]). Let  $H$  be a Hilbert space,  $K$  a closed convex subset of  $H$ , and  $T : K \rightarrow K$  a nonexpansive mapping with nonempty fixed point set  $F(T)$ . If  $\{x_n\}$  is a sequence in  $K$  weakly converging to  $x$  and if  $x_n - Tx_n$  converges strongly to 0, then  $x = Tx$ .

**Lemma 1.4** (see [6]). Assume that  $\{a_n\}$  is a sequence of nonnegative real numbers such that

$$a_{n+1} \leq (1 - \gamma_n)a_n + \gamma_n\delta_n, \quad (1.9)$$

where  $\{\gamma_n\}$  is a sequence in  $(0, 1)$  and  $\{\delta_n\}$  is a real sequence such that

- (i)  $\sum_{n=1}^{\infty} \gamma_n = \infty$ ,
- (ii)  $\limsup_{n \rightarrow \infty} \delta_n \leq 0$  or  $\sum_{n=1}^{\infty} \gamma_n |\delta_n| < \infty$ .

Then  $\lim_{n \rightarrow \infty} a_n = 0$ .

## 2. Main Results

**Theorem 2.1.** Suppose that  $A$  is strongly positive operator with coefficient  $\gamma > 0$  as given in (1.2). Suppose that the sequences  $\{\alpha_n\}, \{\beta_n\}$  satisfy the conditions  $(C_1), (C_2)$  and  $0 < a \leq \beta_n \leq b < 1$  for some constants  $a, b$ . Then the sequence  $\{x_n\}$  generated by algorithm (1.5) converges strongly to the unique solution  $x^*$  of the minimization problem (1.1).

*Proof.* First we show that  $\{x_n\}$  is bounded. As a matter of fact, take  $p \in F(T)$  and use Lemma 1.2 to obtain

$$\begin{aligned} \|x_{n+1} - p\| &= \|(I - \alpha_n A)(\beta_n x_n + (1 - \beta_n)Tx_n - p) + \alpha_n(u - Ap)\| \\ &\leq (1 - \gamma\alpha_n)\|x_n - p\| + \alpha_n\|u - Ap\|. \end{aligned} \quad (2.1)$$

By induction we can get

$$\|x_n - p\| \leq \max\left\{\|x_0 - p\|, \frac{1}{\gamma}\|u - Ap\|\right\}, \quad n \geq 0. \quad (2.2)$$

Hence,  $\{x_n\}$  is bounded and so is  $\{y_n\}$ . Next rewrite  $x_{n+1}$  in the form

$$x_{n+1} = (1 - \lambda_n)x_n + \lambda_n z_n, \quad (2.3)$$

where

$$\lambda_n = 1 - (1 - \alpha_n)\beta_n, \quad (2.4)$$

$$z_n = \frac{\alpha_n\beta_n}{\lambda_n}(I - A)x_n + \frac{1 - \beta_n}{\lambda_n}(I - \alpha_n A)Tx_n + \frac{\alpha_n}{\lambda_n}u. \quad (2.5)$$

Since  $\alpha_n \rightarrow 0$  and  $0 < a \leq \beta_n \leq b < 1$ , then

$$0 < \liminf_{n \rightarrow \infty} \lambda_n \leq \limsup_{n \rightarrow \infty} \lambda_n < 1. \quad (2.6)$$

Next some manipulations give us that

$$\begin{aligned} z_{n+1} - z_n &= \frac{\beta_{n+1}\alpha_{n+1}}{\lambda_{n+1}}(I - A)x_{n+1} - \frac{\beta_n\alpha_n}{\lambda_n}(I - A)x_n + \left(\frac{\alpha_{n+1}}{\lambda_{n+1}} - \frac{\alpha_n}{\lambda_n}\right)u \\ &\quad + \frac{1 - \beta_{n+1}}{\lambda_{n+1}}(Tx_{n+1} - Tx_n) - \frac{(1 - \beta_{n+1})\alpha_{n+1}}{\lambda_{n+1}}A(Tx_{n+1} - Tx_n) \\ &\quad + \left(\frac{1 - \beta_{n+1}}{\lambda_{n+1}} - \frac{1 - \beta_n}{\lambda_n}\right)Tx_n - \left(\frac{\alpha_{n+1}}{\lambda_{n+1}} - \frac{\alpha_n}{\lambda_n}\right)(1 - \beta_n)ATx_n \\ &\quad - \frac{\alpha_{n+1}}{\lambda_{n+1}}(\beta_n - \beta_{n+1})ATx_n. \end{aligned} \quad (2.7)$$

Therefore,

$$\begin{aligned} \|z_{n+1} - z_n\| - \|x_{n+1} - x_n\| &\leq \frac{\beta_{n+1}\alpha_{n+1}}{\lambda_{n+1}}\|(I - A)x_{n+1}\| + \frac{\beta_n\alpha_n}{\lambda_n}\|(I - A)x_n\| + \left|\frac{\alpha_{n+1}}{\lambda_{n+1}} - \frac{\alpha_n}{\lambda_n}\right\| \|u\| \\ &\quad + \left(\frac{1 - \beta_{n+1}}{\lambda_{n+1}} - 1\right)\|x_{n+1} - x_n\| + \frac{(1 - \beta_{n+1})\alpha_{n+1}}{\lambda_{n+1}}\|A\|\|x_{n+1} - x_n\| \\ &\quad + \left|\frac{1 - \beta_{n+1}}{\lambda_{n+1}} - \frac{1 - \beta_n}{\lambda_n}\right\| \|Tx_n\| + \left|\frac{\alpha_{n+1}}{\lambda_{n+1}} - \frac{\alpha_n}{\lambda_n}\right\| \|(1 - \beta_n)ATx_n\| \\ &\quad + \frac{\alpha_{n+1}}{\lambda_{n+1}}|\beta_n - \beta_{n+1}|\|ATx_n\|. \end{aligned} \quad (2.8)$$

Since  $\lambda_n = 1 - (1 - \alpha_n)\beta_n$  and  $\alpha_n \rightarrow 0$ , then

$$\lim_{n \rightarrow \infty} \frac{1 - \beta_n}{\lambda_n} = \lim_{n \rightarrow \infty} \left(1 - \frac{\alpha_n\beta_n}{\lambda_n}\right) = 1. \quad (2.9)$$

Then last inequality implies that

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0, \quad (2.10)$$

and so an application of Lemma 1.1 asserts that

$$\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0. \quad (2.11)$$

By (2.5) we have that

$$z_n - Tx_n = \frac{\alpha_n \beta_n}{\lambda_n} (I - A)x_n + \left( \frac{1 - \beta_n}{\lambda_n} - 1 \right) Tx_n - \frac{(1 - \beta_n) \alpha_n}{\lambda_n} ATx_n + \frac{\alpha_n}{\lambda_n} u. \quad (2.12)$$

Again since  $\alpha_n \rightarrow 0$ ,  $\{x_n\}$  is bounded, and  $\lambda_n = 1 - (1 - \alpha_n)\beta_n$ , then we deduce from (2.12) that

$$\lim_{n \rightarrow \infty} \|z_n - Tx_n\| = 0. \quad (2.13)$$

This together with (2.11) yields

$$\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0. \quad (2.14)$$

By using Lemma 1.3, we obtain  $\omega_w(x_n) \subset F(T)$ , where  $\omega_w(x_n) = \{z : \exists x_{n_k} \rightarrow z\}$  is the set of weak  $\omega$ -limit points of sequence  $\{x_n\}$ .

Let  $x^*$  be the unique solution to the minimization (1.1). Then by the definition of algorithm (1.5), we can write

$$x_{n+1} - x^* = (I - \alpha_n A)(\beta_n x_n + (1 - \beta_n)Tx_n - x^*) + \alpha_n(u - Ax^*). \quad (2.15)$$

Since  $H$  is a Hilbert space, then we have that

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &\leq \|(I - \alpha_n A)(\beta_n x_n + (1 - \beta_n)Tx_n - x^*)\|^2 + 2\alpha_n \langle u - Ax^*, x_{n+1} - x^* \rangle \\ &\leq (1 - \gamma \alpha_n) \|x_n - x^*\|^2 + 2\alpha_n \langle u - Ax^*, x_{n+1} - x^* \rangle. \end{aligned} \quad (2.16)$$

However, we can take a subsequence  $\{x_{n_k}\}$  of  $\{x_n\}$  such that

$$\limsup_{n \rightarrow \infty} \langle u - Ax^*, x_n - x^* \rangle = \lim_{k \rightarrow \infty} \langle u - Ax^*, x_{n_k} - x^* \rangle, \quad (2.17)$$

and also  $\{x_{n_k}\}$  converges weakly to a fixed point  $p \in F(T)$ . It follows from optimality condition (1.3) that

$$\limsup_{n \rightarrow \infty} \langle u - Ax^*, x_n - x^* \rangle = \langle u - Ax^*, p - x^* \rangle \leq 0. \quad (2.18)$$

Therefore, by using Lemma 1.4 and noticing (2.18), we conclude that  $x_n \rightarrow x^*$ . This completes the proof.  $\square$

## References

- [1] G. Marino and H.-K. Xu, "A general iterative method for nonexpansive mappings in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 318, no. 1, pp. 43–52, 2006.
- [2] H.-K. Xu, "An iterative approach to quadratic optimization," *Journal of Optimization Theory and Applications*, vol. 116, no. 3, pp. 659–678, 2003.
- [3] T. Suzuki, "Strong convergence theorems for infinite families of nonexpansive mappings in general Banach spaces," *Fixed Point Theory and Applications*, vol. 2005, no. 1, pp. 103–123, 2005.
- [4] T. Suzuki, "Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter nonexpansive semigroups without Bochner integrals," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.
- [5] K. Goebel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, vol. 28 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge, UK, 1990.
- [6] H.-K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society*, vol. 66, no. 1, pp. 240–256, 2002.