

RESEARCH

Open Access

# On the translations of quasimonotone maps and monotonicity

AP Farajzadeh<sup>1</sup>, A Karamian<sup>1</sup> and S Plubtieng<sup>2\*</sup>

\*Correspondence:  
somyotp@nu.ac.th

<sup>2</sup>Department of Mathematics,  
Faculty of Science, Naresuan  
University, Phitsanulok, 65000,  
Thailand

Full list of author information is  
available at the end of the article

## Abstract

We show that given a convex subset  $K$  of a topological vector space  $X$  and a multivalued map  $T : K \rightrightarrows X^*$ , if there exists a nonempty subset  $S$  of  $X^*$  with the surjective property on  $K$  and  $T + w$  is quasimonotone for each  $w \in S$ , then  $T$  is monotone. Our result is a new version of the result obtained by N. Hadjisavvas (Appl. Math. Lett. 19:913-915, 2006).

**Keywords:** monotone map; pseudomonotone map; quasimonotone map; surjective property

## 1 Introduction and some definitions

Throughout the paper,  $X$  and  $X^*$  denote a real topological vector space and the dual space of  $X$ , respectively. Suppose  $K \subseteq X$  is a nonempty subset of  $X$  and  $T : K \rightrightarrows X^*$  is a multivalued map from  $K$  to  $X^*$ . Recall that  $T$  is said to be monotone if for all  $x^* \in T(x)$ ,  $y^* \in T(y)$  one has

$$\langle x^* - y^*, x - y \rangle \geq 0.$$

$T$  is said to be pseudomonotone and quasimonotone, in the sense of Karamardian (see [1, 2]), respectively, if for any  $x^* \in T(x)$ ,  $y^* \in T(y)$  the following implications hold:

$$\langle y^*, x - y \rangle \geq 0 \quad \Rightarrow \quad \langle x^*, x - y \rangle \geq 0$$

and

$$\langle y^*, x - y \rangle > 0 \quad \Rightarrow \quad \langle x^*, x - y \rangle \geq 0.$$

It is clear that a monotone map is pseudomonotone, while a pseudomonotone map is quasimonotone. The converse is not true. If  $T$  is pseudomonotone (quasimonotone) and  $w \in X^* \setminus \{0\}$ , then  $T + w$  is not pseudomonotone (quasimonotone) in general. In the case of a single-valued linear map  $T$  defined on the whole space  $\mathbb{R}^n$ , it is known that if  $T + w$  is quasimonotone, then  $T$  is monotone [2]. Many authors (see, e.g., [4, 5]) extended this result for a nonlinear Gateaux differentiable map defined on a convex subset  $K$  (of a Hilbert space) with a nonempty interior.

Recently, Hadjisavvas [3] extended the above result to the multivalued maps defined on a convex subset of a real topological vector space with no assumption of differentiability

or even continuity on the map  $T$  whose domain need not have a nonempty interior. In this paper, we first introduce the surjective property of a subset of  $X^*$  on a segment of  $K$ . By using this concept, we can extend the corresponding result obtained in [3]. Before stating the main result, we recall some definitions.

**Definition 1** Let  $x, y$  be two elements of  $K$ . We say that  $S \subseteq X^*$  has the surjective property on  $x$  and  $y$  whenever the following equality holds:

$$\langle S, x - y \rangle = \{ \langle x^*, x - y \rangle : x^* \in S \} = \mathbb{R}.$$

Remark that we can consider  $x - y$  as a linear functional (denoted by  $\widehat{x - y}$ ) on  $X^*$  which is defined by

$$\langle \widehat{x - y}, f \rangle = \langle f, x - y \rangle.$$

Hence if  $S$  has the surjective property on  $x, y$ , then the image of  $S$  under the linear functional  $\widehat{x - y}$  is all of the real numbers, and that is why we used the phrase surjective property.

**Definition 2** Let  $K \subseteq X$  be a nonempty set and  $S \subseteq X^*$ . We say that  $S$  has the surjective property on  $K$  if for every  $x \in K$  there exists  $y \in K$  such that  $S$  has the surjective property on  $x$  and  $y$ .

**Definition 3** [3] Let  $K$  be a convex subset of  $X$ . An element  $\nu$  of  $X^*$  is called perpendicular to  $K$  if  $\nu$  is constant on  $K$ , i.e.,

$$\langle \nu, x \rangle = \langle \nu, y \rangle, \quad \forall x, y \in K.$$

Also the straight line  $S = \{u + t\nu : t \in \mathbb{R}\}$ , where  $u, \nu \in X^*$  with  $\nu \neq 0$ , is said to be perpendicular to  $K$  if  $\nu$  is perpendicular to  $K$ .

**Remark 1** If  $K \subseteq X$  is a nonempty convex set and  $u, \nu \in X^*$  with  $\nu$  is not perpendicular to  $K$ , then the straight line  $S = \{u + t\nu : t \in \mathbb{R}\}$  has the surjective property on  $K$ . Indeed, let  $x \in K$  be an arbitrary member of  $K$ . Because  $\nu$  is not perpendicular to  $K$ , there exists  $y \in K$  such that  $c = \langle \nu, x - y \rangle \neq 0$ . For each  $a \in \mathbb{R}$ , we put  $t = \frac{a - \langle u, x - y \rangle}{c}$  and so  $a = \langle u + t\nu, x - y \rangle$ . Hence  $\langle S, x - y \rangle = \mathbb{R}$ . This means that  $S$  has the surjective property. Therefore,  $\nu$  being not perpendicular to  $K$  implies the surjective property while the simple example  $X = \mathbb{R}^2$ ,  $S = \{(t, t) = (0, 0) + (1, 1)t : t \in \mathbb{R}\}$  and  $K = \{(x, -x) : x \in \mathbb{R}\}$  shows that the converse does not hold in general. In this example, one can see that  $S$  has the surjective property and  $\nu = (1, 1)$  is perpendicular to  $K$  (note  $\langle \nu = (1, 1), (x, -x) \rangle = \langle \nu = (1, 1), (y, -y) \rangle = 0$ ). The notion  $\nu$  is not perpendicular to  $K$ , which plays a crucial rule in proving the main results in [3]; while in this note, the surjective property has an essential rule in the main result. Hence one can consider this paper as an improvement of [3] (slightly, of course).

We need the following lemma in the sequel.

**Lemma 1** *Let  $X$  be a real topological vector space,  $K$  a nonempty convex subset of  $X$  and  $T : K \rightrightarrows X^*$  a multivalued map. Suppose  $x, y \in K$ ,  $S \subseteq X^*$  has the surjective property on  $x, y$  and  $T + w$  is quasimonotone on the line segment  $[x, y] = \{tx + (1 - t)y : t \in [0, 1]\}$  for all  $w \in S$ . Then  $T$  is monotone on  $[x, y]$ .*

*Proof* We can define an order on  $[x, y]$  as follows:

$$a \leq b \iff t_1 \leq t_2, \quad \text{where } a = x + t_1(y - x), b = x + t_2(y - x).$$

On the contrary, assume  $T$  is not monotone on  $[x, y]$ . So there exist  $a, b \in [x, y]$  and  $a^* \in T(a), b^* \in T(b)$  with  $a < b$  and  $\langle a^* - b^*, a - b \rangle < 0$ . Hence we have

$$\langle a^*, y - x \rangle > \langle b^*, y - x \rangle.$$

Since  $S$  is surjective on  $x, y$ , there exists  $w \in S$  such that

$$\langle a^*, y - x \rangle > \langle w, x - y \rangle > \langle b^*, y - x \rangle.$$

Therefore,

$$\langle a^* + w, y - x \rangle > 0, \quad \langle b^* + w, y - x \rangle < 0,$$

which is a contradiction. This completes the proof. □

Now we are ready to present the main result.

**Theorem 1** *Let  $X$  be a real topological vector space,  $K$  a nonempty convex subset of  $X$  and  $T : X \rightrightarrows X^*$  a multivalued map. Assume  $S \subseteq X^*$  is connected and has the surjective property on  $K$ . If  $T + w$  is quasimonotone for all  $w \in S$ , then  $T$  is monotone on  $K$ .*

*Proof* Let  $x, y \in K$ ,  $x^* \in T(x)$  and  $y^* \in T(y)$  be arbitrary elements. If  $S$  is surjective on  $x, y$  then, by Lemma 1,  $T$  is monotone on  $[x, y]$  and the proof is complete. Assume  $S$  does not have the surjective property on  $x, y$ . So  $S(x - y) \neq \mathbb{R}$ . Since  $S$  has the surjective property on  $K$ , then there exists  $z \in K$  such that  $S$  is surjective on  $x, z$ ; and since  $S$  is connected, then  $\langle S, y - z \rangle$  is a connected subset of the real numbers unbounded from above and below, and so it is equal to the real numbers. This means that  $S$  has the surjective property on  $y, z$  and also on  $\frac{x+y}{2}, z$ . Therefore, it follows from Lemma 1 that  $T$  is monotone on  $[y, z]$  and  $[\frac{x+y}{2}, z]$ . Similarly,  $T$  is monotone on the segments  $[x, z_s]$  and  $[y, z_s]$ , for all  $s \in ]0, 1[$ , where  $z_s = sz + (1 - s)\frac{x+y}{2}$ . Therefore, for any  $z_s^* \in T(z_s)$  and  $z^* \in T(z)$ , we have

$$\langle z_s^*, z_s - x \rangle \geq \langle x^*, z_s - x \rangle, \tag{1}$$

$$\langle z_s^*, z_s - y \rangle \geq \langle y^*, z_s - y \rangle, \tag{2}$$

$$\langle z^*, z - z_s \rangle \geq \langle z_s^*, z - z_s \rangle. \tag{3}$$

From (1) and (2), we deduce that

$$2s \left\langle z_s^*, z - \frac{x+y}{2} \right\rangle \geq \langle x^*, z_s - x \rangle + \langle y^*, z_s - y \rangle. \tag{4}$$

Now from  $z - z_s = (1 - s)(z - \frac{x+y}{2})$  and (3), we obtain

$$\left\langle z^*, z - \frac{x+y}{2} \right\rangle \geq \left\langle z_s^*, z - \frac{x+y}{2} \right\rangle. \quad (5)$$

Combining (4) and (5), we have

$$2s \left\langle z^*, z - \frac{x+y}{2} \right\rangle \geq \langle x^*, z_s - x \rangle + \langle y^*, z_s - y \rangle.$$

So if in the previous inequality we tend  $s \rightarrow 0$ , then  $z_s \rightarrow \frac{x+y}{2}$ , and hence we deduce

$$0 \geq \langle x^*, y - x \rangle + \langle y^*, y - x \rangle.$$

This means  $T$  is monotone and the proof is now complete.  $\square$

Remark 1 shows that Theorem 1 is a new version of Theorem 1 in [3], although our proof is, in fact, completely similar to it.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

All authors contributed equally and all authors read and approved the final manuscript.

#### Author details

<sup>1</sup>Department of Mathematics, Razi University, Kermanshah, 67149, Iran. <sup>2</sup>Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok, 65000, Thailand.

Received: 4 April 2012 Accepted: 15 August 2012 Published: 31 August 2012

#### References

1. Karamardian, S, Schaible, S: Seven kinds of monotone maps. *J. Optim. Theory Appl.* **66**, 37-46 (1990)
2. Karamardian, S, Schaible, S, Crouzeix, JP: Characterizations of generalized monotone maps. *J. Optim. Theory Appl.* **76**, 399-413 (1993)
3. Hadjisavvas, N: Translations of quasimonotone maps and monotonicity. *Appl. Math. Lett.* **19**, 913-915 (2006)
4. He, Y: A relationship between pseudomonotone and monotone mappings. *Appl. Math. Lett.* **17**, 459-461 (2004)
5. Isac, G, Motreanu, D: Pseudomonotonicity and quasimonotonicity by translations versus monotonicity in Hilbert spaces. *Aust. J. Math. Anal. Appl.* **1**(1), 1-8 (2004)

doi:10.1186/1029-242X-2012-192

Cite this article as: Farajzadeh et al.: On the translations of quasimonotone maps and monotonicity. *Journal of Inequalities and Applications* 2012 2012:192.

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)