

RESEARCH

Open Access

Minimum generalized degree distance of n -vertex tricyclic graphs

Asma Hamzeh, Ali Iranmanesh* and Samaneh Hossein-Zadeh

*Correspondence:
Iranmanesh@modares.ac.ir
Department of Mathematics,
Faculty of Mathematical Sciences,
Tarbiat Modares University, P.O. Box
14115-137, Tehran, Iran

Abstract

In (Hamzeh *et al.* in Stud. Univ. Babeş-Bolyai, Chem., 4:73-85, 2012), we introduced a generalization of a degree distance of graphs as a new topological index. In this paper, we characterize the n -vertex tricyclic graphs which have the minimum generalized degree distance.

MSC: 05C12; 05C35; 05C05

Keywords: generalized degree distance; tricyclic graphs; Harary index; join; extremal graphs

1 Introduction

Topological indices and graph invariants based on the distances between the vertices of a graph are widely used in theoretical chemistry to establish relations between the structure and the properties of molecules. They provide correlations with physical, chemical and thermodynamic parameters of chemical compounds [1–3]. The Wiener index is a well-known topological index, which equals to the sum of distances between all pairs of vertices of a molecular graph [4]. It is used to describe molecular branching and cyclicity and establish correlations with various parameters of chemical compounds. In this paper, we only consider simple and connected graphs. Let G be a connected graph with the vertex and edge sets $V(G)$ and $E(G)$, respectively, and the number of vertices and edges of G are denoted, respectively, by n and m . As usual, the distance between the vertices u and v of G is denoted by $d_G(u, v)$ ($d(u, v)$ for short). It is defined as the length of any shortest path connecting u and v in G . We let $d_G(v)$ be the degree of a vertex v in G . The eccentricity of v , denoted by $\varepsilon(v)$, is the maximum distance from vertex v to any other vertex. The diameter of a graph G is denoted by $\text{diam}(G)$ and is the maximum eccentricity over all vertices in a graph G . The radius $r = r(G)$ is defined as the minimum of $\varepsilon(v)$ over all vertices $v \in V(G)$. A connected graph G with n vertices and m edges is called tricyclic if $m = n + 2$. The join $G = G_1 + G_2$ of two graphs G_1 and G_2 with disjoint vertex sets V_1 and V_2 and edge sets E_1 and E_2 is the graph union $G_1 \cup G_2$ together with all the edges joining V_1 and V_2 .

Additively weighted Harary index is defined as follows in [5]

$$H_A(G) = \sum_{\{u,v\} \subseteq V(G)} d^{-1}(u,v)(d_G(u) + d_G(v)).$$

There are two papers [6, 7], which introduced a new graph invariant with the name degree distance. It is defined as follows:

$$D'(G) = \sum_{\{u,v\} \subseteq V(G)} d(u,v)(d_G(u) + d_G(v)).$$

Generalized degree distance is denoted by $H_\lambda(G)$ and defined as follows in [8].

For every vertex x , $H_\lambda(x)$ is defined by $H_\lambda(x) = D^\lambda(x)d_G(x)$, where $D^\lambda(x) = \sum_{y \in V(G)} d^\lambda(x, y)$, and to avoid confusion, we denote $H_\lambda(x)$ in graph G with $H_\lambda(x, G)$. So we have

$$H_\lambda(G) = \sum_{x \in V(G)} H_\lambda(x) = \sum_{x \in V(G)} D^\lambda(x)d_G(x) = \sum_{\{u,v\} \subseteq V(G)} d^\lambda(u,v)(d_G(u) + d_G(v)),$$

where λ is a real number. If $\lambda = 0$, then $H_\lambda(G) = 4m$. When $\lambda = 1$, this new topological index $H_\lambda(G)$ equals the degree distance index (i.e., the Dobrynin or the Schultz index). The properties of the degree distance index were studied in [9–12]. Also, if $\lambda = -1$, then $H_\lambda(G) = H_A(G)$ (see above). The relation of our new index with other intensely studied indices motivated our present (and future) study.

Throughout this paper, K_n and $K_{1,n-1}$ denote the complete and star graphs on n vertices, respectively. Our other notations are standard and taken mainly from [1, 2, 13].

Extremal graph theory is a branch of graph theory that studies extremal (maximal or minimal) graphs, which satisfy a certain property. Extremality can be taken with respect to different graph invariants, such as order, size or girth. The problem of determining extremal values and corresponding extremal graphs of some graph invariants is the topic of several papers, for example, see [9–12, 14–16].

2 Main results

In this paper, we characterize all of n -vertex tricyclic graphs which have the minimum generalized degree distance.

It is well known that the natural numbers $d_1 \geq d_2 \geq \dots \geq d_n \geq 1$ are the degrees of the vertices of a tree if and only if $\sum_{i=1}^n d_i = 2n - 2$ [17, 18]. The next lemma characterizes connected tricyclic graphs by their degree sequence.

Lemma 2.1 [19] *Let $n \geq 4$ and G be an n -vertex tricyclic graph. The integers $d_1 \geq d_2 \geq \dots \geq d_n \geq 1$ are the degrees of the vertices of a graph G if and only if*

- (i) $\sum_{i=1}^n d_i = 2n + 4$,
- (ii) *at least four of them are greater than or equal to 2,*
- (iii) $(d_1, d_2, \dots, d_n) \notin \{(n-1, 4, 3, 2, 1, \dots, 1), (n-1, 5, 2, 2, 1, \dots, 1)\}$ *for sufficiently large n .*

Let x_i be the number of vertices of degree i of G for $1 \leq i \leq n-1$. If $d_G(v) = k$, then

$$\begin{aligned} D^\lambda(v) &= \sum_{u \in V(G)} d^\lambda(u, v) = \sum_{u \in V(G), d(u,v)=1} d^\lambda(u, v) + \sum_{u \in V(G), d(u,v) \geq 2} d^\lambda(u, v) \\ &\geq k + 2^\lambda(n - k - 1) \\ &= 2^\lambda n - k(2^\lambda - 1) - 2^\lambda, \end{aligned}$$

so

$$\begin{aligned} H_\lambda(G) &= \sum_{v \in V(G)} d_G(v) D^\lambda(v) \\ &\geq \sum_{k=1}^{n-1} k x_k (2^\lambda n - k(2^\lambda - 1) - 2^\lambda). \end{aligned}$$

We define

$$F_\lambda(x_1, x_2, \dots, x_{n-1}) = \sum_{k=1}^{n-1} k x_k (2^\lambda n - k(2^\lambda - 1) - 2^\lambda).$$

We obtain the minimum of $F_\lambda(x_1, x_2, \dots, x_{n-1})$ over all integers numbers $x_1, x_2, \dots, x_{n-1} \geq 0$, which satisfy one of the conditions of Lemma 2.1.

We rewrite Lemma 2.1 in terms of the notations above as follows.

Lemma 2.2 [19] *Let $n \geq 4$ and G be an n -vertex tricyclic graph. The integers $x_1, x_2, \dots, x_{n-1} \geq 0$ are the multiplicities of the degrees of a graph G if and only if*

- (i) $\sum_{i=1}^{n-1} x_i = n$,
- (ii) $\sum_{i=1}^{n-1} i x_i = 2n + 4$,
- (iii) $x_1 \leq n - 4$,
- (iv) $(x_1, x_2, \dots, x_{n-1}) \notin \{(n-4, 1, 1, 1, 0, \dots, 0, 1), (n-4, 2, 0, 0, 1, 0, \dots, 0, 1)\}$ for sufficiently large n .

We denote the set of all vectors (x_1, \dots, x_{n-1}) , which satisfy the conditions above, by Δ . Let G be a connected graph with multiplicities of the degrees (x_1, \dots, x_{n-1}) , and let $m \geq 2$, $p > 0$, $m + p \leq n - 2$, $x_m \geq 1$ and $x_{m+p} \geq 1$. Now we consider the transformation of t_1 , which is defined as follows [11]:

$$\begin{aligned} t_1(x_1, \dots, x_{n-1}) &= (x'_1, \dots, x'_{n-1}) \\ &= (x_1, \dots, x_{m-1} + 1, x_m - 1, \dots, x_{m+p} - 1, x_{m+p+1} + 1, \dots, x_{n-1}). \end{aligned}$$

We have $x'_i = x_i$ for $i \notin \{m-1, m, m+p, m+p+1\}$ and $x'_{m-1} = x_{m-1} + 1$, $x'_m = x_m - 1$, $x'_{m+p} = x_{m+p} - 1$, $x'_{m+p+1} = x_{m+p+1} + 1$.

Let $2 \leq m \leq n - 2$, $x_m \geq 2$. Now consider the transformation t_2 defined as follows [11]:

$$\begin{aligned} t_2(x_1, \dots, x_{n-1}) &= (x'_1, \dots, x'_{n-1}) \\ &= (x_1, \dots, x_{m-1} + 1, x_m - 2, x_{m+1} + 1, \dots, x_{n-1}). \end{aligned}$$

That is $x'_i = x_i$ for $i \notin \{m-1, m, m+1\}$ and $x'_{m-1} = x_{m-1} + 1$, $x'_m = x_m - 2$, $x'_{m+1} = x_{m+1} + 1$.

Lemma 2.3 *Suppose that λ is a positive integer number, and consider the set of vectors (x_1, \dots, x_{n-1}) .*

- (1) *If $(x_1, \dots, x_{n-1}) \in \Delta$, then $t_1(x_1, \dots, x_{n-1}) \in \Delta$, unless $m = 2$ and $x_1 = n - 4$,*
- (2) *$F_\lambda(t_1(x_1, x_2, \dots, x_{n-1})) < F_\lambda(x_1, x_2, \dots, x_{n-1})$.*

Proof (1) We can easily see that $\sum_{i=1}^{n-1} x_i = \sum_{i=1}^{n-1} x'_i$, $\sum_{i=1}^{n-1} ix_i = \sum_{i=1}^{n-1} ix'_i$. If $m = 2$, $x_1 = n - 4$, then $x'_1 > n - 4$, and in this case, $t_1(x_1, \dots, x_{n-1}) \notin \Delta$. Now if $x'_1 > n - 4$, according to $x_1 \leq n - 4$, we have $m = 2$ and $x_1 = n - 4$. Therefore, we conclude that if $(x_1, \dots, x_{n-1}) \in \Delta$, then $x'_1 > n - 4$ if and only if $m = 2$ and $x_1 = n - 4$.

(2) With a simple calculation, we have

$$F_\lambda(x_1, x_2, \dots, x_{n-1}) - F_\lambda(t_1(x_1, x_2, \dots, x_{n-1})) = (2^\lambda - 1)(2p + 2) > 0.$$

The proof is now completed. \square

Lemma 2.4 Suppose that λ is a positive integer number, and consider the set of vectors (x_1, \dots, x_{n-1}) .

- (1) If $(x_1, \dots, x_{n-1}) \in \Delta$, then $t_2(x_1, \dots, x_{n-1}) \in \Delta$, unless $m = 2$ and $x_1 = n - 4$,
- (2) $F_\lambda(t_2(x_1, x_2, \dots, x_{n-1})) < F_\lambda(x_1, x_2, \dots, x_{n-1})$.

Proof The proof is similar to the proof of the previous lemma, by taking $p = 0$. \square


Theorem 2.5 Let $n \geq 4$ and λ be a positive integer. If G belongs to the class of connected tricyclic graphs on n vertices, then we have

$$\min H_\lambda(G) = (n^2 - n + 24) + 2^\lambda(n^2 + 3n - 28),$$

and all the extremal graphs are isomorphic to H or H' , where H is obtained by identifying the center of a star $K_{1,n-4}$ with an arbitrary vertex of a complete graph K_4 , and H' is obtained by identifying the center of a star $K_{1,n-5}$ with an arbitrary four degree vertex of a graph A_2 , which is shown in Figure 1, respectively.

Proof To find minimum $H_\lambda(G)$, over the classes of connected tricyclic graphs with n vertices, it is enough to find $\min_{(x_1, x_2, \dots, x_{n-1}) \in \Delta} F_\lambda(x_1, x_2, \dots, x_{n-1})$. At first, we consider the case $n = 4$. In this case, only tricyclic graph with 4 vertices is K_4 , that is, $H_\lambda(K_4) = 36$, and the theorem is proved in this case.

Now let $n \geq 5$. Let us consider $n = 5$. The all connected tricyclic graphs with 5 vertices, is A_1, A_2, A_3 and A_4 . Meanwhile, $H_\lambda(A_1) = 44 + 3 \cdot 2^{\lambda+2} = \theta(5)$, $H_\lambda(A_2) = 44 + 3 \cdot 2^{\lambda+2} = \theta(5)$, where $\theta(n) = (n^2 - n + 24) + 2^\lambda(n^2 + 3n - 28)$, $H_\lambda(A_3) = 40 + 2^{\lambda+4}$, and $H_\lambda(A_4) = 42 + 2^\lambda \cdot 14$, respectively, where A_i ($i = 1, \dots, 4$) are receipted in Figure 1. So A_1 and A_2 are extremal graphs, and the theorem is proved in this case.



Finally, let $n \geq 6$. If $x_{n-1} \geq 2$, consider two different vertices $x, y \in V(G)$ such that $d_G(x) = d_G(y) = n - 1$. Since $n \geq 6$, we can choose four different vertices $p, q, r, s \in V(G) - \{x, y\}$. Actually, the vertices p, q, r, s are all adjacent to x and y , hence G has at least four cycles $xypx, xyqx, xyrx$, and $xysx$, which contradicts the hypothesis. Therefore, $x_{n-1} \leq 1$.

Let us analyze the possible values for x_5, \dots, x_{n-2} . If there exist $5 \leq i < j \leq n - 2$ such that $x_i \geq 1$ and $x_j \geq 1$, then, by applying t_1 for the positions i and j , we obtain a new vector $(x'_1, \dots, x'_{n-1}) \in \Delta$, for which $F_\lambda(x'_1, \dots, x'_{n-1}) < F_\lambda(x_1, \dots, x_{n-1})$. Similarly, if there exists $5 \leq i \leq n - 2$ such that $x_i \geq 2$, then by t_2 , we obtain a new degree sequence in Δ , for which $F_\lambda(x'_1, \dots, x'_{n-1}) < F_\lambda(x_1, \dots, x_{n-1})$, that is, x_i ($i = 5, \dots, n - 2$) are not greater than 1. Let us show that all vectors $(x_1, \dots, x_{n-1}) \in \Delta$, realizing the minimum of F_λ , have $x_5 = \dots = x_{n-2} = 0$.

Actually, suppose that there is an index $5 \leq i \leq n - 2$ such that $x_i = 1$ and $x_k = 0$ for all $5 \leq k \leq n - 2$, $k \neq i$. In this case, if $x_4 \geq 1$, we can apply t_1 for positions 4 and i and obtain a smaller value for F_λ . Suppose that $x_4 = 0$. Since $x_{n-1} \in \{0, 1\}$, we will separately analyze the two cases: (a) $x_{n-1} = 1$ and (b) $x_{n-1} = 0$.

(a) In this case, $x_{n-1} = x_i = 1$, where $i \geq 5$ and $x_4 = 0$. We can consider different vertices $x, y, p, q, r, s \in V(G)$ such that $d_G(x) = n - 1 \geq 5$, $d_G(y) = i \geq 5$, p, q, r, s , all are adjacent to x and y , respectively. Meanwhile, x and y are adjacent. We have found four cycles $xypx, xyqx, xyrx$, and $xysx$, which contradicts the hypothesis.

(b) If $x_{n-1} = 0$, then $x_3 = 0$, $x_i = 1$ ($5 \leq i \leq n - 2$), and Δ is characterized by the equations $x_1 + x_2 = n - 1$, $x_1 + 2x_2 + i = 2n + 4$, and $x_1 \leq n - 8$. Suppose that $x_3 \geq 1$. In this case, we can use t_1 for position 3 and i and deduce a smaller value for F_λ . Thus $x_1 + x_2 = n - 1$ and $x_1 + 2x_2 + i = 2n + 4$. We can obtain $x_1 = i - 6 \leq n - 8$. Moreover, $x_2 = n - i + 5 \geq 7$. In this case, we can use t_1 for position 2 and i and deduce a smaller value for F_λ .

To sum up, we have $x_5 = \dots = x_{n-2} = 0$ and $x_{n-1} \in \{0, 1\}$. We will separately analyze the two cases: (a') $x_{n-1} = 0$ and (b') $x_{n-1} = 1$ again.

(a') If $x_{n-1} = 0$, then $x_2 + 2x_3 + 3x_4 = n + 4$. This equation does not hold. If all x_2, x_3 , and x_4 are not greater than 2, $x_2 + 2x_3 + 3x_4 \leq 12$, which contradicts the hypothesis $n \geq 6$. If one of x_2, x_3 , and x_4 is greater than 2, then by using t_2 for the corresponding position, we can obtain a smaller value for F_λ .

(b') If $x_{n-1} = 1$, then $x_1 + x_2 + x_3 + x_4 = n - 1$ and $x_1 + 2x_2 + 3x_3 + 4x_4 = n + 5$, namely $x_2 + 2x_3 + 3x_4 = 6$. If $x_4 \geq 3$, then $x_2 + 2x_3 + 3x_4 \geq 9$. If $x_4 = 2$, then $x_2 = x_3 = 0$ and $x_1 = n - 1$, which contradicts the condition (iii) in Lemma 2.2. Thus, $x_4 = 0$ or $x_4 = 1$. We now distinguish the following two subcases:

Subcase 1. If $x_4 = 0$, then $x_3 = 3 - \frac{x_2}{2}$. All its solutions are $x_2 = 0, x_3 = 3$ or $x_2 = 2, x_3 = 2$ or $x_2 = 4, x_3 = 1$ or $x_2 = 6, x_3 = 0$. In the third and fourth cases, by using t_2 for the position 2, we can obtain a smaller value for F_λ . In the second case, by using t_2 for the position 2, we obtain the first case. But the first case cannot be changed, since by using t_2 for the position 3, the vector $(n - 4, 0, 3, 0, \dots, 0, 1)$ can be $(n - 4, 1, 1, 1, 0, \dots, 0, 1)$, which contradicts Lemma 2.2. Thus, $(x_1, x_2, \dots, x_{n-1}) = (n - 4, 0, 3, 0, \dots, 0, 1)$, and the corresponding unique graph is denoted by H .

Subcase 2. If $x_4 = 1$, then $x_2 + 2x_3 = 3$. All its solutions are $x_2 = 1, x_3 = 1$ or $x_2 = 3, x_3 = 0$. The first case should be removed. By a similar reasoning as before, the second case cannot be transferred by t_2 . Meanwhile, the first transformation cannot be used. Otherwise, the sequence can be changed into $(n - 4, 2, 0, 0, 1, 0, \dots, 0, 1)$, which contradicts Lemma 2.2.

Thus, $(x_1, x_2, \dots, x_{n-1}) = (n-5, 3, 0, 1, 0, \dots, 0, 1)$, and the corresponding unique graph is denoted by H' .

We can easily obtain $H_\lambda(H)$ and $H_\lambda(H')$. Thus, $H_\lambda(H) = H_\lambda(H') = (n^2 - n + 24) + 2^\lambda(n^2 + 3n - 28)$. So, case (b) holds. This completes the proof of Theorem 2.5. \square

In [19], the authors proved the following theorem.

Theorem 2.6 *Let $n \geq 4$. If G belongs to the class of connected tricyclic graphs on n vertices, then:*

- (a) *If $n = 4$, then $\min D'(G) = 36$ and the unique extremal graph is isomorphic to K_4 ;*
- (b) *If $n \geq 5$, then $\min D'(G) = 3n^2 + 5n - 32$, and all the extremal graphs are isomorphic to H or H' , where H is obtained by identifying the center of a star $K_{1,n-4}$ with an arbitrary vertex of a complete graph K_4 , and H' is obtained by identifying the center of a star $K_{1,n-5}$ with an arbitrary degree 4 vertex of a graph A_2 , which is shown in Figure 1, respectively.*

If we choose $\lambda = 1$ in Theorem 2.5, then we can obtain the same result (Theorem 3.1 in [19]).

Theorem 2.7 *Let G belong to the class of connected graphs on n vertices, and let λ be a positive integer number, then for every $n \geq 2$, we have*

$$\min H_\lambda(G) = (n^2 - n) + (n^2 - 3n + 2)2^\lambda,$$

and the unique extremal graph is $K_{1,n-1}$.

Proof Since for $n = 2$ the property is trivial, so let $n \geq 3$. If G is a connected graph with n vertices, then $\sum_{i=1}^{n-1} x_i = n$ and $\sum_{i=1}^{n-1} ix_i \geq 2(n-1)$.

We will find the minimum of $F_\lambda(x_1, x_2, \dots, x_{n-1})$ over all natural numbers $x_1, x_2, \dots, x_{n-1} \geq 0$ satisfying $\sum_{i=1}^{n-1} x_i = n$ and $\sum_{i=1}^{n-1} ix_i \geq 2(n-1)$. First, we shall prove that under these conditions, all systems $(x_1, x_2, \dots, x_{n-1})$ of natural numbers reaching $\min F_\lambda(x_1, x_2, \dots, x_{n-1})$ must satisfy (i) and (ii) given below:

- (i) $x_1 + \dots + x_{n-1} = n$;
- (ii) $x_1 + 2x_2 + \dots + (n-1)x_{n-1} = 2n-2$.

Indeed, suppose that $x_1 + 2x_2 + \dots + (n-1)x_{n-1} > 2n-2$. It follows that there exists a smallest index m , $2 \leq m \leq n-1$ such that $x_m > 0$. We shall define the system (y_1, \dots, y_{n-1}) such that $y_{m-1} = x_{m-1} + 1$, $y_m = x_m - 1$ and $y_i = x_i$ for every $i \neq m-1, m$. We have $\sum_{i=1}^{n-1} y_i = n$, $\sum_{i=1}^{n-1} iy_i = \sum_{i=1}^{n-1} ix_i - 1 \geq 2(n-1)$ and $F_\lambda(x_1, x_2, \dots, x_{n-1}) - F_\lambda(y_1, y_2, \dots, y_{n-1}) = 2^\lambda(-2m+n) + (2m-1) > 0$.

Hence F_λ cannot be minimum for systems $(x_1, x_2, \dots, x_{n-1})$, for which $\sum_{i=1}^{n-1} ix_i > 2(n-1)$. Since natural numbers d_1, \dots, d_n are the degrees of the vertices of a tree of order n if and only if $\sum_{i=1}^n d_i = 2n-2$, we can consider that the numbers x_1, x_2, \dots, x_{n-1} are the multiplicities of the degrees $1, \dots, n-1$ of the vertices of a tree T of order n , and we shall denote also $F_\lambda(T)$ instead of $F_\lambda(x_1, x_2, \dots, x_{n-1})$. Hence the domain where F_λ must be minimized is defined by the multiplicities of the degrees $1, \dots, n-1$ of the vertices of a tree T of order n .

If T is not isomorphic to $K_{1,n-1}$, i.e., if $(x_1, x_2, \dots, x_{n-1}) \neq (n-1, 0, \dots, 0, 1)$, then $\text{diam}(T) \geq 3$. Let $z, t \in V(T)$ such that $d(z, t) = \text{diam}(T)$. It follows that $d_T(z) = d_T(t) = 1$,

and denote by u and v , $u \neq v$, the vertices adjacent with t and z , respectively. Suppose that $d_T(u) = m + p$ and $d_T(v) = m$, where $m \geq 2$ and $p \geq 0$.

Let T_1 be the tree of order n defined in the following way: $V(T_1) = V(T)$ and $E(T_1) = E(T) \cup \{zu\} - \{zv\}$. By replacing T with T_1 , the degrees of u and v change: $d_{T_1}(u) = m + p + 1$, $d_{T_1}(v) = m - 1$, and the degrees of $n - 2$ vertices remain unchanged. We get $F_\lambda(T) - F_\lambda(T_1) = (2^\lambda - 1)(2p + 2) > 0$, and $F_\lambda(T)$ cannot be minimum.

It follows that $F_\lambda(T)$ is minimum only if $T \cong K_{1,n-1}$, i.e., $x_1 = n - 1$, $x_2 = \dots = x_{n-2} = 0$ and $x_{n-1} = 1$, when $F_\lambda(n - 1, 0, \dots, 0, 1) = (n^2 - n) + (n^2 - 3n + 2)2^\lambda$. By concluding, $\min F_\lambda(x_1, \dots, x_{n-1}) = (n^2 - n) + (n^2 - 3n + 2)2^\lambda$, and equality holds if and only if $x_1 = n - 1$, $x_2 = \dots = x_{n-2} = 0$ and $x_{n-1} = 1$, i.e., $G \cong K_{1,n-1}$ since $H_\lambda(G) = (n^2 - n) + (n^2 - 3n + 2)2^\lambda$. \square

In [12], the authors proved the following theorem.

Theorem 2.8 *Let G belong to the class of connected graphs on n vertices, then for every $n \geq 2$, we have*

$$\min D'(G) = 3n^2 - 7n + 4,$$

and the unique extremal graph is $K_{1,n-1}$.

If we choose $\lambda = 1$ in Theorem 2.7, then we can obtain the same result (Theorem 2.1 in [12]).

A Moore graph is a graph of diameter k with girth $2k + 1$. Those graphs have the minimum number of vertices possible for a regular graph with given diameter and the maximum degree. We first bring the following results in [20, 21].

Lemma 2.9 [20] *Let G be a connected graph of order $n \geq 2$ and size $m \geq 1$, and let λ be a negative integer. Then $H_\lambda(G) \leq (1 - 2^\lambda)M_1(G) + 2^{\lambda+1}mn - 2^{\lambda+1}m$, and the equality holds if and only if $d \leq 2$, where d is the diameter of G .*

Lemma 2.10 [21] *Let G be a triangle- and quadrangle-free graph on n vertices with radius r . Then $M_1(G) \leq n(n + 1 - r)$. The equality is valid if and only if G is a Moore graph of diameter 2 or $G = C_6$.*

Theorem 2.11 *Let G be a triangle- and quadrangle-free graph on n vertices and m edges with radius r , and let λ be a negative integer. Then $H_\lambda(G) \leq (1 - 2^\lambda)n(n + 1 - r) + 2^{\lambda+1}mn - 2^{\lambda+1}m$. The equality is valid if and only if G is a Moore graph of diameter 2 or $G = C_6$.*

Proof By the lemmas above, we have

$$\begin{aligned} H_\lambda(G) &\leq (1 - 2^\lambda)M_1(G) + 2^{\lambda+1}mn - 2^{\lambda+1}m \\ &\leq (1 - 2^\lambda)n(n + 1 - r) + 2^{\lambda+1}mn - 2^{\lambda+1}m. \end{aligned}$$

The first equality holds if and only if the diameter of G is at most 2, and the second one holds if and only if G is a Moore graph of diameter 2 or $G = C_6$. So, the equality holds if and only if G is a Moore graph of diameter 2 or $G = C_6$. This completes the proof. \square

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank the referee for the valuable comments.

Received: 1 May 2013 Accepted: 29 August 2013 Published: 19 Nov 2013

References

1. Diudea, MV, Gutman, I, Jantschi, L: Molecular Topology. Huntington, New York (2001)
2. Trinajstić, N: Chemical Graph Theory. CRC Press, Boca Raton (1992)
3. Todeschini, R, Consonni, V: Handbook of Molecular Descriptors. Wiley, New York (2000)
4. Wiener, H: Structural determination of the paraffin boiling points. *J. Am. Chem. Soc.* **69**, 17-20 (1947)
5. Alizadeh, Y, Iranmanesh, A, Došlić, T: Additively weighted Harary index of some composite graphs. *Discrete Math.* **313**, 26-34 (2013)
6. Dobrynin, AA, Kochetova, AA: Degree distance of a graph: a degree analogue of the Wiener index. *J. Chem. Inf. Comput. Sci.* **34**, 1082-1086 (1994)
7. Gutman, I: Selected properties of the Schultz molecular topological index. *J. Chem. Inf. Comput. Sci.* **34**, 1087-1089 (1994)
8. Hamzeh, A, Iranmanesh, A, Hossein-Zadeh, S, Diudea, MV: Generalized degree distance of trees, unicyclic and bicyclic graphs. *Stud. Univ. Babeş-Bolyai, Chem.* **4**, 73-85 (2012)
9. Dankelmann, P, Gutman, I, Mukwembli, S, Swart, HC: On the degree distance of a graph. *Discrete Appl. Math.* **157**, 2773-2777 (2009)
10. Hossein-Zadeh, S, Hamzeh, A, Ashrafi, AR: Extremal properties of Zagreb coindices and degree distance of graphs. *Miskolc Math. Notes* **11**, 129-137 (2010)
11. Tomescu, I: Unicyclic and bicyclic graphs having minimum degree distance. *Discrete Appl. Math.* **156**, 125-130 (2008)
12. Tomescu, I: Some extremal properties of the degree distance of a graph. *Discrete Appl. Math.* **98**, 159-163 (1999)
13. Harary, F: Graph Theory. Addison-Wesley, Reading (1969)
14. Ashrafi, AR, Došlić, T, Hamzeh, A: Extremal graphs with respect to the Zagreb coindices. *MATCH Commun. Math. Comput. Chem.* **65**, 85-92 (2011)
15. Bucicovschi, O, Cioabă, SM: The minimum degree distance of graphs of given order and size. *Discrete Appl. Math.* **156**, 3518-3521 (2008)
16. Khormali, O, Iranmanesh, A, Gutman, I, Ahmadi, A: Generalized Schultz index and its edge versions. *MATCH Commun. Math. Comput. Chem.* **64**, 783-798 (2010)
17. Moon, JW: Counting Labelled Trees. Canadian Mathematical Monographs, vol. 1. Clowes, London (1970)
18. Senior, JK: Partitions and their representative graphs. *Am. J. Math.* **73**, 663-689 (1951)
19. Zhu, W, Hu, S, Ma, H: A note on tricyclic graphs with minimum degree distance. *Discrete Math. Algorithms Appl.* **3**, 25-32 (2011)
20. Hamzeh, A, Iranmanesh, A, Hossein-Zadeh, S: On the generalization of degree distance of graphs (submitted)
21. Yamaguchi, S: Estimating the Zagreb indices and the spectral radius of triangle-and quadrangle-free connected graphs. *Chem. Phys. Lett.* **458**, 396-398 (2008)

10.1186/1029-242X-2013-548

Cite this article as: Hamzeh et al.: Minimum generalized degree distance of n -vertex tricyclic graphs. *Journal of Inequalities and Applications* 2013, **2013**:548

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com