

RESEARCH

Open Access

Berinde-Borcut tripled fixed point theorem in partially ordered (intuitionistic) fuzzy normed spaces

Poom Kumam¹, Juan Martínez-Moreno^{2*}, Antonio-Francisco Roldán-López-de-Hierro² and Concepción Roldán-López-de-Hierro³

*Correspondence:

jmmoreno@ujaen.es

²Department of Mathematics,
University of Jaén, Campus Las
Lagunillas, s/n, Jaén, 23071, Spain
Full list of author information is
available at the end of the article

Abstract

In this paper, we prove some tripled fixed point theorems in fuzzy normed spaces. Our results improve and restate the proof lines of the main results given in the paper (Abbas *et al.* in *Fixed Point Theory Appl.* 2012:187, 2012).

Keywords: fuzzy metric space; tripled fixed point; intuitionistic space

1 Introduction

Once the notion of *coupled fixed point* was given by Gnana Bhaskar and Lakshmikantham in [1], the theory of multidimensional fixed points has attracted much attention (see, for instance, [2–8]), specially in the tripled case (see [9–17]).

Recently, many authors have shown the existence of tripled fixed points and common tripled fixed points for some contractions in cone metric spaces, partially ordered metric spaces, fuzzy metric spaces, fuzzy normed spaces, intuitionistic fuzzy normed spaces and others. Especially in [18], Abbas *et al.* proved some tripled fixed point theorem for contractive mappings in partially complete intuitionistic fuzzy normed spaces. But the authors found some mistakes in the proof lines of their main result. In this paper we give a corrected version of the main theorem.

A *t*-norm (resp., a *t*-conorm) is a mapping $*$: $[0, 1]^2 \rightarrow [0, 1]$ (resp., \diamond : $[0, 1]^2 \rightarrow [0, 1]$) that is associative, commutative, and non-decreasing in both arguments and has 1 (resp., 0) as identity.

Definition 1 ([19, 20]) For any $a \in [0, 1]$, let the sequence $\{*^n a\}_{n=1}^{\infty}$ be defined by $*^1 a = a$ and $*^n a = (*^{n-1} a) * a$. Then a *t*-norm $*$ is said to be of *H*-type if the sequence $\{*^n a\}_{n=1}^{\infty}$ is equicontinuous at $a = 1$.

Definition 2 A *fuzzy normed space* (briefly, FNS) is a triple $(X, \mu, *)$, where X is a vector space, $*$ is a continuous *t*-norm and $\mu : X \times (0, \infty) \rightarrow [0, 1]$ is a fuzzy set such that, for all $x, y \in X$ and $t, s > 0$,

$$(F1) \quad \mu(x, t) > 0;$$

$$(F2) \quad \mu(x, t) = 1 \text{ for all } t > 0 \text{ if and only if } x = 0;$$

$$(F3) \quad \mu(\alpha x, t) = \mu(x, \frac{t}{|\alpha|}) \text{ for all } \alpha \neq 0;$$

- (F4) $\mu(x, t) * \mu(y, s) \leq \mu(x + y, t + s)$;
 (F5) $\mu(x, \cdot) : (0, \infty) \rightarrow [0, 1]$ is continuous;
 (F6) $\lim_{t \rightarrow \infty} \mu(x, t) = 1$ and $\lim_{t \rightarrow 0} \mu(x, t) = 0$.

Using the continuous t -norms and t -conorms, Saadati and Park [21] introduced the concept of an intuitionistic fuzzy normed space.

Definition 3 ([21, 22]) An intuitionistic fuzzy normed space (briefly, IFNS) is a 5-tuple $(X, \mu, \nu, *, \diamond)$ where X is a vector space, $*$ is a continuous t -norm, \diamond is a continuous t -conorm and $\mu, \nu : X \times (0, \infty) \rightarrow [0, 1]$ are fuzzy sets such that, for all $x, y \in X$ and $t, s > 0$,

- (IF1) $\mu(x, t) + \nu(x, t) \leq 1$;
 (IF2) $\mu(x, t) > 0$ and $\nu(x, t) < 1$;
 (IF3) $\mu(x, t) = 1$ for all $t > 0$ if and only if $x = 0$ if and only if $\nu(x, t) = 0$ for all $t > 0$;
 (IF4) $\mu(\alpha x, t) = \mu(x, \frac{t}{|\alpha|})$ and $\nu(\alpha x, t) = \nu(x, \frac{t}{|\alpha|})$ for all $\alpha \neq 0$;
 (IF5) $\mu(x, t) * \mu(y, s) \leq \mu(x + y, t + s)$ and $\nu(x, t) \diamond \nu(y, s) \geq \nu(x + y, t + s)$;
 (IF6) $\mu(x, \cdot), \nu(x, \cdot) : (0, \infty) \rightarrow [0, 1]$ are continuous;
 (IF7) $\lim_{t \rightarrow \infty} \mu(x, t) = 1 = \lim_{t \rightarrow 0} \nu(x, t)$ and $\lim_{t \rightarrow 0} \mu(x, t) = 0 = \lim_{t \rightarrow \infty} \nu(x, t)$.

Obviously, if $(X, \mu, \nu, *, \diamond)$ is a IFNS, then $(X, \mu, *)$ is a FNS. We refer to this space as its *support*.

Lemma 4 $\mu(x, \cdot)$ is a non-decreasing function on $(0, \infty)$ and $\nu(x, \cdot)$ is a non-increasing function on $(0, \infty)$.

Some properties and examples of IFNS and the concepts of *convergence* and a *Cauchy sequence* in IFNS are given in [21].

Definition 5 Let $(X, \mu, \nu, *, \diamond)$ be an IFNS.

- (1) A sequence $\{x_n\} \subset X$ is called a *Cauchy sequence* if, for any $\epsilon > 0$ and $t > 0$, there exists $n_0 \in \mathbb{N}$ such that $\mu(x_n - x_m, t) > 1 - \epsilon$ and $\nu(x_n - x_m, t) < \epsilon$ for all $n, m \geq n_0$.
- (2) A sequence $\{x_n\} \subset X$ is said to be *convergent to a point* $x \in X$, denoted by $x_n \rightarrow x$ or by $\lim_{n \rightarrow \infty} x_n = x$, if, for any $\epsilon > 0$ and $t > 0$, there exists $n_0 \in \mathbb{N}$ such that $\mu(x_n - x, t) > 1 - \epsilon$ and $\nu(x_n - x, t) < \epsilon$ for all $n \geq n_0$.
- (3) An IFNS in which every Cauchy sequence is convergent is said to be *complete*.

Definition 6 ([7]) Let $F : X^3 \rightarrow X$ and $g : X \rightarrow X$ be two mappings.

- We say that F and g are *commuting* if $gF(x, y, z) = F(gx, gy, gz)$ for all $x, y, z \in X$.
- A point $(x, y, z) \in X^3$ is called a *tripled coincidence point of the mappings F and g* if $F(x, y, z) = gx, F(y, x, y) = gy$ and $F(z, y, x) = gz$. If g is the identity, (x, y, z) is called a *tripled fixed point of F* .
- If (X, \sqsubseteq) is a partially ordered set, then F is said to have the *mixed g -monotone property* if it verifies the following properties:

$$\begin{aligned} x_1, x_2 \in X, \quad gx_1 \sqsubseteq gx_2 &\implies F(x_1, y, z) \sqsubseteq F(x_2, y, z), \quad \forall y \in X, \\ y_1, y_2 \in X, \quad gy_1 \sqsubseteq gy_2 &\implies F(x, y_1, z) \supseteq F(x, y_2, z), \quad \forall x \in X, \\ z_1, z_2 \in X, \quad gz_1 \sqsubseteq gz_2 &\implies F(x, y, z_1) \sqsubseteq F(x, y, z_2), \quad \forall x \in X. \end{aligned}$$

If g is the identity mapping, then F is said to have the *mixed monotone property*.

- If (X, \sqsubseteq) is a partially ordered set, then X is said to have the *sequential g -monotone property* if it verifies the following properties:

(B1) If $\{x_n\}$ is a non-decreasing sequence and $\lim_{n \rightarrow \infty} x_n = x$, then $gx_n \sqsubseteq gx$ for all $n \in \mathbb{N}$.

(B2) If $\{x_n\}$ is a non-increasing sequence and $\lim_{n \rightarrow \infty} x_n = y$, then $gy_n \supseteq gy$ for all $n \in \mathbb{N}$.

If g is the identity mapping, then X is said to have the *sequential monotone property*.

Definition 7 Let X and Y be two IFNS. A function $f : X \rightarrow Y$ is said to be *continuous at a point* $x_0 \in X$ if, for any sequence $\{x_n\}$ in X converging to x_0 , the sequence $\{f(x_n)\}$ in Y converges to $f(x_0)$. If f is continuous at each $x \in X$, then f is said to be *continuous on X* .

The following lemma proved by Haghi *et al.* [23] is useful for our main results:

Lemma 8 Let X be a nonempty set and $g : X \rightarrow X$ be a mapping. Then there exists a subset $E \subset X$ such that $g(E) = g(X)$ and $g : E \rightarrow X$ is one-to-one.

Definition 9 Let $(X, \mu, \nu, *, \diamond)$ be an IFNS. The pair (μ, ν) is said to satisfy the *n -property* on $X \times (0, \infty)$ if $\lim_{n \rightarrow \infty} [\mu(x, k^n t)]^{n^p} = 1$ and $\lim_{n \rightarrow \infty} [\nu(x, k^n t)]^{n^p} = 0$ whenever $x \in X, k > 1$ and $p > 0$.

In order to state our results, we recall the main result given in [18].

Theorem 10 (Abbas *et al.*, Theorem 2.2) Let (X, \sqsubseteq) be a partially ordered set and suppose that $a \diamond a = a, ab \leq a * b$ for all $a, b \in [0, 1]$. Let $(X, \mu, \nu, *, \diamond)$ be a complete IFNS such that (μ, ν) has the *n -property*. Let $F : X \times X \times X \rightarrow X$ and $g : X \rightarrow X$ be two mappings such that F has the *mixed g -monotone property* and

$$\begin{aligned} \mu(F(x, y, z) - F(u, v, w), kt) &\geq \mu(gx - gu, t) * \mu(gy - gv, t) * \mu(gz - gw, t), \\ \nu(F(x, y, z) - F(u, v, w), kt) &\leq \nu(gx - gu, t) \diamond \nu(gy - gv, t) \diamond \nu(gz - gw, t), \end{aligned}$$

for which $gx \sqsubseteq gu$ and $gy \supseteq gv$ and $gz \sqsubseteq gw$, where $0 < k < 1$. Suppose either

- F is continuous or
- X has the *sequential g -monotone property*.

If there exist $x_0, y_0, z_0 \in X$ such that $gx_0 \sqsubseteq F(x_0, y_0, z_0), gy_0 \supseteq F(y_0, x_0, y_0)$ and $gz_0 \sqsubseteq F(z_0, y_0, x_0)$, then F and g have a *tripled coincidence point*.

2 Comments and revised tripled fixed point theorem

Firstly, we show that the conditions of Theorem 10 are inadequate and, further, the proof lines of Theorem 10 are not correct. We also would like to point out that the results in [18] can be corrected under the appropriate conditions on the t -norm and the FNS. The proof lines of Theorem 10 are not correct (see pp.7 and 8):

$$\mu(x_n - x_m, t) \geq \left[\mu \left(x_0 - x_1, (1-k) \frac{t}{k^n} \right) \right]^m \geq \left[\mu \left(x_0 - x_1, (1-k) \frac{t}{k^n} \right) \right]^{n^p} \rightarrow 1,$$

where $p > 0$ such that $m < n^p$. Hence the sequence $\{x_n\}$ is a Cauchy sequence. This is not correct since the same p would not be valid for all positive integers $m > n \geq n_0$. For example, let $(X, \|\cdot\|)$ be an ordinary normed space, define $\mu(x, t) = \frac{t}{t + \|x\|}$ for any $x \in X$ and $t > 0$ and $a * b = ab$ for all $a, b \in [0, 1]$. Then $(X, \mu, 1 - \mu, *, *)$ is an IFNS. If $k = 1/2$ and $m = 2^n$, we have

$$\left[\mu \left(x_0 - x_1, (1-k) \frac{t}{k^n} \right) \right]^m = \left[\frac{2^{n-1}t}{2^{n-1}t + \|x_0 - x_1\|} \right]^{2^n} \rightarrow e^{-\frac{2\|x_0 - x_1\|}{t}} < 1.$$

Now, by replacing in Theorem 10 the hypothesis that μ satisfies the n -property with the one that the t -norm is of H -type, we state and prove a tripled fixed point theorem as a modification.

Theorem 11 *Let (X, \sqsubseteq) be a partially ordered set and $(X, \mu, *)$ be a complete FNS such that $*$ is of H -type and $a * a \geq a$ for all $a \in [0, 1]$. Let $k \in (0, 1)$ be a number and $F : X \times X \times X \rightarrow X$ be mapping such that F has the mixed monotone property and*

$$\mu(F(x, y, z) - F(u, v, w), kt) \geq \mu(x - u, t) * \mu(y - v, t) * \mu(z - w, t), \tag{1}$$

for which $x \sqsubseteq u, y \sqsupseteq v$ and $z \sqsubseteq w$. Suppose that either:

- (a) F is continuous or
- (b) X has the sequential monotone property.

If there exist $x_0, y_0, z_0 \in X$ such that $x_0 \sqsubseteq F(x_0, y_0, z_0), y_0 \sqsupseteq F(y_0, x_0, y_0)$ and $z_0 \sqsubseteq F(z_0, y_0, x_0)$, then F has a tripled fixed point. Furthermore, if x_0 and y_0 are comparable, then $x = y$, that is, $x = F(x, x)$.

Proof As in [18] starting with $x_0, y_0, z_0 \in X$ such that $x_0 \sqsubseteq F(x_0, y_0, z_0), y_0 \sqsupseteq F(y_0, x_0, y_0)$ and $z_0 \sqsubseteq F(z_0, y_0, x_0)$, one can define inductively three sequences $\{x_n\}, \{y_n\}, \{z_n\} \subset X$ such that $x_{n+1} = F(x_n, y_n, z_n), y_{n+1} = F(y_n, x_n, y_n)$ and $z_{n+1} = F(z_n, y_n, x_n)$.

Define

$$\delta_n(t) = \mu(x_n - x_{n+1}, t) * \mu(y_n - y_{n+1}, t) * \mu(z_n - z_{n+1}, t).$$

Continuing as in [18], we have

$$\begin{aligned} \mu(x_n - x_{n+1}, kt) &\geq \delta_{n-1}(t), & \mu(y_n - y_{n+1}, kt) &\geq \delta_{n-1}(t) \quad \text{and} \\ \mu(z_n - z_{n+1}, kt) &\geq \delta_{n-1}(t) * \delta_{n-1}(t). \end{aligned} \tag{2}$$

Since $a * a \geq a$ for all $a \in [0, 1]$, it follows that

$$\delta_n(kt) = \mu(x_n - x_{n+1}, kt) * \mu(y_n - y_{n+1}, kt) * \mu(z_n - z_{n+1}, kt) \geq \delta_{n-1}(t).$$

This implies that

$$1 \geq \delta_n(t) \geq \delta_{n-1} \left(\frac{t}{k} \right) \geq \delta_{n-2} \left(\frac{t}{k^2} \right) \geq \dots \geq \delta_0 \left(\frac{t}{k^n} \right).$$

Since $\lim_{n \rightarrow \infty} \delta_0 \left(\frac{t}{k^n} \right) = 1$ for all $t > 0$, we have $\lim_{n \rightarrow \infty} \delta_n(t) = 1$ for all $t > 0$.

Now, we claim that, for any $p \geq 1$ and $n \geq 1$,

$$\begin{aligned} \mu(x_n - x_{n+p}, t) &\geq *^p \delta_{n-1}(t - kt), \\ \mu(y_n - y_{n+p}, t) &\geq *^p \delta_{n-1}(t - kt), \quad \text{and} \\ \mu(z_n - z_{n+p}, t) &\geq *^p \delta_{n-1}(t - kt). \end{aligned} \tag{3}$$

In fact, it is obvious for $p = 1$ by (2), $a * a \geq a$ and Lemma 4 since $t/k \geq t - kt$ and δ_{n-1} is non-decreasing. Assume that (3) holds for some $p \geq 1$. By (2), we have

$$\mu(x_n - x_{n+1}, t) \geq \mu(x_n - x_{n+1}, kt) \geq \delta_{n-1}(t)$$

and so

$$\mu(x_n - x_{n+1}, t - kt) \geq \delta_{n-1}(t - kt).$$

Thus, from (1), (3) and $a * a \geq a$, we have

$$\begin{aligned} \mu(x_{n+1} - x_{n+p+1}, kt) &\geq \mu(x_n - x_{n+p}, t) * \mu(y_n - y_{n+p}, t) * \mu(z_n - z_{n+p}, t) \\ &\geq *^p \delta_{n-1}(t - kt). \end{aligned} \tag{4}$$

Hence, by the monotonicity of the t -norm $*$, we have

$$\begin{aligned} \mu(x_n - x_{n+p+1}, t) &= \mu(x_n - x_{n+p+1}, t - kt + kt) \\ &\geq \mu(x_n - x_{n+1}, t - kt) * \mu(x_{n+1} - x_{n+p+1}, kt) \\ &\geq \delta_{n-1}(t - kt) * (*^p \delta_{n-1}(t - kt)) = *^{p+1} \delta_{n-1}(t - kt). \end{aligned}$$

Similarly, we have

$$\begin{aligned} \mu(y_{n+1} - y_{n+p+1}, kt) &\geq *^{p+1} \delta_{n-1}(t - kt) \quad \text{and} \\ \mu(z_{n+1} - z_{n+p+1}, kt) &\geq *^{p+1} \delta_{n-1}(t - kt). \end{aligned} \tag{5}$$

Therefore, by induction, (3) holds for all $p \geq 1$. Suppose that $t > 0$ and $\epsilon \in (0, 1]$ are given. By hypothesis, since $*$ is a t -norm of H -type, there exists $0 < \eta < 1$ such that $*^p(a) > 1 - \epsilon$ for all $a \in (1 - \eta, 1]$ and $p \geq 1$. Since $\lim_{n \rightarrow \infty} \delta_n(t) = 1$, there exists n_0 such that $\delta_n(t - kt) > 1 - \eta$ for all $n \geq n_0$. Hence, from (3), we get

$$\mu(x_n - x_{n+p}, t) > 1 - \epsilon, \quad \mu(y_n - y_{n+p}, t) > 1 - \epsilon, \quad \mu(z_n - z_{n+p}, t) > 1 - \epsilon, \quad \forall n \geq n_0.$$

Therefore, $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ are Cauchy sequences. We can continue as in [18] to complete the proof. \square

Theorem 12 *Let (X, \sqsubseteq) be a partially ordered set and $(X, \mu, *)$ be a complete FNS such that $*$ is of H -type and $a * a \geq a$ for all $a \in [0, 1]$. Let $k \in (0, 1)$ be a number and $F : X \times X \times X \rightarrow$*

X and $g : X \rightarrow X$ be two mappings such that F has the mixed g -monotone property and

$$\begin{aligned}\mu(F(x, y, z) - F(u, v, w), kt) &\geq \mu(gx - gu, t) * \mu(gy - gv, t) * \mu(gz - gw, t), \\ v(F(x, y, z) - F(u, v, w), kt) &\leq v(gx - gu, t) \diamond v(gy - gv, t) \diamond v(gz - gw, t),\end{aligned}$$

for which $gx \sqsubseteq gu$ and $gy \sqsupseteq gv$ and $gz \sqsubseteq gw$, where $0 < k < 1$. Suppose either

- (a) F is continuous or
- (b) X has the sequential g -monotone property.

If there exist $x_0, y_0, z_0 \in X$ such that $gx_0 \sqsubseteq F(x_0, y_0, z_0)$, $gy_0 \sqsupseteq F(y_0, x_0, y_0)$ and $gz_0 \sqsubseteq F(z_0, y_0, x_0)$, then F and g have a tripled coincidence point.

Proof As in Theorem 2.2 in [18]. □

Of course, all the results are valid if X is intuitionistic.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally and significantly in writing this paper. All authors read and approved the final manuscript.

Author details

¹Department of Mathematics, Faculty of Science, King Mongkut's University of Technology Thonburi (KMUTT), Bang Mod, Thung Khru, Bangkok, Thailand. ²Department of Mathematics, University of Jaén, Campus Las Lagunillas, s/n, Jaén, 23071, Spain. ³Department of Statistics and Operations Research, University of Granada, Granada, Spain.

Acknowledgements

The first author was supported by the Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission (Under NRU-CSEC Project No. NRU56000508). The last three authors have been partially supported by Junta de Andalucía, by projects FQM-268, FQM-178 and FQM-235 of the Andalusian CICYE.

Received: 1 October 2013 Accepted: 8 January 2014 Published: 30 Jan 2014

References

1. Gnana Bhaskar, T, Lakshmikantham, V: Fixed point theorems in partially ordered metric spaces and applications. *Nonlinear Anal.* **65**, 1379-1393 (2006)
2. Berinde, V, Borcut, M: Tripled fixed point theorems for contractive type mappings in partially ordered metric spaces. *Nonlinear Anal.* **74**, 4889-4897 (2011)
3. Borcut, M, Berinde, V: Tripled coincidence theorems for contractive type mappings in partially ordered metric spaces. *Appl. Math. Comput.* **218**, 5929-5936 (2012)
4. Aydi, H, Karapinar, E, Postolache, M: Tripled coincidence theorems for weak φ -contractions in partially ordered metric spaces. *Fixed Point Theory Appl.* **2012**, 44 (2012)
5. Karapinar, E, Luong, N: Quadruple fixed point theorems for nonlinear contractions. *Comput. Math. Appl.* **64**, 1839-1848 (2012)
6. Karapinar, E, Berinde, V: Quadruple fixed point theorems for nonlinear contractions in partially ordered metric spaces. *Banach J. Math. Anal.* **6**, 74-89 (2012)
7. Roldán, A, Martínez-Moreno, J, Roldán, C: Multidimensional fixed point theorems in partially ordered complete metric spaces. *J. Math. Anal. Appl.* **396**, 536-545 (2012)
8. Roldán, A, Martínez-Moreno, J, Roldán, C: Multidimensional coincidence point results for compatible mappings in partially ordered fuzzy metric spaces. *Fuzzy Sets Syst.* (2013). doi:10.1016/j.fss.2013.10.009
9. Aydi, H, Karapinar, E, Shatanawi, W: Tripled fixed point results in generalized metric spaces. *J. Appl. Math.* **2012**, 314279 (2012)
10. Karapinar, E, Aydi, H, Mustafa, Z: Some tripled coincidence point theorems for almost generalized contractions in ordered metric spaces. *Tamkang J. Math.* **44**, 233-251 (2013)
11. Aydi, H, Karapinar, E, Radenovic, S: Tripled coincidence fixed point results for Boyd-Wong and Matkowski type contractions. *Rev. R. Acad. Cienc. Exactas Fis. Nat., Ser. A Mat.* **107**, 339-353 (2013)
12. Aydi, H, Karapinar, E, Shatanawi, W: Tripled common fixed point results for generalized contractions in ordered generalized metric spaces. *Fixed Point Theory Appl.* **2012**, 101 (2012)
13. Aydi, H, Karapinar, E: New Meir-Keeler type tripled fixed point theorems on ordered partial metric spaces. *Math. Probl. Eng.* **2012**, 409872 (2012)
14. Aydi, H, Karapinar, E, Shatanawi, W: Tripled fixed point results in generalized metric spaces. *J. Appl. Math.* **2012**, 314279 (2012)

15. Abbas, M, Aydi, H, Karapinar, E: Tripled fixed points of multi-valued nonlinear contraction mappings in partially ordered metric spaces. *Abstr. Appl. Anal.* **2011**, 812690 (2011)
16. Aydi, H, Karapinar, E: Tripled coincidence point theorems for a class of contractions in ordered metric spaces. *Acta Univ. Apulensis* **32**, 193-205 (2012)
17. Aydi, H, Karapinar, E: Tripled fixed points in ordered metric spaces. *Bull. Math. Anal. Appl.* **4**, 197-207 (2012)
18. Abbas, M, Ali, B, Sintunavarat, W, Kumam, P: Tripled fixed point and tripled coincidence point theorems in intuitionistic fuzzy normed spaces. *Fixed Point Theory Appl.* **2012**, 187 (2012)
19. Hadžić, O, Pap, E: *Fixed Point Theory in Probabilistic Metric Spaces*. Kluwer Academic, Dordrecht (2001)
20. Roldán, A, Martínez-Moreno, J, Roldán, C: Tripled fixed point theorem in fuzzy metric spaces and applications. *Fixed Point Theory Appl.* **2013**, 29 (2013)
21. Saadati, R, Park, J: On the intuitionistic fuzzy topological spaces. *Chaos Solitons Fractals* **27**, 231-344 (2006)
22. Roldán, A, Martínez-Moreno, J, Roldán, C: On interrelationships between fuzzy metric structures. *Iran. J. Fuzzy Syst.* **10**, 133-150 (2013)
23. Haghi, R, Rezapour, H, Shahzad, N: Some fixed point generalizations are not real generalizations. *Nonlinear Anal.* **74**, 1799-1803 (2011)

10.1186/1029-242X-2014-47

Cite this article as: Kumam et al.: Berinde-Borcut tripled fixed point theorem in partially ordered (intuitionistic) fuzzy normed spaces. *Journal of Inequalities and Applications* 2014, **2014**:47

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
