

RESEARCH

Open Access



New results for the upper bounds of the distance between adjacent zeros of first-order differential equations with several variable delays

Emad R. Attia^{1,3} and Bassant M. El-Matary^{2,3*}

*Correspondence:

b.elmatary@qu.edu.sa;
bassantmarof@yahoo.com

²Department of Mathematics,
College of Science and Arts,
Al-Badaya, Qassim University,
Buraidah 51951, Saudi Arabia

³Department of Mathematics,
Faculty of Science, Damietta
University, New Damietta 34517,
Egypt

Full list of author information is
available at the end of the article

Abstract

The distance between consecutive zeros of a first-order differential equation with several variable delays is studied. Here, we show that the distribution of zeros of differential equations with variable delays is not an easy extension of the case of constant delays. We obtain new upper bounds for the distance between zeros of all solutions of a differential equation with several delays, which extend and improve some existing results. Two illustrative examples are given to show the advantages of the proposed results over the known ones.

Mathematics Subject Classification: 34K11; 34K06

Keywords: Differential equations; Variable delays; Distance between zeros; Oscillation

1 Introduction

Consider the differential equation with several variable delays

$$x'(t) + \sum_{j=1}^n a_j(t)x(g_j(t)) = 0, \quad t \geq t_0, \quad (1)$$

where $a_j, g_j \in C([t_0, \infty), [0, \infty))$, $g_j(t)$ is a strictly increasing function such that $g_j(t) \leq t$, $\lim_{t \rightarrow \infty} g_j(t) = \infty$, $j = 1, 2, \dots, n$. We make use of the following notation:

$$h_i(t) = \max_{1 \leq j \leq i} g_j(t), \quad w_i(t) = \min_{1 \leq j \leq i} g_j(t), \quad i = 1, 2, \dots, n.$$

Therefore,

$$h_j^{-k}(t) \geq h_i^{-k}(t) \quad \text{and} \quad w_j^{-k}(t) \leq w_i^{-k}(t), \quad i \geq j, i, j = 1, 2, \dots, n, k = 1, 2, \dots,$$

where $h_j^{-1}(t)$ and $w_j^{-1}(t)$ are the inverse of the functions $h_j(t)$ and $w_j(t)$, $j = 1, 2, \dots, n$.

© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Consequently,

$$\max_{1 \leq j \leq i} w_j^{-k}(t) = w_i^{-k}(t) \quad \text{and} \quad \max_{1 \leq j \leq i} h_j^{-k}(t) = h_1^{-k}(t), \quad i = 1, 2, \dots, n, k = 1, 2, \dots$$

Let $t^* \geq t_0$ and $x(t)$ be a continuous function on $[t^*, \infty)$. The function $x(t)$ is said to be a solution of Eq. (1) on $[t^*, \infty)$ if $x(t)$ is continuously differentiable on $[w_n^{-1}(t^*), \infty)$ and satisfying Eq. (1) for $t \geq w_n^{-1}(t^*)$. Any solution of Eq. (1) is called oscillatory if it has arbitrarily large zeros; otherwise, it is called nonoscillatory. Equation (1) is called oscillatory if all its solutions are oscillatory; otherwise, it is called nonoscillatory.

The oscillation theory of delay differential equations has received a great deal of attention in recent years; see the monographs [1, 2, 13–15] and the papers [3–12, 16–27] for more details. Many efforts have been made to establish sufficient and/or necessary oscillation criteria for Eq. (1); see [1, 3, 9, 11, 13, 15, 17]. In oscillation theory, the distribution of zeros of delay differential equations has always been an important problem. In this topic, not only is the existence of zeros demonstrated, but efforts are also being made to determine their locations. In fact, the study of the distribution of zeros raises many challenges. This explains the few studies that concern the distance between zeros compared to the oscillation.

Many upper bounds for the distance between consecutive zeros of the delay differential equations

$$x'(t) + a(t)x(t - \sigma) = 0, \quad t \geq t_0, \tag{2}$$

and

$$x'(t) + a(t)x(g(t)) = 0, \quad t \geq t_0, \tag{3}$$

where $\sigma > 0$, $a, g \in C([t_0, \infty), [0, \infty))$, $g(t)$ is a strictly increasing function such that $\lim_{t \rightarrow \infty} g(t) = \infty$, have been obtained by [6–11, 17, 18, 20–27]. Further, some results concerning the lower bounds for the distance between consecutive zeros of all solutions of Eqs. (2) and (3) were investigated in [6–10, 17]. For example, Barr [6] showed that the lower bound of the distance between zeros of an oscillatory solution of Eq. (3) goes to infinity when $t - g(t)$ is not bounded. Therefore, we will restrict our attention to the case when $t - g_j(t) < \infty, j = 1, 2, \dots, n$. In this work, we obtain new upper bounds for the distance between consecutive zeros of all solutions of Eq. (1), which would improve the above-mentioned ones. We conclude by providing two illustrative examples to show the applicability and importance of some of our findings.

2 Main results

Let $t_1 \geq t_0$ and $D_{t_1}(x)$ be the upper bound of the distance between consecutive zeros of all solutions of Eq. (1) on the interval $[t_1, \infty)$. Throughout this paper, it is assumed that

$$\sup_{t \geq t_1} \{t - g_j(t)\} < \infty \quad \text{for } j = 1, 2, \dots, n.$$

Let $r \in \{1, 2, \dots, n\}$ and the sequence $\{R^k(\eta_r)\}_{k \geq 0}$ be defined by $R^0(\eta_r) = 1$ and

$$\begin{aligned}
 R^1(\eta_r) &= \frac{1}{1 - \eta_r}, \\
 R^k(\eta_r) &= \frac{1}{1 - \eta_r - \frac{1}{2}\eta_r^2 R^{k-1}(\eta_r)}, \quad k = 2, 3, \dots,
 \end{aligned}
 \tag{4}$$

where

$$\int_{h_r(t)}^t \sum_{j=1}^r a_j(s) ds \geq \eta_r \quad \text{for } t \geq h_r^{-1}(t_1).$$

Lemma 2.1 *Let $k \in \mathbb{N}_0$, $r \in \{1, 2, \dots, n\}$ and $x(t)$ be a solution of Eq. (1) such that $x(t) > 0$ on $[T_0, T_1]$, $T_0 \geq t_1$, $T_1 \geq h_r^{-1}(w_r^{-k}(w_n^{-1}(T_0)))$. Then*

$$\frac{x(h_r(t))}{x(t)} \geq R^k(\eta_r) \quad \text{for } t \in [h_r^{-(k-1)}(w_r^{-k}(w_n^{-1}(T_0))), T_1],
 \tag{5}$$

where $w_r^0(T_0) = T_0$.

Proof Since $x(t) > 0$ on $[T_0, T_1]$, it follows from Eq. (1) that $x'(t) \leq 0$ on $[w_n^{-1}(T_0), T_1]$, and hence

$$\frac{x(h_r(t))}{x(t)} \geq 1 = R^0(\eta_r) \quad \text{for } t \in [h_r^{-1}(w_n^{-1}(T_0)), T_1].$$

In view of Eq. (1) and the positivity of $x(t)$ on $[T_0, T_1]$, we have

$$x'(t) + \sum_{j=1}^r a_j(t)x(g_j(t)) \leq 0 \quad \text{for } t \in [w_n^{-1}(T_0), T_1].
 \tag{6}$$

Integrating from $h_r(t)$ to t , we get

$$x(t) - x(h_r(t)) + \int_{h_r(t)}^t \sum_{j=1}^r a_j(s)x(g_j(s)) ds \leq 0 \quad \text{for } t \in [h_r^{-1}(w_n^{-1}(T_0)), T_1].
 \tag{7}$$

Since $h_r(t) \geq g_j(t)$, so $h_r(t) \geq g_j(s)$ for $h_r(t) \leq s \leq t$, $j = 1, 2, \dots, r$, it follows from (6) that

$$x(g_j(s)) \geq x(h_r(t)) + \int_{g_j(s)}^{h_r(t)} \sum_{j_1=1}^r a_{j_1}(s_1)x(g_{j_1}(s_1)) ds_1, \quad t \in [h_r^{-1}(w_n^{-1}(w_n^{-1}(T_0))), T_1].$$

Substituting into (7), we obtain

$$\begin{aligned}
 &x(t) - x(h_r(t)) + x(h_r(t)) \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) ds \\
 &+ \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) \int_{g_j(s)}^{h_r(t)} \sum_{j_1=1}^r a_{j_1}(s_1)x(g_{j_1}(s_1)) ds_1 ds \leq 0
 \end{aligned}
 \tag{8}$$

for $t \in [h_r^{-1}(w_r^{-1}(w_n^{-1}(T_0))), T_1]$. Therefore,

$$x(t) - x(h_r(t)) + x(h_r(t)) \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) ds \leq 0 \quad \text{for } t \in [h_r^{-1}(w_r^{-1}(w_n^{-1}(T_0))), T_1].$$

That is,

$$\begin{aligned} \frac{x(h_r(t))}{x(t)} &\geq \frac{1}{1 - \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) ds} \geq \frac{1}{1 - \eta_r} \\ &= R^1(\eta_r) \quad \text{for } t \in [h_r^{-1}(w_r^{-1}(w_n^{-1}(T_0))), T_1]. \end{aligned} \tag{9}$$

Also, since $h_r^2(t) \geq g_{j_1}(s_1)$ for $h_r(t) \leq s \leq t$, $g_j(s) \leq s_1 \leq h_r(t)$, $j, j_1 = 1, 2, \dots, r$. Then

$$x(g_{j_1}(s_1)) \geq x(h_r^2(t)) + \int_{g_{j_1}(s_1)}^{h_r^2(t)} \sum_{j_2=1}^r a_{j_2}(s_2) x(g_{j_2}(s_2)) ds_2, \quad g_{j_1}(s_1) \leq s_2 \leq h_r^2(t)$$

for $t \in [h_r^{-1}(w_r^{-2}(w_n^{-1}(T_0))), T_1]$. From this and (8), it follows that

$$\begin{aligned} &x(t) - x(h_r(t)) + x(h_r(t)) \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) ds \\ &+ x(h_r^2(t)) \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) \int_{g_j(s)}^{h_r(t)} \sum_{j_1=1}^r a_{j_1}(s_1) ds_1 ds \\ &+ \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) \int_{g_j(s)}^{h_r(t)} \sum_{j_1=1}^r a_{j_1}(s_1) \int_{g_{j_1}(s_1)}^{h_r^2(t)} \sum_{j_2=1}^r a_{j_2}(s_2) x(g_{j_2}(s_2)) ds_2 ds_1 ds \leq 0 \end{aligned} \tag{10}$$

for $t \in [h_r^{-1}(w_r^{-2}(w_n^{-1}(T_0))), T_1]$. Using the positivity of $x(t)$ on $[T_0, T_1]$, we have

$$\begin{aligned} &x(t) - x(h_r(t)) + x(h_r(t)) \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) ds \\ &+ x(h_r^2(t)) \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) \int_{g_j(s)}^{h_r(t)} \sum_{j_1=1}^r a_{j_1}(s_1) ds_1 ds \leq 0 \end{aligned}$$

for $t \in [h_r^{-1}(w_r^{-2}(w_n^{-1}(T_0))), T_1]$. Therefore,

$$\frac{x(h_r(t))}{x(t)} \geq \frac{1}{1 - \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) ds - \frac{x(h_r^2(t))}{x(h_r(t))} \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) \int_{g_j(s)}^{h_r(t)} \sum_{j_1=1}^r a_{j_1}(s_1) ds_1 ds} \tag{11}$$

for $t \in [h_r^{-1}(w_r^{-2}(w_n^{-1}(T_0))), T_1]$. Clearly,

$$\begin{aligned} \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) \int_{g_j(s)}^{h_r(t)} \sum_{j_1=1}^r a_{j_1}(s_1) ds_1 ds &\geq \int_{h_r(t)}^{\bar{t}} \sum_{j=1}^r a_j(s) \int_{g_j(s)}^s \sum_{j_1=1}^r a_{j_1}(s_1) ds_1 ds \\ &\quad - \int_{h_r(t)}^{\bar{t}} \sum_{j=1}^r a_j(s) \int_{h_r(t)}^s \sum_{j_1=1}^r a_{j_1}(s_1) ds_1 ds \\ &\geq \eta_r^2 - \int_{h_r(t)}^{\bar{t}} \sum_{j=1}^r a_j(s) \int_{h_r(t)}^s \sum_{j_1=1}^r a_{j_1}(s_1) ds_1 ds, \end{aligned} \tag{12}$$

where $\bar{t} \in (h_r(t), t]$ such that $\int_{h_r(t)}^{\bar{t}} \sum_{j=1}^r a_j(s) ds = \eta_r$. It is easy to see that (see [13, Lemma 2.1.3])

$$\int_{h_r(t)}^{\bar{t}} \sum_{j=1}^r a_j(s) \int_{h_r(t)}^s \sum_{j_1=1}^r a_{j_1}(s_1) ds_1 ds = \frac{1}{2} \left(\int_{h_r(t)}^{\bar{t}} \sum_{j=1}^r a_j(s) ds \right)^2 = \frac{1}{2} \eta_r^2.$$

From this and (12), we get

$$\int_{h_r(t)}^t \sum_{j=1}^r a_j(s) \int_{g_j(s)}^{h_r(t)} \sum_{j_1=1}^r a_{j_1}(s_1) ds_1 ds \geq \frac{1}{2} \eta_r^2.$$

Substituting into (11), we have

$$\frac{x(h_r(t))}{x(t)} \geq \frac{1}{1 - \eta_r - \frac{1}{2} \eta_r^2 \frac{x(h_r^2(t))}{x(h_r(t))}} \quad \text{for } t \in [h_r^{-1}(w_r^{-2}(w_n^{-1}(T_0))), T_1]. \tag{13}$$

In view of (9), we have

$$\frac{x(h_r^2(t))}{x(h_r(t))} \geq R^1(\eta_r) \quad \text{for } t \in [h_r^{-2}(w_r^{-1}(w_n^{-1}(T_0))), T_1].$$

This together with (13) implies that

$$\frac{x(h_r(t))}{x(t)} \geq \frac{1}{1 - \eta_r - \frac{1}{2} \eta_r^2 R^1(\eta_r)} = R^2(\eta_r) \quad \text{for } t \in [h_r^{-1}(w_r^{-2}(w_n^{-1}(T_0))), T_1].$$

Therefore,

$$\frac{x(h_r^2(t))}{x(h_r(t))} \geq R^2(\eta_r) \quad \text{for } t \in [h_r^{-2}(w_r^{-2}(w_n^{-1}(T_0))), T_1].$$

From this and (13), we get

$$\frac{x(h_r(t))}{x(t)} \geq \frac{1}{1 - \eta_r - \frac{1}{2} \eta_r^2 R^2(\eta_r)} = R^3(\eta_r)$$

for $t \in [h_r^{-1}(w_r^{-3}(w_n^{-1}(T_0))), T_1] \subseteq [h_r^{-2}(w_r^{-2}(w_n^{-1}(T_0))), T_1]$.

Repeating this procedure k times, we obtain (5). The proof is complete. □

Let $r \in \{1, 2, \dots, n\}$ and the sequence $\{B_{j,r}^i(s, t)\}_{i \geq 1, j = 1, 2, \dots, r}$ be defined by

$$B_{j,r}^1(s, t) = a_j(s), \quad h_r(t) \leq s \leq t \text{ for } t \geq w_r^{-1}(t_1)$$

$$B_{j,r}^i(s, t) = a_j(s) \int_{g_j(s)}^{h_r(t)} \sum_{j_1=1}^r B_{j_1,r}^{i-1}(s_1, h_r(t)) ds_1, \quad h_r(t) \leq s \leq t, i = 2, 3, \dots$$

for $t \geq w_r^{-i}(t_1)$.

Theorem 2.1 *Assume that $k \in \mathbb{N}$ and $r \in \{1, 2, \dots, n\}$. If*

$$\sum_{i=1}^k \prod_{j=2}^i R^{k+1-j}(\eta_r) \int_{h_r(t)}^t \sum_{j=1}^r B_{j,r}^i(s, h_r(t)) ds \geq 1 \quad \text{for all } t \geq w_r^{-k-1}(t_1),$$

then Eq. (1) oscillates and $D_{t_1}(x) \leq \sup_{t \geq t_1} \{h_r^{-1}(w_r^{-k}(w_n^{-1}(t))) - t\}$.

Proof Suppose the contrary, let $x(t)$ be a positive solution of Eq. (1) on $[T_0, T_1]$, $T_0 \geq t_1$, $T_1 > h_r^{-1}(w_r^{-k}(w_n^{-1}(T_0)))$. Using a similar argument as in the proof of Lemma 2.1, we obtain (10). That is,

$$x(t) - x(h_r(t)) + \sum_{i=1}^2 x(h_r^i(t)) \int_{h_r(t)}^t \sum_{j=1}^r B_{j,r}^i(s, h_r(t)) ds$$

$$+ \int_{h_r(t)}^t \sum_{j=1}^r a_j(s) \int_{g_j(s)}^{h_r(t)} \sum_{j_1=1}^r a_{j_1}(s_1) \int_{g_{j_1}(s_1)}^{h_r^2(t)} \sum_{j_2=1}^r a_{j_2}(s_2) x(g_{j_2}(s_2)) ds_2 ds_1 ds \leq 0,$$

for $t \in [h_r^{-1}(w_r^{-2}(w_n^{-1}(T_0))), T_1]$, where $h_r^1(t) = h_r(t)$. It follows that

$$x(t) - x(h_r(t)) + \sum_{i=1}^2 \int_{h_r(t)}^t x(h_r^i(t)) \sum_{j=1}^r B_{j,r}^i(s, h_r(t)) ds \leq 0$$

for $t \in [h_r^{-1}(w_r^{-2}(w_n^{-1}(T_0))), T_1]$. By repeating this argument k times, we get

$$x(t) - x(h_r(t)) + \sum_{i=1}^k \int_{h_r(t)}^t x(h_r^i(t)) \sum_{j=1}^r B_{j,r}^i(s, h_r(t)) ds \leq 0 \tag{14}$$

for $t \in [h_r^{-1}(w_r^{-k}(w_n^{-1}(T_0))), T_1]$. Since

$$x(h_r^i(t)) = \left(\prod_{j=2}^i \frac{x(h_r^j(t))}{x(h_r^{j-1}(t))} \right) x(h_r(t)), \quad i = 1, 2, \dots$$

By using (5) and the fact that

$$h_r^{j-1}(t) \in [h_r^{-1}(w_r^{-k+(j-1)}(w_n^{-1}(T_0))), T_1]$$

for $t \in [h_r^{-1}(w_r^{-k}(w_n^{-1}(T_0))), T_1]$, we obtain

$$\frac{x(h_r^j(t))}{x(h_r^{j-1}(t))} \geq R^{k+1-j}(\eta_r) \quad \text{for } t \in [h_r^{-1}(w_r^{-k}(w_n^{-1}(T_0))), T_1].$$

Then

$$x(h_r^i(t)) \geq \left(\prod_{j=2}^i R^{k+1-j}(\eta_r) \right) x(h_r(t)), \quad i = 1, 2, \dots$$

Substituting into (14), we get

$$x(t) - x(h_r(t)) + x(h_r(t)) \sum_{i=1}^k \prod_{j=2}^i R^{k+1-j}(\eta_r) \int_{h_r(t)}^t \sum_{j=1}^r B_{j,r}^i(s, h_r(t)) ds \leq 0$$

for $t \in [h_r^{-1}(w_r^{-k}(w_n^{-1}(T_0))), T_1]$, that is,

$$x(t) + \left(\sum_{i=1}^k \prod_{j=2}^i R^{k+1-j}(\eta_r) \int_{h_r(t)}^t \sum_{j=1}^r B_{j,r}^i(s, h_r(t)) ds - 1 \right) x(h_r(t)) \leq 0$$

for $t \in [h_r^{-1}(w_r^{-k}(w_n^{-1}(T_0))), T_1]$. This contradiction completes the proof. □

Theorem 2.2 Assume that $k \in \mathbb{N}_0$. If

$$\prod_{i=1}^n \left(\prod_{j=1}^n \int_{h_i(t)}^t a_j(s) e^{\int_{g_j(s)}^{h_j(t)} \sum_{j_1=1}^n R^{k}(\eta_{j_1}) a_{j_1}(s_1) ds_1} ds \right)^{\frac{1}{n}} \geq \frac{1}{n^n} \quad \text{for } t \geq h_1^{-1}(w_n^{-1}(t_1)), \tag{15}$$

then Eq. (1) oscillates and $D_{t_1}(x) \leq \sup_{t \geq t_1} \{h_1^{-2}(w_n^{-(k+2)}(t)) - t\}$.

Proof Assume that $x(t)$ is a solution of Eq. (1) such that $x(t) > 0$ on $[T_0, T_1]$, $T_0 \geq t_1$, $T_0 > h_1^{-2}(w_n^{-(k+2)}(T_1))$. Integrating Eq. (1) from $h_i(t)$ to t , $i = 1, 2, \dots, n$, we get

$$x(t) - x(h_i(t)) + \int_{h_i(t)}^t \sum_{j=1}^n a_j(s) x(g_j(s)) ds = 0 \quad \text{for } t \in [h_1^{-1}(w_n^{-1}(T_0)), T_1]. \tag{16}$$

It follows from Eq. (1) and $h_j(t) \geq g_j(s)$, $h_i(t) \leq s \leq t$, $j = 1, 2, \dots, n$, that

$$x(g_j(s)) = x(h_j(t)) e^{\int_{g_j(s)}^{h_j(t)} \sum_{j_1=1}^n a_{j_1}(s_1) \frac{x(g_{j_1}(s_1))}{x(s_1)} ds_1}.$$

Substituting into Eq. (16), we get

$$x(t) - x(h_i(t)) + \sum_{j=1}^n x(h_j(t)) \int_{h_i(t)}^t a_j(s) e^{\int_{g_j(s)}^{h_j(t)} \sum_{j_1=1}^n a_{j_1}(s_1) \frac{x(g_{j_1}(s_1))}{x(s_1)} ds_1} ds = 0$$

for $t \in [h_1^{-1}(w_n^{-2}(T_0)), T_1]$. (17)

By using (5), we have

$$\frac{x(h_j(s_1))}{x(s_1)} \geq \frac{x(g_j(s_1))}{x(s_1)} \geq R^k(\eta_j), \quad g_j(s) \leq s_1 \leq h_j(t), h_i(t) \leq s \leq t, i, j = 1, 2, \dots, n,$$

for $t \in [h_j^{-1}(w_j^{-1}(h_j^{-1}(w_j^{-k}(w_n^{-1}(T_0))))), T_1] \subseteq [h_j^{-2}(w_j^{-(k+1)}(w_n^{-1}(T_0))), T_1]$. This together with (17) leads to

$$x(t) - x(h_i(t)) + \sum_{j=1}^n x(h_j(t)) \int_{h_i(t)}^t a_j(s) e^{\int_{g_j(s)}^{h_j(t)} \sum_{j_1=1}^n a_{j_1}(s_1) R^k(\eta_{j_1}) ds_1} ds \leq 0$$

for $t \in [h_1^{-2}(w_n^{-(k+2)}(T_0)), T_1]$.

That is,

$$x(h_i(t)) > \sum_{j=1}^n x(h_j(t)) \int_{h_i(t)}^t a_j(s) e^{\int_{g_j(s)}^{h_j(t)} \sum_{j_1=1}^n a_{j_1}(s_1) R^k(\eta_{j_1}) ds_1} ds$$

for $t \in [h_1^{-2}(w_n^{-(k+2)}(T_0)), T_1]$.

By using the arithmetic–geometric mean, we obtain

$$x(h_i(t)) > n \left(\prod_{j=1}^n x(h_j(t)) \right)^{\frac{1}{n}} \left(\prod_{j=1}^n \int_{h_i(t)}^t a_j(s) e^{\int_{g_j(s)}^{h_j(t)} \sum_{j_1=1}^n a_{j_1}(s_1) R^k(\eta_{j_1}) ds_1} ds \right)^{\frac{1}{n}}$$

for $t \in [h_1^{-2}(w_n^{-(k+2)}(T_0)), T_1]$. Taking the product of both sides

$$\prod_{j=1}^n x(h_j(t)) > n^n \left(\prod_{j=1}^n x(h_j(t)) \right) \prod_{i=1}^n \left(\prod_{j=1}^n \int_{h_i(t)}^t a_j(s) e^{\int_{g_j(s)}^{h_j(t)} \sum_{j_1=1}^n a_{j_1}(s_1) R^k(\eta_{j_1}) ds_1} ds \right)^{\frac{1}{n}}$$

for $t \in [h_1^{-2}(w_n^{-(k+2)}(T_0)), T_1]$. Therefore,

$$\prod_{i=1}^n \left(\prod_{j=1}^n \int_{h_i(t)}^t a_j(s) e^{\int_{g_j(s)}^{h_j(t)} \sum_{j_1=1}^n a_{j_1}(s_1) R^k(\eta_{j_1}) ds_1} ds \right)^{\frac{1}{n}} < \frac{1}{n^n} \quad \text{for } t \in [h_1^{-2}(w_n^{-(k+2)}(T_0)), T_1],$$

which contradicts (15). The proof is complete. □

Remark 2.1

- (i) It should be noted that $w_n^{-1}(t) - t < \infty$ when $\sup_{t \geq t_1} \{t - g_j(t)\} < \infty$ for $j = 1, 2, \dots, n$. Therefore, all upper bounds of the distance between zeros of all solutions of Eq. (1) obtained in this work are bounded. For example,

$$\begin{aligned} h_r^{-1}(w_r^{-k}(w_n^{-1}(t))) - t &\leq w_n^{-(k+2)}(t) - t \\ &= w_n^{-1}(w_n^{-(k+1)}(t)) - w_n^{-(k+1)}(t) + w_n^{-(k+1)}(t) - \dots + w_n^{-1}(t) - t \\ &< \infty. \end{aligned}$$

(ii) Since

$$R^k(d) \geq f_k(d), \quad k = 0, 1, \dots,$$

for some values of d , where

$$\int_{h_n(t)}^t \sum_{j=1}^n a_j(s) ds \geq d \quad \text{for } t \geq h_n^{-1}(t_1),$$

and the sequence $\{R^k(d)\}_{k \geq 1}$ is defined by (4), and

$$f_0(d) = 1, \quad f_1(d) = \frac{1}{1-d}, \quad f_k(d) = \frac{f_{k-2}(d)}{f_{k-2}(d) + 1 - e^{df_{k-2}(d)}}, \quad k = 2, 3, \dots$$

Then, by using a similar argument as in the proof of Lemma 2.1, we can improve [11, Lemma 2.4] and consequently all results that use it, as [11, Theorem 2.23].

3 Numerical examples

This section is devoted to validating the main theoretical findings through several examples. We first begin with the following example:

Example 3.1 Consider the differential equation with multiple delays

$$x'(t) + a_1(t)x(g_1(t)) + a_2(t)x(g_2(t)) = 0, \quad t \geq 3, \tag{18}$$

where $a_1(t) = \mu, a_2(t) = \rho, \mu, \rho > 0$,

$$g_1(t) = \begin{cases} t - 2 & \text{if } t \in [3i, 3i + 1], \\ \frac{1}{4}(5t - 3i - 9) & \text{if } t \in [3i + 1, 3i + 2], \\ \frac{1}{4}(3t + 3i - 5) & \text{if } t \in [3i + 2, 3i + 3], \end{cases} \quad i \in \mathbb{N},$$

$g_2(t) = t - \frac{1}{4}$. Clearly,

$$t - 2 \leq g_1(t) \leq t - \frac{7}{4}.$$

Since $h_1(t) = g_1(t)$ and $h_2(t) = g_2(t)$, so $w_1(t) = g_1(t)$ and $w_2(t) = \min_{1 \leq j \leq 2} g_j(t) = g_1(t)$. It follows that

$$\max_{1 \leq j \leq 2} w_j^{-i}(t) = w_2^{-i}(t) \leq t + 2i.$$

Let

$$I(t) = \prod_{i=1}^2 \left(\prod_{j=1}^2 \int_{h_i(t)}^t a_j(s) e^{\int_{g_j(s)}^{h_j(t)} \sum_{\nu=1}^2 R^k(\eta_{j\nu}) a_{j\nu}(s_1) ds_1} ds \right)^{\frac{1}{2}}.$$

Then

$$I(t) \geq \left(\mu\rho \int_{h_1(t)}^t e^{(\mu+\rho)(h_1(t)-g_1(s))} ds \times \int_{h_1(t)}^t e^{(\mu+\rho)(h_2(t)-g_2(s))} ds \right)^{\frac{1}{2}} \\ \times \left(\mu\rho \int_{h_2(t)}^t e^{(\mu+\rho)(h_1(t)-g_1(s))} ds \times \int_{h_2(t)}^t e^{(\mu+\rho)(h_2(t)-g_2(s))} ds \right)^{\frac{1}{2}}.$$

Therefore,

$$I(t) \geq \left(\mu\rho \int_{t-\frac{175}{100}}^t e^{(\mu+\rho)(t-s-\frac{1}{4})} ds \times \int_{t-\frac{175}{100}}^t e^{(\mu+\rho)(t-s)} ds \right)^{\frac{1}{2}} \\ \times \left(\mu\rho \int_{t-\frac{1}{4}}^t e^{(\mu+\rho)(t-s-\frac{1}{4})} ds \times \int_{t-\frac{1}{4}}^t e^{(\mu+\rho)(t-s)} ds \right)^{\frac{1}{2}} \\ = \frac{\mu\rho e^{-\frac{1}{4}(\mu+\rho)}(e^{\frac{7}{4}(\mu+\rho)} - 1)(e^{\frac{1}{4}(\mu+\rho)} - 1)}{(\mu + \rho)^2} > \frac{1}{4} \quad \text{for } \mu \geq \frac{1}{2}, \rho \geq \frac{56}{115}.$$

Consequently, Theorem 2.2 with $k = 0$ implies that $D_3(x) \leq \sup_{t \geq 3} \{w_2^{-4}(t) - t\} \leq 8$ for $\mu \geq \frac{1}{2}, \rho \geq \frac{56}{115}$.

Observe that none of the results in [11] apply to Eq. (18) when $0 < \mu + \rho \leq \frac{4}{e}$. The reason for this is that

$$\max_{1 \leq j \leq 2} g_j(t) = t - \frac{1}{4},$$

which leads to

$$\int_{\max_{1 \leq j \leq 2} g_j(t)}^t (a_1(s) + a_2(s)) ds \leq \frac{1}{4}(\mu + \rho) < \frac{1}{e} \quad \text{for } \mu + \rho < \frac{4}{e}.$$

Next, we move to the next example.

Example 3.2 Consider the differential equation

$$x'(t) + \frac{1}{2}x\left(t - \frac{11}{10}\right) + \frac{1}{2}x(t - 1) + x(t - \epsilon) = 0, \quad t \geq \frac{11}{10}, \tag{19}$$

where $0 < \epsilon < \frac{1}{2}$. This equation is of the form (1) with $a_1(t) = a_2(t) = \frac{1}{2}, a_3(t) = 1, g_1(t) = t - \frac{11}{10}, g_2(t) = t - 1,$ and $g_3(t) = t - \epsilon$. Clearly,

$$h_2(t) = \max_{1 \leq j \leq 2} g_j(t) = t - 1, \quad w_2(t) = \min_{1 \leq j \leq 2} g_j(t) = t - \frac{11}{10}, \quad w_3(t) = \min_{1 \leq j \leq 3} g_j(t) = t - \frac{11}{10}$$

and

$$h_2^{-k}(t) = t + k, \quad w_2^{-k}(t) = \max_{1 \leq j \leq 2} w_j^{-k}(t) = t + \frac{11}{10}k, \quad w_3^{-k}(t) = \max_{1 \leq j \leq 3} w_j^{-k}(t) = t + \frac{11}{10}k.$$

Since

$$\sum_{i=1}^1 \prod_{j=2}^i R^{k+1-j}(\eta_r) \int_{h_2(t)}^t \sum_{j=1}^2 B_{j,2}^i(s, h_2(t)) ds = \int_{h_2(t)}^t \sum_{j=1}^2 a_j(s) ds = 1.$$

Then, according to Theorem 2.1 with $k = 0$, Eq. (19) is oscillatory and $D_{\frac{11}{10}}(x) \leq \sup_{t \geq \frac{11}{10}} \{h_2^{-1}(w_2^{-1}(w_3^{-1}(t))) - t\} = \frac{16}{5}$.

Observe, however, that

$$\max_{1 \leq j \leq 3} g_j(t) = t - \epsilon.$$

It is not difficult to show that all results of [11], [3, Theorem 3] and [3, Theorem 4] fail to apply to Eq. (19) for sufficiently small ϵ . Also, observe that

$$\int_{g_j(t)}^t a_j(s) ds < 1 \quad \text{for } j = 1, 2, 3.$$

Therefore, [3, Theorem 2] cannot give an approximation to $D_{\frac{11}{10}}(x)$ for sufficiently small ϵ better than $\frac{16}{5}$.

4 Conclusion

In this paper, we studied the distribution of zeros of first-order delay differential equations. Also, we obtained upper bounds for the zeros of a first-order differential equation with several delays. Finally, some examples are demonstrated to prove the theoretical results.

Funding

This study is supported via funding from Prince Sattam bin Abdulaziz University project number (PSAU/2023/R/1444). The researchers would like to thank the Deanship of Scientific Research of Qassim University for funding the publication of this project.

Availability of data and materials

Not applicable.

Declarations

Competing interests

The authors declare no competing interests.

Author contributions

Supervision, E.A. and B.E.-M.; writing—original draft, E.A.; writing—review editing, B.E.-M. All authors have read and agreed to the published version of the manuscript.

Author details

¹Department of Mathematics, College of Sciences and Humanities, Prince Sattam Bin Abdulaziz University, Alkharj 11942, Saudi Arabia. ²Department of Mathematics, College of Science and Arts, Al-Badaya, Qassim University, Buraidah 51951, Saudi Arabia. ³Department of Mathematics, Faculty of Science, Damietta University, New Damietta 34517, Egypt.

Received: 13 April 2023 Accepted: 24 July 2023 Published online: 16 August 2023

References

1. Agarwal, R.P., Berezansky, L., Braverman, E., Domoshnitsky, A.: Non-oscillation Theory of Functional Differential Equations with Applications. Springer, New York (2012)
2. Agarwal, R.P., Grace, S.R., ÓRegan, D.: Oscillation Theory for Difference and Functional Differential Equations. Springer, Berlin (2013)
3. Attia, E.R., Chatzarakis, G.E.: Upper bounds for the distance between adjacent zeros of first-order linear differential equations with several delays. *Mathematics* **10**, 648 (2022)

4. Baker, F.A., El-Morshedy, H.A.: The distribution of zeros of all solutions of first order neutral differential equations. *Appl. Math. Comput.* **259**, 777–789 (2015)
5. Baker, F.A., El-Morshedy, H.A.: On the distribution of zeros of solutions of a first order neutral differential equation. *Sci. J. Damietta Fac. Sci.* **4**, 1–9 (2015)
6. Barr, T.H.: Oscillations in linear delay differential equations. *J. Math. Anal. Appl.* **195**, 261–277 (1995)
7. Berezansky, L., Domshlak, Y.: Can a solution of a linear delay differential equation have an infinite number of isolated zeros on a finite interval? *Appl. Math. Lett.* **19**, 587–589 (2006)
8. Domoshnitsky, A., Drakhlin, M., Stavroulakis, I.P.: Distribution of zeros of solutions to functional equations. *Math. Comput. Model.* **42**, 193–205 (2005)
9. Domshlak, Y., Aliev, A.I.: On oscillatory properties of the first order differential equations with one or two retarded arguments. *Hiroshima Math. J.* **18**, 31–46 (1988)
10. El-Morshedy, H.A.: On the distribution of zeros of solutions of first order delay differential equations. *Nonlinear Anal.* **74**, 3353–3362 (2011)
11. El-Morshedy, H.A., Attia, E.R.: On the distance between adjacent zeros of solutions of first order differential equations with distributed delays. *Electron. J. Qual. Theory Differ. Equ.* **2016**, 8 (2016)
12. Elbert, A., Stavroulakis, I.P.: Oscillation of first order differential equations with deviating arguments. In: *Recent Trends in Differential Equations*. World Science Series in Applied Analysis, vol. 1, pp. 163–178. World Scientific, River Edge (1992)
13. Erbe, L.H., Kong, Q., Zhang, B.G.: *Oscillation Theory for Functional Differential Equations*. Dekker, New York (1995)
14. Gopalsamy, K.: *Stability and Oscillation in Delay Differential Equations of Population Dynamics*. Kluwer Academic, Dordrecht (1992)
15. Györi, I., Ladas, G.: *Oscillation Theory of Delay Differential Equations with Applications*. Clarendon, Oxford (1991)
16. Jaroš, J., Stavroulakis, I.P.: Oscillation tests for delay equations. *Rocky Mt. J. Math.* **29**, 197–207 (1999)
17. McCalla, C.: Zeros of the solutions of first order functional differential equations. *SIAM J. Math. Anal.* **9**, 843–847 (1978)
18. Pituk, M., Stavroulakis, J.I.: The first positive root of the fundamental solution is an optimal oscillation bound for linear delay differential equations. *J. Math. Anal.* **507**, 125789 (2022)
19. Sficas, Y.G., Stavroulakis, I.P.: Oscillation criteria for first-order delay equations. *Bull. Lond. Math. Soc.* **35**, 239–246 (2003)
20. Tang, X., Yu, J.: Distribution of zeros of solutions of first order delay differential equations. *Appl. Math. J. Chin. Univ. Ser. B* **14**, 375–380 (1999)
21. Wu, H.W., Cheng, S.S.: Upper bounds for the distances between adjacent zeros of solutions of delay differential equations. *Appl. Math. Comput.* **218**, 3379–3386 (2011)
22. Wu, H.W., Cheng, S.S., Wang, Q.R.: Distribution of zeros of solutions of functional differential equations. *Appl. Math. Comput.* **193**, 154–161 (2007)
23. Wu, H.W., Erbe, L.: On the distance between consecutive zeros of solutions of first order delay differential equations. *Appl. Math. Comput.* **219**, 8622–8631 (2013)
24. Wu, H.W., Erbe, L., Peterson, A.: Upper bounds for the distances between consecutive zeros of solutions of first order delay differential equations. *Appl. Math. Comput.* **429**, 562–575 (2015)
25. Wu, H.W., Xu, Y.T.: The distribution of zeros of solutions of neutral differential equations. *Appl. Math. Comput.* **156**, 665–677 (2004)
26. Zhang, B.G., Zhou, Y.: The distribution of zeros of solutions of differential equations with a variable delay. *J. Math. Anal. Appl.* **256**, 216–228 (2001)
27. Zhou, Y.: The distribution of zeros of solutions of first order functional differential equations. *Bull. Aust. Math. Soc.* **59**, 305–314 (1999)

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)
