(2023) 2023:161

RESEARCH

Open Access

C

New refinements of the Cauchy–Bunyakovsky inequality



*Correspondence: saeed.montazeri@outlook.com ¹Faculty of Mechanical Engineering, Shahid Rajaee Teacher Training University, Tehran, Iran

Abstract

This paper presents new refinements on the integral form of Cauchy–Schwartz inequality known as Cauchy–Bunyakovsky inequality. It is proved that when we possess a weighted sum of a set of Cauchy–Bunyakovsky inequalities, there are two forms of refinements enhancing the precision of the original inequality. The superiority of one refinement over the other depends on the problem in which the presented theorem is utilized.

Keywords: Cauchy–Bunyakovsky; Cauchy–Schwartz; Inequality; Refinement

1 Introduction

The most common applications of inequalities in science and engineering occur when there is incomplete information about a system, yet we can utilize an inequality to estimate a certain quantity. It is obvious that the closer the two sides of an inequality are to each other, the more accurate our estimation becomes. This is why refining inequalities is advantageous. Up to the present date, numerous generalizations and refinements for the Cauchy–Schwartz and Cauchy–Bunyakovsky inequalities have been extensively studied [1-10].

The integral form of Cauchy–Schwartz inequality known as Cauchy–Bunyakovsky inequality for two real functions f(x) and g(x) states that [11, Ch. 1]

$$\left(\int_{a}^{b} f(x)g(x)\,dx\right)^{2} \leq \left(\int_{a}^{b} f^{2}(x)\,dx\right)\left(\int_{a}^{b} g^{2}(x)\,dx\right). \tag{1}$$

Accordingly, it is obvious that for a set of *n* functions $\{f_i(x)\}_{i=1}^n$, we have:

$$\sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b f_i(x) g(x) \, dx \right)^2 \le \sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b f_i^2(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right), \tag{2}$$

© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.



where $\{p_i\}_{i=1}^n > 0$ and $P = \sum_{i=1}^n p_i$. In this paper, it is proved that Inequality (2) can be refined into the following two forms:

$$\sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b f_i(x) g(x) \, dx \right)^2 + \begin{cases} q_1^2 \\ q_2^2 \end{cases} \le \sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b f_i^2(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right), \tag{3}$$

where q_1^2 and q_2^2 are functions of $f_i(x)$, p_i and g(x).

2 Main results

The main result of this paper is contained in the following theorem.

Theorem Consider a set of *n* Cauchy–Bunyakovsky inequalities applied to functions $\{f_i(x)\}_{i=1}^n \in R \text{ and } g(x) \in R \text{ as follows:}$

$$\left(\int_{a}^{b} f_{i}(x)g(x)\,dx\right)^{2} \leq \left(\int_{a}^{b} f_{i}^{2}(x)\,dx\right)\left(\int_{a}^{b} g^{2}(x)\,dx\right), \quad i=1,\ldots,n.$$

$$\tag{4}$$

Then, there are two refinements for the weighted sum of n Cauchy–Bunyakovsky inequalities, as follows:

$$\sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b f_i(x) g(x) \, dx \right)^2 + \begin{cases} q_1^2 \\ q_2^2 \end{cases} \le \sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b f_i^2(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right), \tag{5}$$

where $\{p_i\}_{i=1}^n > 0$, $P = \sum_{i=1}^n p_i$ and

$$q_1^2 = \int_a^b F^2(x) \, dx \int_a^b g^2(x) \, dx - \left(\int_a^b F(x)g(x) \, dx\right)^2,\tag{6}$$

$$q_2^2 = \sum_{i=1}^n \frac{p_i}{P} \left(\int_a^b r_i^2(x) \, dx \int_a^b g^2(x) \, dx - \left(\int_a^b r_i(x) g(x) \, dx \right)^2 \right),\tag{7}$$

$$F(x) = \frac{\sum_{i=1}^{n} f_i(x) p_i}{P},$$
(8)

$$r_i(x) = f_i(x) - F(x).$$
 (9)

Proof The theorem is proved through a three-step process. In the first and second steps, the right and left sides of Inequality (2) are obtained, respectively. In the third step, the obtained values are subtracted from each other.

Step 1: Obtaining the right side of the Inequality.

Let S_1 be equal to the right side of Inequality (2) as follows:

$$S_1 = \sum_{i=1}^n \frac{p_i}{P} \left(\int_a^b f_i^2(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right). \tag{10}$$

Substituting $f_i(x)$ from Equation (9) into Equation (10) gives:

$$S_{1} = \sum_{i=1}^{n} \frac{p_{i}}{P} \left(\int_{a}^{b} F^{2}(x) dx \right) \left(\int_{a}^{b} g^{2}(x) dx \right) + 2 \sum_{i=1}^{n} \frac{p_{i}}{P} \left(\int_{a}^{b} r_{i}(x) F(x) dx \right) \left(\int_{a}^{b} g^{2}(x) dx \right) + \sum_{i=1}^{n} \frac{p_{i}}{P} \left(\int_{a}^{b} r_{i}^{2}(x) dx \right) \left(\int_{a}^{b} g^{2}(x) dx \right).$$
(11)

The first summation in Equation (11) can be simplified as:

$$\sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b F^2(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right) = \left(\int_a^b F^2(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right). \tag{12}$$

The second summation in Equation (11) is obtained as follows:

$$2\sum_{i=1}^{n} \frac{p_{i}}{p} \left(\int_{a}^{b} r_{i}(x)F(x) dx \right) \left(\int_{a}^{b} g^{2}(x) dx \right)$$

= $2 \left(\int_{a}^{b} g^{2}(x) dx \right) \sum_{i=1}^{n} \frac{p_{i}}{p} \int_{a}^{b} F(x)r_{i}(x) dx$
= $2 \left(\int_{a}^{b} g^{2}(x) dx \right) \int_{a}^{b} \sum_{i=1}^{n} \frac{p_{i}}{p} F(x)r_{i}(x) dx$
= $2 \left(\int_{a}^{b} g^{2}(x) dx \right) \int_{a}^{b} \frac{F(x)}{p} \sum_{i=1}^{n} p_{i}r_{i}(x) dx.$ (13)

From Equations (8) and (9), it is obvious that $\sum_{i=1}^{n} p_i r_i(x) = 0$. Hence,

$$2\sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b r_i(x) F(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right) = 0.$$
(14)

Therefore, the value S_1 in Equation (11) can be rewritten as follows:

$$S_{1} = \left(\int_{a}^{b} F^{2}(x) \, dx\right) \left(\int_{a}^{b} g^{2}(x) \, dx\right) + \sum_{i=1}^{n} \frac{p_{i}}{P} \left(\int_{a}^{b} r_{i}^{2}(x) \, dx\right) \left(\int_{a}^{b} g^{2}(x) \, dx\right).$$
(15)

Step 2: Obtaining the left side of the Inequality.

Let S_2 be equal to the left side of Inequality (2) as follows:

$$S_2 = \sum_{i=1}^n \frac{p_i}{P} \left(\int_a^b f_i(x) g(x) \, dx \right)^2.$$
(16)

Substituting $f_i(x)$ from Equation (9) into Equation (16) gives:

$$S_{2} = \sum_{i=1}^{n} \frac{p_{i}}{P} \left(\int_{a}^{b} F(x)g(x) \, dx + \int_{a}^{b} r_{i}(x)g(x) \, dx \right)^{2}$$

$$= \sum_{i=1}^{n} \frac{p_{i}}{P} \left(\int_{a}^{b} F(x)g(x) \, dx \right)^{2} + 2 \sum_{i=1}^{n} \frac{p_{i}}{P} \left(\int_{a}^{b} F(x)g(x) \, dx \right) \left(\int_{a}^{b} r_{i}(x)g(x) \, dx \right)$$

$$+ \sum_{i=1}^{n} \frac{p_{i}}{P} \left(\int_{a}^{b} r_{i}(x)g(x) \, dx \right)^{2}.$$
(17)

The first summation in Equation (17) can be simplified as:

$$\sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b F(x) g(x) \, dx \right)^2 = \left(\int_a^b F(x) g(x) \, dx \right)^2.$$
(18)

The second summation in Equation (17) can be obtained as follows:

$$2\sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b F(x)g(x) \, dx \right) \left(\int_a^b r_i(x)g(x) \, dx \right)$$

$$= 2 \left(\int_a^b F(x)g(x) \, dx \right) \sum_{i=1}^{n} \frac{p_i}{P} \left(\int_a^b r_i(x)g(x) \, dx \right)$$

$$= 2 \left(\int_a^b F(x)g(x) \, dx \right) \int_a^b \sum_{i=1}^{n} \frac{p_i}{P} r_i(x)g(x) \, dx$$

$$= 2 \left(\int_a^b F(x)g(x) \, dx \right) \int_a^b \frac{g(x)}{P} \sum_{i=1}^{n} p_i r_i(x) \, dx.$$
(19)

We know that $\sum_{i=1}^{n} p_i r_i(x) = 0$. Thus, the second summation in the right side of Equation (17) is equal to zero similar to step 1. Therefore, the value S_2 in Equation (17) can be rewritten as follows:

$$S_2 = \left(\int_a^b F(x)g(x)\,dx\right)^2 + \sum_{i=1}^n \frac{p_i}{P} \left(\int_a^b r_i(x)g(x)\,dx\right)^2.$$
 (20)

Step 3:

In this step, the value of $S_1 - S_2$ is obtained. From Equations (15) and (20), we can write:

$$S_{1} - S_{2} = \left(\left(\int_{a}^{b} F^{2}(x) \, dx \right) \left(\int_{a}^{b} g^{2}(x) \, dx \right) - \left(\int_{a}^{b} F(x)g(x) \, dx \right)^{2} \right) \\ + \sum_{i=1}^{n} \frac{p_{i}}{P} \left(\left(\int_{a}^{b} r_{i}^{2}(x) \, dx \right) \left(\int_{a}^{b} g^{2}(x) \, dx \right) - \left(\int_{a}^{b} r_{i}(x)g(x) \, dx \right)^{2} \right).$$
(21)

Equation (21) is the sum of two positive parts, according to the Cauchy–Bunyakovsky inequality. We can name them q_1^2 and q_2^2 as follows:

$$q_1^2 = \left(\left(\int_a^b F^2(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right) - \left(\int_a^b F(x) g(x) \, dx \right)^2 \right), \tag{22}$$

and

$$q_2^2 = \sum_{i=1}^n \frac{p_i}{P} \left(\left(\int_a^b r_i^2(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right) - \left(\int_a^b r_i(x) g(x) \, dx \right)^2 \right). \tag{23}$$

Given that values of S_1 and S_2 are respectively right and left side of Inequality (2), the inequality can be refined into two following forms:

$$S_1 - S_2 = \sum_{i=1}^n \frac{p_i}{P} \left(\int_a^b f_i^2(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right) - \sum_{i=1}^n \frac{p_i}{P} \left(\int_a^b f_i(x) g(x) \, dx \right)^2 \ge q_1^2, \quad (24)$$

or

$$S_1 - S_2 = \sum_{i=1}^n \frac{p_i}{P} \left(\int_a^b f_i^2(x) \, dx \right) \left(\int_a^b g^2(x) \, dx \right) - \sum_{i=1}^n \frac{p_i}{P} \left(\int_a^b f_i(x) g(x) \, dx \right)^2 \ge q_2^2.$$
(25)

The proof is complete.

3 Conclusion

It is obvious that a weighted sum of a set of Cauchy–Bunyakovsky inequalities results in a new Cauchy–Bunyakovsky inequality. The theorem presented in this paper demonstrates that there exist two distinct refinement forms for the inequality obtained from a weighted sum of *n* Cauchy–Bunyakovsky inequalities.

Funding

There is no funding to declare.

Data availability

There is no data to declare.

Declarations

Competing interests

The authors declare no competing interests.

Author contributions

This article has only one author.

Received: 9 October 2023 Accepted: 19 December 2023 Published online: 22 December 2023

References

- 1. Han, J., Shi, J.: Refinements of Cauchy–Schwarz norm inequality. J. Math. Inequal. 13, 1095–1103 (2019)
- 2. Alzer, H.: A refinement of the Cauchy–Schwarz inequality. J. Math. Anal. Appl. 168(2), 596–604 (1992)
- 3. Masjed-Jamei, M., Hussain, N.: More results on a functional generalization of the Cauchy–Schwarz inequality. J. Inequal. Appl. 2012, 239 (2012)
- Masjed-Jamei, M.: A functional generalization of the Cauchy–Schwarz inequality and some subclasses. Appl. Math. Lett. 22(9), 1335–1339 (2009)
- 5. Mercer, P.R.: A refined Cauchy–Schwarz inequality. Int. J. Math. Educ. Sci. Technol. 38(6), 839–843 (2007)
- 6. Wada, S.: On some refinement of the Cauchy–Schwarz inequality. Linear Algebra Appl. 420(2–3), 433–440 (2007)
- 7. Mercer, P.R.: A refined Cauchy–Schwarz inequality. Int. J. Math. Educ. Sci. Technol. 38(6), 839–843 (2007)
- 8. Choi, D.: A generalization of the Cauchy–Schwarz inequality. J. Math. Inequal. 10(4), 1009–1012 (2016)
- 9. Tuo, L.: Generalizations of Cauchy-Schwarz inequality in unitary spaces. J. Inequal. Appl. 2015(1), 201 (2015)
- 10. Sever Dragomir, S.: Improving Schwarz inequality in inner product spaces. arXiv e-prints, p. arXiv-1709 (2017)
- 11. Steele, J.M.: The Cauchy–Schwarz Master Class: An Introduction to the Art of Mathematical Inequalities. Cambridge University Press, Cambridge (2004)

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.