

Research Article

Hermite-Hadamard-Type Inequalities for Increasing Positively Homogeneous Functions

G. R. Adilov and S. Kemali

Received 20 October 2006; Accepted 6 June 2007

Recommended by Kok Lay Teo

We study Hermite-Hadamard-type inequalities for increasing positively homogeneous functions. Some examples of such inequalities for functions defined on special domains are given.

Copyright © 2007 G. R. Adilov and S. Kemali. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Recently, Hermite-Hadamard-type inequalities and their applications have attracted considerable interest, as shown in the book [1], for example. These inequalities have been studied for various classes of functions such as convex functions [1], quasiconvex functions [2–4], p -functions [3, 5], Godnova-Levin type functions [5], r -convex functions [6], increasing convex-along-rays functions [7], and increasing radiant functions [8], and it is shown that these inequalities are sharp.

For instance, if $f : [0, 1] \rightarrow \mathbb{R}$ is an arbitrary nonnegative quasiconvex function, then for any $u \in (0, 1)$ one has (see [3])

$$f(u) \leq \frac{1}{\min(u, 1-u)} \int_0^1 f(x) dx, \quad (1.1)$$

and the inequality (1.1) is sharp.

In this paper, we consider one generalization of Hermite-Hadamard-type inequalities for the class of increasing positively homogeneous of degree one functions defined on $\mathbb{R}_{++}^n = \{x \in \mathbb{R}^n : x_i > 0, i = 1, 2, 3, \dots, n\}$.

The structure of the paper is as follows: in Section 2, certain concepts of abstract convexity, definition of increasing positively homogeneous of degree one functions and its important properties are given. In Section 3, Hermite-Hadamard-type inequalities for

the class of increasing positively homogeneous of degree one functions are considered. Some examples of such inequalities for functions defined on \mathbb{R}_{++}^2 are given in Section 4.

2. Preliminaries

First we recall some definitions from abstract convexity. Let \mathbb{R} be a real line and $\mathbb{R}_{+\infty} = \mathbb{R} \cup \{+\infty\}$. Consider a set X and a set H of function $h : X \rightarrow \mathbb{R}$ defined on X . A function $f : X \rightarrow \mathbb{R}_{+\infty}$ is called abstract convex with respect to H (or H -convex) if there exists a set $U \subset H$ such that

$$f(x) = \sup \{h(x) : h \in U\} \quad \forall x \in X. \quad (2.1)$$

Clearly, f is H -convex if and only if

$$f(x) = \sup \{h(x) : h \leq f\} \quad \forall x \in X. \quad (2.2)$$

Let Y be a set of functions $f : X \rightarrow \mathbb{R}_{+\infty}$. A set $H \subset Y$ is called a supremal generator of the set Y , if each function $f \in Y$ is abstract convex with respect to H .

In some cases, the investigation of Hermite-Hadamard-type inequalities is based on the principle of preservation of inequalities [9].

PROPOSITION 2.1 (principle of preservation of inequalities). *Let H be a supremal generator of Y and let Ψ be an increasing functional defined on Y . Then*

$$(h(u) \leq \Psi(h) \quad \forall h \in H) \iff (f(u) \leq \Psi(f) \quad \forall f \in Y). \quad (2.3)$$

A function f defined on \mathbb{R}_{++}^n is called increasing (with respect to the coordinate-wise order relation) if $x \geq y$ implies $f(x) \geq f(y)$.

The function f is positively homogeneous of degree one if $f(\lambda x) = \lambda f(x)$ for all $x \in \mathbb{R}_{++}^n$ and $\lambda > 0$.

Let L be the set of all min-type functions defined on \mathbb{R}_{++}^n , that is, the set L consists of identical zero and all the functions of the form

$$l(x) = \langle l, x \rangle = \min_i \frac{x_i}{l_i}, \quad x \in \mathbb{R}_{++}^n \quad (2.4)$$

with all $l \in \mathbb{R}_{++}^n$.

One has (see [9]) that a function $f : \mathbb{R}_{++}^n \rightarrow \mathbb{R}$ is L -convex if and only if f is increasing and positively homogeneous of degree one (shortly IPH).

Let us present the important property of IPH functions.

PROPOSITION 2.2. *Let f be an IPH function defined on \mathbb{R}_{++}^n . Then the following inequality holds for all $x, l \in \mathbb{R}_{++}^n$:*

$$f(l)\langle l, x \rangle \leq f(x). \quad (2.5)$$

Proof. Since $\langle l, x \rangle = \min_{1 \leq i \leq n} (x_i/l_i)$, then $\langle l, x \rangle l_i \leq x_i$ is proved for all $i = 1, 2, 3, \dots, n$.
Consequently, we get $\langle l, x \rangle l \leq x$. Because f is an IPH function,

$$f(x) \geq f(\langle l, x \rangle l) = \langle l, x \rangle f(l) \quad \forall l, x \in \mathbb{R}_{++}^n. \quad (2.6)$$

□

Let f be an IPH function defined on \mathbb{R}_{++}^n and $D \subset \mathbb{R}_{++}^n$. It can be easily shown by Proposition 2.2 that the function

$$f_D(x) = \sup_{l \in D} (f(l)\langle l, x \rangle) \quad (2.7)$$

is IPH and it possesses the properties

$$f_D(x) \leq f(x) \quad \forall x \in \mathbb{R}_{++}^n, \quad f_D(x) = f(x) \quad \forall x \in D. \quad (2.8)$$

Let $D \subset \mathbb{R}_{++}^n$. A function $f : D \rightarrow [0, \infty]$ is called IPH on D if there exists an IPH function F defined on \mathbb{R}_{++}^n such that $F|_D = f$, that is, $F(x) = f(x)$ for all $x \in D$.

PROPOSITION 2.3. *Let $f : D \rightarrow [0, \infty]$ be a function on $D \subset \mathbb{R}_{++}^n$, then the following assertions are equivalent:*

- (i) f is abstract convex with respect to the set of functions $c\langle l, \cdot \rangle : D \rightarrow [0, \infty)$ with $l \in D, c \geq 0$;
- (ii) f is IPH function on D ;
- (iii) $f(l)\langle l, x \rangle \leq f(x)$ for all $l, x \in D$.

Proof. (i) \Rightarrow (ii) It is obvious since any function $l(x) = c\langle l, x \rangle$ defined on D can be considered as elementary function $l(x) \in L$ defined on \mathbb{R}_{++}^n .

(ii) \Rightarrow (iii) By definition, there exists an IPH function $F : \mathbb{R}_{++}^n \rightarrow [0, \infty]$ such that $F(x) = f(x)$ for all $x \in D$. Then by (2.7) we have

$$f(x) = F_D(x) = \sup_{l \in D} (F(l)\langle l, x \rangle) = \sup_{l \in D} (f(l)\langle l, x \rangle) \quad (2.9)$$

for all $x \in D$, which implies the assertion (iii).

(iii) \Rightarrow (i) Consider the function f_D defined on D , $\sup_{l \in D} (f(l)\langle l, x \rangle) = f_D(x)$. It is clear that f_D is abstract convex with respect to the set of functions $\{c\langle l, \cdot \rangle : l \in D, c \geq 0\}$ defined on D . Further, using (iii) we get that for all $x \in D$,

$$f_D(x) \leq f(x) = f(x)\langle x, x \rangle \leq \sup_{l \in D} (f(l)\langle l, x \rangle) = f_D(x). \quad (2.10)$$

So, $f_D(x) = f(x)$ for all $x \in D$ and we have the defined statement (i). □

3. Hermite-Hadamard-type inequalities for IPH functions

Now, we will research to Hermite-Hadamard-type inequality for IPH functions.

PROPOSITION 3.1. *Let $D \subset \mathbb{R}_{++}^n$, $f : D \rightarrow [0, \infty]$ is IPH function, and f is integrable on D . Then*

$$f(u) \int_D \langle u, x \rangle dx \leq \int_D f(x) dx \tag{3.1}$$

for all $u \in D$.

Proof. It can be seen via Proposition 2.3. Since $f(l)\langle l, x \rangle \leq f(x)$ for all $l, x \in D$, (3.1) is clear. □

Let us investigate Hermite-Hadamard-type inequalities via $Q(D)$ sets given in [7, 8].

Let $D \subset \mathbb{R}_{++}^n$ be a closed domain, that is, D is bounded set such that $cl \text{int} D = D$. Denote by $Q(D)$ the set of all points $x^* \in D$ such that

$$\frac{1}{A(D)} \int_D \langle x^*, x \rangle dx = 1, \tag{3.2}$$

where $A(D) = \int_D dx$.

PROPOSITION 3.2. *Let f be an IPH function defined on D . If the set $Q(D)$ is nonempty and f is integrable on D , then*

$$\sup_{x^* \in Q(D)} f(x^*) \leq \frac{1}{A(D)} \int_D f(x) dx. \tag{3.3}$$

Proof. If we take $f(x^*) = +\infty$, by using the equality (2.5), it can be easily shown that f cannot be integrable. So $f(x^*) < +\infty$. According to Proposition 2.3,

$$f(x^*) \langle x^*, x \rangle \leq f(x) \quad \forall x \in D. \tag{3.4}$$

Since $x^* \in Q(D)$, then by (3.2) we get

$$\begin{aligned} f(x^*) &= f(x^*) \frac{1}{A(D)} \int_D \langle x^*, x \rangle dx \\ &= \frac{1}{A(D)} \int_D \langle x^*, x \rangle f(x^*) dx \leq \frac{1}{A(D)} \int_D f(x) dx. \end{aligned} \tag{3.5}$$

□

Remark 3.3. For each $x^* \in Q(D)$ we have also the following inequality, which is weaker than (3.3):

$$f(x^*) \leq \frac{1}{A(D)} \int_D f(x) dx. \tag{3.6}$$

However, even the inequality (3.6) is sharp. For example, if $f(x) = \langle x^*, x \rangle$, then (3.6) holds as the equality.

Remark 3.4. Let $Q(D)$ be a nonempty set. We can define a set $Q_k(D)$ for every positive real number k such that $Q_k(D) = \{u \in D : u = k \cdot x^*, x^* \in Q(D)\}$. The set $Q_k(D)$ above can be easily defined as follows: $Q_k(D) = \{u \in D : (k/A(D)) \int_D \langle u, x \rangle dx = 1\}$.

Considering the property that an IPH function is positively homogeneous of degree one, we can generalize the inequality (3.3) as follows:

$$\sup_{u \in Q_k(D)} f(u) \leq \frac{k}{A(D)} \int_D f(x) dx. \quad (3.7)$$

Let us try to derive inequalities similar to the right hand of the statement which is derived for convex functions (see [1]).

Let f be an IPH function defined on a closed domain $D \subset \mathbb{R}_{++}^n$, and f is integrable on D . Then $f(l)\langle l, x \rangle \leq f(x)$ for all $l, x \in D$. Hence for all $l, x \in D$,

$$f(l) \leq \frac{f(x)}{\langle l, x \rangle} = \langle x, l \rangle^+ f(x), \quad (3.8)$$

where $\langle x, l \rangle^+ = \max_{1 \leq i \leq n} l_i/x_i$ is the so-called max-type function.

We have established the following result.

PROPOSITION 3.5. *Let f be IPH and integrable function on D . Then*

$$\int_D f(x) dx \leq \inf_{u \in D} \left[f(u) \int_D \langle u, x \rangle^+ dx \right]. \quad (3.9)$$

For every $u \in D$, inequality

$$\int_D f(x) dx \leq f(u) \int_D \langle u, x \rangle^+ dx \quad (3.10)$$

is sharp.

4. Examples

On some special domains D of the cones \mathbb{R}_{++} and \mathbb{R}_{++}^2 , Hermite-Hadamard-type inequalities have been stated for ICAR and InR functions (see [7, 8]). Let us derive the set $Q(D)$ and the inequalities (3.1), (3.6), (3.9), for IPH functions, too.

Before the examples, for a region $D \subset \mathbb{R}_{++}^2$ and every $u \in D$, let us derive the computation formula of the integral $\int_D \langle u, x \rangle dx$.

Let $D \subset \mathbb{R}_{++}^2$ and $u = (u_1, u_2) \in D$. In order to calculate the integral, we represent the set D as $D_1(u) \cup D_2(u)$, where

$$D_1(u) = \left\{ x \in D : \frac{x_2}{u_2} \leq \frac{x_1}{u_1} \right\}, \quad D_2(u) = \left\{ x \in D : \frac{x_2}{u_2} \geq \frac{x_1}{u_1} \right\}. \quad (4.1)$$

Then

$$\begin{aligned} \int_D \langle u, x \rangle dx &= \int_{D_1(u)} \langle u, x \rangle dx + \int_{D_2(u)} \langle u, x \rangle dx \\ &= \frac{1}{u_2} \int_{D_1(u)} x_2 dx_1 dx_2 + \frac{1}{u_1} \int_{D_2(u)} x_1 dx_1 dx_2. \end{aligned} \tag{4.2}$$

Example 4.1. Consider the triangle D defined as

$$D = \{(x_1, x_2) \in \mathbb{R}_{++}^2 : 0 < x_1 \leq a, 0 < x_2 \leq vx_1\}. \tag{4.3}$$

Let $u \in D$. Assume that the \mathbb{R}_u is ray defined by the equation $x_2 = (u_2/u_1)x_1$. Since $u \in D$, we get $0 < u_2/u_1 \leq v$. Hence \mathbb{R}_u intersects the set D and divides the set into two parts D_1 and D_2 given as

$$\begin{aligned} D_1(u) &= \left\{ (x_1, x_2) \in \mathbb{R}_{++}^2 : 0 < x_1 \leq a, 0 < x_2 \leq \frac{u_2}{u_1}x_1 \right\} = \left\{ (x_1, x_2) \in D : \frac{x_2}{u_2} \leq \frac{x_1}{u_1} \right\}, \\ D_2(u) &= \left\{ (x_1, x_2) \in \mathbb{R}_{++}^2 : 0 < x_1 \leq a, \frac{u_2}{u_1}x_1 \leq x_2 \leq vx_1 \right\} = \left\{ (x_1, x_2) \in D : \frac{x_2}{u_2} \geq \frac{x_1}{u_1} \right\}. \end{aligned} \tag{4.4}$$

By (4.2) we get

$$\begin{aligned} \int_D \langle u, x \rangle dx &= \frac{1}{u_2} \int_{D_1(u)} x_2 dx_1 dx_2 + \frac{1}{u_1} \int_{D_2(u)} x_1 dx_1 dx_2 \\ &= \frac{1}{u_2} \int_0^a \int_0^{(u_2/u_1)x_1} x_2 dx_2 dx_1 + \frac{1}{u_1} \int_0^a \int_{(u_2/u_1)x_1}^{vx_1} x_1 dx_2 dx_1 \\ &= \frac{a^3 u_2}{6u_1^2} + \frac{(u_1 v - u_2)a^3}{3u_1^2} = \frac{(2u_1 v - u_2)a^3}{6u_1^2}. \end{aligned} \tag{4.5}$$

Thus, for the given region D , the inequality (3.1) will be as follows:

$$f(u_1, u_2) \leq \frac{6u_1^2}{a^3(2u_1 v - u_2)} \int_D f(x_1, x_2) dx_1 dx_2. \tag{4.6}$$

Since $A(D) = va^2/2$, then a point $x^* \in D$ belongs to $Q(D)$ if and only if

$$\frac{2}{va^2} \frac{(2x_1^* v - x_2^*)a^3}{6(x_1^*)^2} = 1 \iff x_2^* = -\frac{3v}{a}(x_1^*)^2 + 2vx_1^*. \tag{4.7}$$

Consider now the inequality (3.9) for triangle D . Let us calculate the integral of the function $\langle u, x \rangle^+$ on D :

$$\begin{aligned} \int_D \langle u, x \rangle^+ dx &= \frac{1}{u_1} \int_{D_1(u)} x_1 dx_1 dx_2 + \frac{1}{u_2} \int_{D_2(u)} x_2 dx_1 dx_2 \\ &= \frac{1}{u_1} \int_0^a \int_0^{(u_2/u_1)x_1} x_1 dx_2 dx_1 + \frac{1}{u_2} \int_0^a \int_{(u_2/u_1)x_1}^{vx_1} x_2 dx_2 dx_1 \\ &= \frac{a^3}{6} \left(\frac{u_2}{u_1^2} + \frac{v^2}{u_2} \right). \end{aligned} \quad (4.8)$$

Therefore,

$$\int_D f(x_1, x_2) dx_1 dx_2 \leq \frac{a^3}{6} \inf_{u \in D} \left\{ \left(\frac{u_2}{u_1^2} + \frac{v^2}{u_2} \right) f(u_1, u_2) \right\}. \quad (4.9)$$

Example 4.2. Let $D \subset \mathbb{R}_{++}^2$ be the triangle with vertices $(0, 0)$, $(a, 0)$ and $(0, b)$, that is

$$D = \left\{ x \in \mathbb{R}_{++}^2 : \frac{x_1}{a} + \frac{x_2}{b} \leq 1 \right\}. \quad (4.10)$$

If $u \in D$, then we get

$$\begin{aligned} D_1(u) &= \left\{ x \in \mathbb{R}_{++}^2 : 0 < x_2 < \frac{abu_2}{au_2 + bu_1}, \frac{u_1}{u_2} x_2 \leq x_1 \leq a - \frac{a}{b} x_2 \right\} \\ D_2(u) &= \left\{ x \in \mathbb{R}_{++}^2 : 0 < x_1 < \frac{abu_1}{au_2 + bu_1}, \frac{u_2}{u_1} x_1 \leq x_2 \leq b - \frac{b}{a} x_1 \right\}. \end{aligned} \quad (4.11)$$

By (4.2) we have

$$\begin{aligned} \int_D \langle u, x \rangle dx &= \frac{1}{u_2} \int_{D_1(u)} x_2 dx_1 dx_2 + \frac{1}{u_1} \int_{D_2(u)} x_1 dx_1 dx_2 \\ &= \frac{1}{u_2} \int_0^{abu_2/(au_2+bu_1)} \int_{(u_1/u_2)x_2}^{a-(a/b)x_2} x_2 dx_1 dx_2 + \frac{1}{u_1} \int_0^{abu_1/(au_2+bu_1)} \int_{(u_2/u_1)x_1}^{b-(b/a)x_1} x_1 dx_2 dx_1 \\ &= \frac{a^3 b^2 u_2}{6(au_2 + bu_1)^2} + \frac{a^2 b^3 u_1}{6(au_2 + bu_1)^2} = \frac{a^2 b^2}{6(au_2 + bu_1)} = \frac{ab}{6(u_1/a + u_2/b)}. \end{aligned} \quad (4.12)$$

In this triangular region D , the inequality (3.1) is as follows:

$$f(u_1, u_2) \leq \frac{6}{ab} \left(\frac{u_1}{a} + \frac{u_2}{b} \right) \int_D f(x_1, x_2) dx_1 dx_2. \quad (4.13)$$

Let us derive the set $Q(D)$ for the given triangular region D . Since $A(D) = ab/2$, then for $x^* \in D$,

$$x^* \in Q(D) \iff \frac{x_1^*}{a} + \frac{x_2^*}{b} = \frac{1}{3}. \quad (4.14)$$

Therefore,

$$Q(D) = \left\{ x^* \in D : \frac{x_1^*}{a} + \frac{x_2^*}{b} = \frac{1}{3} \right\}. \tag{4.15}$$

For the same region D , let us compute $\int_D \langle u, x \rangle^+ dx$ in order to derive the inequality (3.9):

$$\begin{aligned} \int_D \langle u, x \rangle^+ dx &= \frac{1}{u_1} \int_{D_1(u)} x_1 dx_1 dx_2 + \frac{1}{u_2} \int_{D_2(u)} x_2 dx_1 dx_2 \\ &= \frac{1}{2u_1} \left[\frac{a^3 bu_2}{au_2 + bu_1} - \frac{a^4 bu_2^2}{(au_2 + bu_1)^2} + \left(\frac{a^2}{b^2} - \frac{u_1^2}{u_2^2} \right) \frac{a^3 b^3 u_2^3}{3(au_2 + bu_1)^3} \right] \\ &\quad + \frac{1}{2u_2} \left[\frac{ab^3 u_1}{au_2 + bu_1} - \frac{b^4 au_1^2}{(au_2 + bu_1)^2} + \left(\frac{b^2}{a^2} - \frac{u_2^2}{u_1^2} \right) \frac{a^3 b^3 u_1^3}{3(au_2 + bu_1)^3} \right] \\ &= \frac{ab}{6} \left(\frac{au_2 + bu_1}{u_1 u_2} - \frac{1}{au_2 + bu_1} \right). \end{aligned} \tag{4.16}$$

Hence,

$$\int_D f(x_1, x_2) dx_1 dx_2 \leq \frac{ab}{6} \inf_{u \in D} \left\{ \left(\frac{au_2 + bu_1}{u_1 u_2} - \frac{1}{au_2 + bu_1} \right) f(u_1, u_2) \right\}. \tag{4.17}$$

Example 4.3. We will now consider the rectangle in \mathbb{R}_{++}^2 . Let D be the rectangle defined as

$$D = \{x \in \mathbb{R}_{++}^2 : x_1 \leq a, x_2 \leq b\}. \tag{4.18}$$

We consider two possible cases for $u \in D$.

(a) If $u_2/u_1 \leq b/a$, then we have

$$\begin{aligned} D_1(u) &= \left\{ x \in \mathbb{R}_{++}^2 : 0 < x_1 \leq a, 0 < x_2 \leq \frac{u_2}{u_1} x_1 \right\}, \\ D_2(u) &= \left\{ x \in \mathbb{R}_{++}^2 : 0 < x_1 \leq a, \frac{u_2}{u_1} x_1 \leq x_2 \leq b \right\}. \end{aligned} \tag{4.19}$$

Therefore,

$$\begin{aligned} \int_D \langle u, x \rangle dx &= \frac{1}{u_2} \int_{D_1(u)} x_2 dx_1 dx_2 + \frac{1}{u_1} \int_{D_2(u)} x_1 dx_1 dx_2 \\ &= \frac{1}{u_2} \int_0^a \int_0^{(u_2/u_1)x_1} x_2 dx_2 dx_1 + \frac{1}{u_1} \int_0^a \int_{(u_2/u_1)x_1}^b x_1 dx_2 dx_1 \\ &= \frac{1}{u_2} \frac{u_2^2 a^3}{6u_1^2} + \frac{1}{u_1} \left(\frac{ba^2}{2} - \frac{u_2 a^3}{u_1 3} \right) = \frac{3ba^2 u_1 - u_2 a^3}{6u_1^2}. \end{aligned} \tag{4.20}$$

By using the equality above, the inequality (3.1) will be as follows:

$$f(u_1, u_2) \leq \frac{6u_1^2}{3ba^2u_1 - u_2a^3} \int_D f(x_1, x_2) dx_1 dx_2. \quad (4.21)$$

Let us derive the set $Q(D)$. Since $A(D) = ab$, then we get the equation for $x^* \in Q(D)$,

$$\frac{1}{ab} \frac{3ba^2x_1^* - x_2^*a^3}{6(x_1^*)^2} = 1 \Leftrightarrow x_2^* = -\frac{6b}{a^2}(x_1^*)^2 + \frac{3b}{a}x_1^*. \quad (4.22)$$

(b) If $u_2/u_1 \geq b/a$, then by analogy

$$\int_D \langle u, x \rangle dx = \frac{3b^2au_2 - u_1b^3}{6u_2^2}. \quad (4.23)$$

Hence,

$$f(u_1, u_2) \leq \frac{6u_2^2}{3ab^2u_2 - u_1b^3} \int_D f(x_1, x_2) dx_1 dx_2. \quad (4.24)$$

We get the symmetric equation for $x^* \in Q(D)$:

$$x_1^* = -\frac{6a}{b^2}(x_2^*)^2 + \frac{3a}{b}x_2^*. \quad (4.25)$$

By taking into account both cases, $Q(D)$ becomes as the following:

$$Q(D) = \left\{ x^* \in D : \frac{x_2^*}{x_1^*} \leq \frac{b}{a}, x_2^* = -\frac{6b}{a^2}(x_1^*)^2 + \frac{3b}{a}x_1^* \right\} \cup \left\{ x^* \in D : \frac{x_2^*}{x_1^*} \geq \frac{b}{a}, x_1^* = -\frac{6a}{b^2}(x_2^*)^2 + \frac{3a}{b}x_2^* \right\}. \quad (4.26)$$

Consider now inequality (3.9). If $u_2/u_1 \leq b/a$, then $D_1(u)$ and $D_2(u)$ are stated as similar to (4.19). Consequently,

$$\int_D \langle u, x \rangle^+ dx = \frac{1}{u_1} \int_{D_1(u)} x_1 dx_1 dx_2 + \frac{1}{u_2} \int_{D_2(u)} x_2 dx_1 dx_2 = \frac{u_2a^3}{6u_1^2} + \frac{ab^2}{2u_2}. \quad (4.27)$$

If $u_2/u_1 \geq b/a$, then by analogy

$$\int_D \langle u, x \rangle^+ dx = \frac{u_1b^3}{6u_2^2} + \frac{ba^2}{2u_1}. \quad (4.28)$$

That is,

$$\int_D \langle u, x \rangle^+ dx = \varphi(u) = \begin{cases} \frac{u_2a^3}{6u_1^2} + \frac{ab^2}{2u_2}, & \text{if } \frac{u_2}{u_1} \leq \frac{b}{a}, \\ \frac{u_1b^3}{6u_2^2} + \frac{ba^2}{2u_1}, & \text{if } \frac{u_2}{u_1} \geq \frac{b}{a}. \end{cases} \quad (4.29)$$

Therefore

$$\int_D f(x_1, x_2) dx_1 dx_2 \leq \inf_{u \in D} \{f(u_1, u_2) \varphi(u_1, u_2)\}. \quad (4.30)$$

Acknowledgment

The authors were supported by the Scientific Research Project Administration Unit of the Akdeniz University (Turkey) and TÜBİTAK (Turkey).

References

- [1] S. S. Dragomir and C. E. M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, Melbourne City, Australia, 2000, <http://rgmia.vu.edu.au/monographs/>.
- [2] S. S. Dragomir and C. E. M. Pearce, "Quasi-convex functions and Hadamard's inequality," *Bulletin of the Australian Mathematical Society*, vol. 57, no. 3, pp. 377–385, 1998.
- [3] C. E. M. Pearce and A. M. Rubinov, "P-functions, quasi-convex functions, and Hadamard-type inequalities," *Journal of Mathematical Analysis and Applications*, vol. 240, no. 1, pp. 92–104, 1999.
- [4] A. M. Rubinov and J. Dutta, "Hadamard type inequality for quasiconvex functions in higher dimensions," preprint, RGMIA Res. Rep. Coll., 4(1) 2001, <http://rgmia.vu.edu.au/v4n1.html>.
- [5] S. S. Dragomir, J. Pečarić, and L. E. Persson, "Some inequalities of Hadamard type," *Soochow Journal of Mathematics*, vol. 21, no. 3, pp. 335–341, 1995.
- [6] P. M. Gill, C. E. M. Pearce, and J. Pečarić, "Hadamard's inequality for r -convex functions," *Journal of Mathematical Analysis and Applications*, vol. 215, no. 2, pp. 461–470, 1997.
- [7] S. S. Dragomir, J. Dutta, and A. M. Rubinov, "Hermite-Hadamard-type inequalities for increasing convex-along-rays functions," *Analysis*, vol. 24, no. 2, pp. 171–181, 2004, <http://rgmia.vu.edu.au/v4n4.html>.
- [8] E. V. Sharikov, "Hermite-Hadamard type inequalities for increasing radiant functions," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 2, pp. 1–13, 2003, article no. 47.
- [9] A. Rubinov, *Abstract Convexity and Global Optimization*, vol. 44 of *Nonconvex Optimization and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2000.

G. R. Adilov: Department of Mathematics, Akdeniz University, 07058 Antalya, Turkey
 Email address: gabil@akdeniz.edu.tr

S. Kemali: Department of Mathematics, Akdeniz University, 07058 Antalya, Turkey
 Email address: skemali@akdeniz.edu.tr