

Research Article

On Subordination Result Associated with Certain Subclass of Analytic Functions Involving Salagean Operator

Sevtap Sümer Eker, Bilal Şeker, and Shigeyoshi Owa

Received 3 February 2007; Accepted 15 May 2007

Recommended by Narendra K. Govil

We obtain an interesting subordination relation for Salagean-type certain analytic functions by using subordination theorem.

Copyright © 2007 Sevtap Sümer Eker et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let \mathcal{A} denote the class of functions $f(z)$ normalized by

$$f(z) = z + \sum_{j=2}^{\infty} a_j z^j, \tag{1.1}$$

which are analytic in the open unit disk $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$. We denote by $\mathcal{S}^*(\alpha)$ and $\mathcal{H}(\alpha)$ ($0 \leq \alpha < 1$) the class of starlike functions of order α and the class of convex functions of order α , respectively, where

$$\begin{aligned} \mathcal{S}^*(\alpha) &= \left\{ f \in \mathcal{A} : \operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) > \alpha, z \in \mathbb{U} \right\}, \\ \mathcal{H}(\alpha) &= \left\{ f \in \mathcal{A} : \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > \alpha, z \in \mathbb{U} \right\}. \end{aligned} \tag{1.2}$$

Note that $f(z) \in \mathcal{H}(\alpha) \Leftrightarrow zf'(z) \in \mathcal{S}^*(\alpha)$.

Sălăgean [1] has introduced the following operator:

$$\begin{aligned}
 D^0 f(z) &= f(z), \\
 D^1 f(z) &= Df(z) = zf'(z), \\
 &\vdots \\
 D^n f(z) &= D(D^{n-1} f(z)), \quad n \in \mathbb{N}_0 = \{0\} \cup \{1, 2, \dots\}.
 \end{aligned}
 \tag{1.3}$$

We note that

$$D^n f(z) = z + \sum_{j=2}^{\infty} j^n a_j z^j \quad (n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}).
 \tag{1.4}$$

We denote by $S_n(\alpha)$ subclass of the class \mathcal{A} which is defined as follows:

$$S_n(\alpha) = \left\{ f : f \in \mathcal{A}, \operatorname{Re} \left\{ \frac{D^{n+1} f(z)}{D^n f(z)} \right\} > \alpha \quad z \in \mathbb{U}; 0 < \alpha \leq 1 \right\}.
 \tag{1.5}$$

The class $S_n(\alpha)$ was introduced by Kadioğlu [2]. We begin by recalling following coefficient inequality associated with the function class $S_n(\alpha)$.

THEOREM 1.1 (Kadioğlu [2]). *If $f(z) \in \mathcal{A}$, defined by (1.1), satisfies the coefficient inequality*

$$\sum_{j=2}^{\infty} (j^{n+1} - \alpha j^n) |a_j| \leq 1 - \alpha, \quad 0 \leq \alpha < 1,
 \tag{1.6}$$

then $f(z) \in S_n(\alpha)$.

In view of Theorem 1.1, we now introduce the subclass

$$\tilde{S}_n(\alpha) \subset S_n(\alpha),
 \tag{1.7}$$

which consists of functions $f(z) \in \mathcal{A}$ whose Taylor-Maclaurin coefficients satisfy the inequality (1.6).

In this paper, we prove an interesting subordination result for the class $\tilde{S}_n(\alpha)$. In our proposed investigation of functions in the class $\tilde{S}_n(\alpha)$, we need the following definitions and results.

Definition 1.2 (Hadamard product or convolution). Given two functions $f, g \in \mathcal{A}$ where $f(z)$ is given by (1.1) and $g(z)$ is defined by

$$g(z) = z + \sum_{j=2}^{\infty} b_j z^j.
 \tag{1.8}$$

The Hadamard product (or convolution) $f * g$ is defined (as usual) by

$$(f * g)(z) = z + \sum_{j=2}^{\infty} a_j b_j z^j = (g * f)(z), \quad z \in \mathbb{U}. \tag{1.9}$$

Definition 1.3 (subordination principle). For two functions f and g analytic in \mathbb{U} , the function $f(z)$ is subordinate to $g(z)$ in \mathbb{U}

$$f(z) \prec g(z), \quad z \in \mathbb{U}, \tag{1.10}$$

if there exists a Schwarz function $w(z)$, analytic in \mathbb{U} with

$$w(0) = 0, \quad |w(z)| < 1, \tag{1.11}$$

such that

$$f(z) = g(w(z)), \quad z \in \mathbb{U}. \tag{1.12}$$

In particular, if the function g is univalent in \mathbb{U} , the above subordination is equivalent to

$$f(0) = g(0), \quad f(\mathbb{U}) \subset g(\mathbb{U}). \tag{1.13}$$

Definition 1.4 (subordinating factor sequence). A sequence $\{b_j\}_{j=1}^{\infty}$ of complex numbers is said to be a *subordinating factor sequence* if whenever $f(z)$ of the form (1.1) is analytic, univalent, and convex in \mathbb{U} , the subordination is given by

$$\sum_{j=1}^{\infty} a_j b_j z^j \prec f(z); \quad z \in \mathbb{U}, \quad a_1 = 1. \tag{1.14}$$

THEOREM 1.5 (Wilf [3]). *The sequence $\{b_j\}_{j=1}^{\infty}$ is subordinating factor sequence if and only if*

$$\operatorname{Re} \left\{ 1 + 2 \sum_{j=1}^{\infty} b_j z^j \right\} > 0 \quad z \in \mathbb{U}. \tag{1.15}$$

2. Main theorem

THEOREM 2.1. *Let the function $f(z)$ defined by (1.1) be in the class $\tilde{\mathcal{S}}_n(\alpha)$. Also, let \mathcal{K} denote familiar class of functions $f(z) \in \mathcal{A}$ which are univalent and convex in \mathbb{U} . Then*

$$\frac{2^n - \alpha 2^{n-1}}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} (f * g)(z) \prec g(z) \quad (z \in \mathbb{U}; n \in \mathbb{N}_0; g(z) \in \mathcal{K}), \tag{2.1}$$

$$\operatorname{Re} f(z) > -\frac{(1 - \alpha) + (2^{n+1} - \alpha 2^n)}{(2^{n+1} - \alpha 2^n)}. \tag{2.2}$$

The following constant factor in the subordination result (2.1):

$$\frac{2^n - \alpha 2^{n-1}}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} \tag{2.3}$$

cannot be replaced by a larger one.

Proof. Let $f(z) \in \tilde{S}_n(\alpha)$ and suppose that

$$g(z) = z + \sum_{j=2}^{\infty} c_j z^j \in \mathcal{H}. \tag{2.4}$$

Then

$$\frac{2^n - \alpha 2^{n-1}}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} (f * g)(z) = \frac{2^n - \alpha 2^{n-1}}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} \left(z + \sum_{j=2}^{\infty} a_j c_j z^j \right). \tag{2.5}$$

Thus, by Definition 1.4, the subordination result (2.1) will hold true if

$$\left\{ \frac{2^n - \alpha 2^{n-1}}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} a_j \right\}_{j=1}^{\infty} \tag{2.6}$$

is a subordinating factor sequences, with $a_1 = 1$. In view of Theorem 1.5, this is equivalent to the following inequality:

$$\operatorname{Re} \left\{ 1 + 2 \sum_{j=1}^{\infty} \frac{2^n - \alpha 2^{n-1}}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} a_j z^j \right\} > 0 \quad z \in \mathbb{U}. \tag{2.7}$$

Now, since $j^{n+1} - j^n$ ($j \geq 2, n \in \mathbb{N}_0$) is an increasing function of j , we have

$$\begin{aligned} & \operatorname{Re} \left\{ 1 + 2 \sum_{j=1}^{\infty} \frac{2^n - \alpha 2^{n-1}}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} a_j z^j \right\} \\ &= \operatorname{Re} \left\{ 1 + \sum_{j=1}^{\infty} \frac{2^{n+1} - \alpha 2^n}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} a_j z^j \right\} \\ &= \operatorname{Re} \left\{ 1 + \frac{2^{n+1} - \alpha 2^n}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} a_1 z + \frac{1}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} \sum_{j=2}^{\infty} (2^{n+1} - \alpha 2^n) a_j z^j \right\} \\ &\geq 1 - \frac{2^{n+1} - \alpha 2^n}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} r - \frac{1}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} \sum_{j=2}^{\infty} (j^{n+1} - \alpha j^n) |a_j| r^j \\ &> 1 - \frac{2^{n+1} - \alpha 2^n}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} r - \frac{1 - \alpha}{(1 - \alpha) + (2^{n+1} - \alpha 2^n)} r > 0 \quad (|z| = r < 1), \end{aligned} \tag{2.8}$$

where we have also made use of the assertion (1.6) of Theorem 1.1. This evidently proves the inequality (2.7), and hence also the subordination result (2.1) asserted by our theorem. The inequality (2.2) follows from (2.1) upon setting

$$g(z) = \frac{z}{1-z} = \sum_{j=1}^{\infty} z^j \in \mathcal{H}. \tag{2.9}$$

Now, consider the function

$$f_0(z) = z - \frac{1-\alpha}{2^{n+1}-\alpha 2^n} z^2 \quad (n \in \mathbb{N}_0, 0 \leq \alpha < 1), \tag{2.10}$$

which is a member of the class $\tilde{\mathcal{S}}_n(\alpha)$. Then by using (2.1), we have

$$\frac{2^n - \alpha 2^{n-1}}{(1-\alpha) + (2^{n+1} - \alpha 2^n)} (f_0 * g)(z) < \frac{z}{1-z}. \tag{2.11}$$

It can be easily verified for the function $f_0(z)$ defined by (2.10) that

$$\min \operatorname{Re} \left\{ \frac{2^n - \alpha 2^{n-1}}{(1-\alpha) + (2^{n+1} - \alpha 2^n)} (f_0 * g)(z) \right\} = -\frac{1}{2}, \quad z \in \mathbb{U}, \tag{2.12}$$

which completes the proof of theorem. □

If we take $n = 0$ in Theorem 2.1, we have the following corollary.

COROLLARY 2.2. *Let the function $f(z)$ defined by (1.1) be in the class $\mathcal{S}^*(\alpha)$ and $g(z) \in \mathcal{H}$, then*

$$\frac{2-\alpha}{2(3-2\alpha)} (f * g)(z) < g(z), \tag{2.13}$$

$$\operatorname{Re} f(z) > -\frac{3-2\alpha}{2-\alpha} \quad (z \in \mathbb{U}). \tag{2.14}$$

The constant factor

$$\frac{2-\alpha}{2(3-2\alpha)} \tag{2.15}$$

in the subordination result (2.13) cannot be replaced by a larger one.

If we take $n = 1$ in Theorem 2.1, we have the following corollary.

COROLLARY 2.3. *Let the function $f(z)$ defined by (1.1) be in the class $\mathcal{K}(\alpha)$ and $g(z) \in \mathcal{K}$, then*

$$\frac{2-\alpha}{5-3\alpha}(f * g)(z) < g(z), \quad (2.16)$$

$$\operatorname{Re} f(z) > -\frac{5-3\alpha}{2(2-\alpha)} \quad (z \in \mathbb{U}). \quad (2.17)$$

The constant factor

$$\frac{2-\alpha}{5-3\alpha} \quad (2.18)$$

in the subordination result (2.16) cannot be replaced by a larger one.

References

- [1] G. S. Sălăgean, "Subclasses of univalent functions," in *Complex Analysis—Proceedings of 5th Romanian-Finnish Seminar—Part 1 (Bucharest, 1981)*, vol. 1013 of *Lecture Notes in Math.*, pp. 362–372, Springer, Berlin, Germany, 1983.
- [2] E. Kadioğlu, "On subclass of univalent functions with negative coefficients," *Applied Mathematics and Computation*, vol. 146, no. 2-3, pp. 351–358, 2003.
- [3] H. S. Wilf, "Subordinating factor sequences for convex maps of the unit circle," *Proceedings of the American Mathematical Society*, vol. 12, pp. 689–693, 1961.

Sevtap Sümer Eker: Department of Mathematics, Faculty of Science and Letters, Dicle University, 21280 Diyarbakir, Turkey
Email address: sevtaps@dicle.edu.tr

Bilal Şeker: Department of Mathematics, Faculty of Science and Letters, Dicle University, 21280 Diyarbakir, Turkey
Email address: bseker@dicle.edu.tr

Shigeyoshi Owa: Department of Mathematics, Kinki University, Higashi-Osaka, Osaka 577-8502, Japan
Email address: owa@math.kindai.ac.jp