

Research Article

New Means of Cauchy's Type

Matloob Anwar¹ and J. Pečarić^{1,2}

¹Abdus Salam School of Mathematical Sciences, GC University, Lahore Gulberg 54660, Pakistan

²Faculty of Textile Technology, University of Zagreb, 10000 Zagreb, Croatia

Correspondence should be addressed to Matloob Anwar, matloob_t@yahoo.com

Received 30 December 2007; Accepted 7 April 2008

Recommended by Wing-Sum Cheung

We will introduce new means of Cauchy's type $M_{r,l}^s(f, \mu)$ defined, for example, as $M_{r,l}^s(f, \mu) = ((l(l-s)/r(r-s))(M_r^r(f, \mu) - M_s^r(f, \mu)/M_l^l(f, \mu) - M_s^l(f, \mu)))^{1/(r-l)}$, in the case when $l \neq r \neq s$, $l, r \neq 0$. We will show that this new Cauchy's mean is monotonic, that is, the following result. *Theorem.* Let $t, r, u, v \in \mathbb{R}$, such that $t \leq v$, $r \leq u$. Then for $M_{r,l}^s(f, \mu)$, one has $M_{t,r}^s \leq M_{v,u}^s$. We will also give some related comparison results.

Copyright © 2008 M. Anwar and J. Pečarić. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let Ω be a convex set equipped with a probability measure μ . Then for a strictly monotonic continuous function f , the integral power mean of order $r \in \mathbb{R}$ is defined as follows:

$$M_r(f, \mu) = \begin{cases} \left(\int_{\Omega} (f(u))^r d\mu(u) \right)^{1/r}, & r \neq 0, \\ \exp \left(\int_{\Omega} \log(f(u)) d\mu(u) \right), & r = 0. \end{cases} \quad (1.1)$$

Throughout our present investigation, we tacitly assume, without further comment, that all the integrals involved in our results exist.

The following inequality for differences of power means was obtained (see [1, Remark 8]):

$$\left| \frac{r(r-s)}{l(l-s)} \right| m \leq \left| \frac{M_r^r(f, \mu) - M_s^r(f, \mu)}{M_l^l(f, \mu) - M_s^l(f, \mu)} \right| \leq \left| \frac{r(r-s)}{l(l-s)} \right| M, \quad (1.2)$$

where $r, l, s \in \mathbb{R}$, $l \neq r \neq s$, $r, l \neq 0$ and where m and M are, respectively, the minimum and the maximum values of the function x^{r-l} on the image of $f(u)$ ($u \in \Omega$).

Let us note that (1.2) was obtained as consequence of the following result (see, e.g., [1, Corollary 1]).

Theorem 1.1. *Let $r, s, l \in \mathbb{R}$, and let Ω be a convex set equipped with a probability measure μ . Then,*

$$\frac{M_r^r(f, \mu) - M_s^r(f, \mu)}{M_l^l(f, \mu) - M_s^l(f, \mu)} = \frac{r(r-s)}{l(l-s)} \eta^{r-l} \quad (1.3)$$

for some η in the image of $f(u)$ ($u \in \Omega$), provided that the denominator on the left-hand side of (1.3) is non-zero.

We can also note that from (1.3) we can get the following form of (1.2):

$$\inf_{u \in \Omega} f(u) \leq \left(\frac{l(l-s)}{r(r-s)} \frac{M_r^r(f, \mu) - M_s^r(f, \mu)}{M_l^l(f, \mu) - M_s^l(f, \mu)} \right)^{1/(r-l)} \leq \sup_{u \in \Omega} f(u), \quad (1.4)$$

where $r, l, s \in \mathbb{R}$, $r \neq l \neq s$, $r, l \neq 0$. Moreover, (1.4) suggests introducing a new mean of Cauchy type. We will prove in Section 3 a comparison theorem for these means. Finally we will, in Section 4, give some applications.

2. New Cauchy's mean

From (1.4), we can define a new mean $M_{r,l}^s$ as follows:

$$M_{r,l}^s(f, \mu) = \left(\frac{l(l-s)}{r(r-s)} \frac{M_r^r(f, \mu) - M_s^r(f, \mu)}{M_l^l(f, \mu) - M_s^l(f, \mu)} \right)^{1/(r-l)}, \quad l \neq r \neq s, l, r \neq 0. \quad (2.1)$$

Now by taking $\lim_{l \rightarrow 0} M_{r,l}^s(f, \mu)$, we will get

$$\begin{aligned} M_{r,0}^s(f, \mu) &= M_{0,r}^s(f, \mu) = \lim_{l \rightarrow 0} M_{r,l}^s(f, \mu) \\ &= \left(\frac{s[M_r^r(f, \mu) - M_s^r(f, \mu)]}{r(r-s)[\log M_s(f, \mu) - \log M_0(f, \mu)]} \right)^{1/r}, \quad r \neq s, r, s \neq 0. \end{aligned} \quad (2.2)$$

Now by taking $\lim_{r \rightarrow s} M_{r,l}^s(f, \mu)$, we will get

$$\begin{aligned} \lim_{r \rightarrow s} M_{r,l}^s(f, \mu) &= M_{s,l}^s(f, \mu) = M_{l,s}^s(f, \mu) \\ &= \left(\frac{l(l-s)}{s} \frac{[\int f(u)^s \log f(u) d\mu(u) - M_s^s(f, \mu) \log M_s(f, \mu)]}{M_l^l(f, \mu) - M_s^l(f, \mu)} \right)^{1/(s-l)}, \quad l \neq s, l, s \neq 0. \end{aligned} \quad (2.3)$$

By similar way, we can calculate all the cases for $r, s, l \in \mathbb{R}$. Finally, we get the following definition of $M_{r,l}^s(f, \mu)$:

$$\begin{aligned}
M_{r,l}^s(f, \mu) &= \left(\frac{l(l-s)}{r(r-s)} \frac{M_r^r(f, \mu) - M_s^r(f, \mu)}{M_l^l(f, \mu) - M_s^l(f, \mu)} \right)^{1/(r-l)}, \quad l \neq r \neq s, l, r \neq 0; \\
M_{r,0}^s(f, \mu) &= M_{0,r}^s(f, \mu) = \left(\frac{s[M_r^r(f, \mu) - M_s^r(f, \mu)]}{r(r-s)[\log M_s(f, \mu) - \log M_0(f, \mu)]} \right)^{1/r}, \quad r \neq s, r, s \neq 0; \\
M_{s,l}^s(f, \mu) &= M_{l,s}^s(f, \mu) = \left(\frac{l(l-s) \int f(u)^s \log f(u) d\mu(u) - M_s^s(f, \mu) \log M_s(f, \mu)}{s[M_l^l(f, \mu) - M_s^l(f, \mu)]} \right)^{1/(s-l)}, \\
& \quad l \neq s, l, s \neq 0; \\
M_{s,0}^s(f, \mu) &= M_{0,s}^s(f, \mu) = \left(\frac{\int f(u)^s \log f(u) d\mu(u) - M_s^s(f, \mu) \log M_s(f, \mu)}{\log M_s(f, \mu) - \log M_0(f, \mu)} \right)^{1/s}, \quad s \neq 0; \\
M_{r,l}^0(f, \mu) &= \left(\frac{l^2(M_r^r(f, \mu) - M_0^r(f, \mu))}{r^2(M_l^l(f, \mu) - M_0^l(f, \mu))} \right)^{1/(r-l)}, \quad l, r \neq 0; \\
M_{r,0}^0(f, \mu) &= M_{0,r}^0(f, \mu) = \left(\frac{2[M_r^r(f, \mu) - M_0^r(f, \mu)]}{r^2[M_2^2(\log f, \mu) - M_1^2(\log f, \mu)]} \right)^{1/r}, \quad r \neq 0; \\
M_{t,t}^s &= \exp \left(-\frac{2t-s}{t(t-s)} + \frac{\int f^t \log f d\mu(u) - M_s^t(f, \mu) \log M_s(f, \mu)}{M_t^t(f, \mu) - M_s^t(f, \mu)} \right), \quad t \neq s; \\
M_{t,t}^0 &= \exp \left(-\frac{2}{t} + \frac{\int f^t \log f d\mu(u) - M_0^t(f, \mu) \log M_0(f, \mu)}{M_t^t(f, \mu) - M_0^t(f, \mu)} \right), \quad t \neq 0; \\
M_{0,0}^0 &= \exp \left(\frac{1 \int (\log f)^3 d\mu(u) - (\log M_0(f, \mu))^3}{3 \int (\log f)^2 d\mu(u) - (\log M_0(f, \mu))^2} \right), \\
M_{s,s}^s &= \exp \left(-\frac{1}{s} + \frac{\int f^s (\log f)^2 d\mu(u) - M_s^s(f, \mu) (\log M_s(f, \mu))^2}{2(\int f^s \log f d\mu(u) - (M_s^s(f, \mu) \log M_s(f, \mu)))} \right), \quad s \neq 0; \\
M_{0,0}^s &= \exp \left(\frac{1}{s} + \frac{\int (\log f)^2 d\mu(u) - (\log M_s(f, \mu))^2}{2(\int \log f d\mu(u) - \log M_s(f, \mu))} \right), \quad s \neq 0.
\end{aligned} \tag{2.4}$$

3. Monotonicity of new means

In this section, we will prove the monotonicity of (2.4). We need the following lemmas for log-convex function.

Lemma 3.1. Let f be log-convex function and if $x_1 \leq y_1$, $x_2 \leq y_2$, $x_1 \neq x_2$, $y_1 \neq y_2$, then the following inequality is valid:

$$\left(\frac{f(x_2)}{f(x_1)}\right)^{1/(x_2-x_1)} \leq \left(\frac{f(y_2)}{f(y_1)}\right)^{1/(y_2-y_1)}. \quad (3.1)$$

Proof. In [2, page 3] we have the following result for convex function f , with $x_1 \leq y_1$, $x_2 \leq y_2$, $x_1 \neq x_2$, $y_1 \neq y_2$:

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} \leq \frac{f(y_2) - f(y_1)}{y_2 - y_1}. \quad (3.2)$$

Putting $f = \log f$, we get

$$\log \left(\frac{f(x_2)}{f(x_1)}\right)^{1/(x_2-x_1)} \leq \log \left(\frac{f(y_2)}{f(y_1)}\right)^{1/(y_2-y_1)}, \quad (3.3)$$

after applying exponential function we get (3.1). \square

The following two lemmas are proved (for functionals) in [3] (Theorem 4 and Lemma 2, for Lemma 3.2 see also [4, Theorem 1]).

Lemma 3.2. Let us consider Λ_t defined as

$$\Lambda_t(g, \mu) = \begin{cases} \frac{M_t^t(g, \mu) - M_1^t(g, \mu)}{t(t-1)}, & t \neq 0, 1; \\ \log M_1(g, \mu) - \log M_0^t(g, \mu), & t = 0; \\ \int g \log g \mu - M_0(g, \mu) \log M_0(g, \mu), & t = 1. \end{cases} \quad (3.4)$$

Then, Λ_t is a log-convex function.

Lemma 3.3. Let us consider Λ_t defined as

$$\Lambda_t = \begin{cases} \frac{1}{t^2}(M_t^t(f, \mu) - M_0^t(f, \mu)), & t \neq 0; \\ \frac{1}{2}(M_2^2(\log f, \mu) - M_1^2(\log f, \mu)), & t = 0. \end{cases} \quad (3.5)$$

Then, Λ_t is a log-convex function.

Theorem 3.4. Let $t, r, u, v \in \mathbb{R}$, such that, $t \leq v$, $r \leq u$. Then for (2.4), we have

$$M_{t,r}^s \leq M_{v,u}^s. \quad (3.6)$$

Proof

Case 1 ($s \neq 0$). Let us consider Λ_t defined as in Lemma 3.2. Λ_t is a continuous and log-convex. So, Lemma 3.1 implies that for $t, r, u, v \in \mathbb{R}$, such that, $t \leq v$, $r \leq u$, $t \neq r$, $v \neq u$, we have

$$\left(\frac{\Lambda_t}{\Lambda_r}\right)^{1/(t-r)} \leq \left(\frac{\Lambda_v}{\Lambda_u}\right)^{1/(v-u)}. \quad (3.7)$$

For $s > 0$ by substituting $g = f^s$, $t = t/s$, $r = r/s$, $u = u/s$, $v = v/s \in \mathbb{R}$, such that, $t/s \leq v/s$, $r/s \leq u/s$, $t \neq r$, $v \neq u$, in (3.4), we get

$$\Lambda_{t,s}(f, \mu) = \begin{cases} \frac{s^2}{t(1-s)} [M_t^t(f, \mu) - M_s^t(f, \mu)], & t \neq 0, s; \\ s(\log M_s(f, \mu) - \log M_0(f, \mu)), & t = 0; \\ s\left(\int f^s \log f - M_0^s(f, \mu) \log M_0(f, \mu)\right), & t = s. \end{cases} \quad (3.8)$$

And (3.7) becomes

$$\left(\frac{\Lambda_{t,s}}{\Lambda_{r,s}}\right)^{1/(t-r)} \leq \left(\frac{\Lambda_{v,s}}{\Lambda_{u,s}}\right)^{1/(v-u)}. \quad (3.9)$$

From (3.9), we get our required result.

Now when $s < 0$ by substituting $g = f^s$, $t = t/s$, $r = r/s$, $u = u/s$, $v = v/s \in \mathbb{R}$, such that, $v/s \leq t/s$, $u/s \leq r/s$, $t \neq r$, $v \neq u$, in (3.4) we get (3.8).

And (3.7) becomes

$$\left(\frac{\Lambda_{v,s}}{\Lambda_{u,s}}\right)^{s/(v-u)} \leq \left(\frac{\Lambda_{t,s}}{\Lambda_{r,s}}\right)^{s/(t-r)}. \quad (3.10)$$

Now $s < 0$, from (3.10), by raising power $-s$, we get

$$\left(\frac{\Lambda_{t,s}}{\Lambda_{r,s}}\right)^{1/(t-r)} \leq \left(\frac{\Lambda_{v,s}}{\Lambda_{u,s}}\right)^{1/(v-u)}. \quad (3.11)$$

From (3.11), we get our required result.

Case 2 ($s = 0$). In this case, we can get our result by taking limit $s \rightarrow 0$ in (3.8) and also in this case we can consider Λ_t defined as in Lemma 3.3.

Λ_t is log-convex function. So, Lemma 3.1 implies that for $t, r, u, v \in \mathbb{R}$, such that, $t \leq v$, $r \leq u$, $t \neq r$, $v \neq u$, we have

$$\left(\frac{\Lambda_t}{\Lambda_r}\right)^{1/(t-r)} \leq \left(\frac{\Lambda_v}{\Lambda_u}\right)^{1/(v-u)}. \quad (3.12)$$

Therefore, we have for $t, r, u, v \in \mathbb{R}$, such that, $t \leq v$, $r \leq u$, $t \neq r$, $v \neq u$:

$$M_{t,r}^0 \leq M_{v,u}^0, \quad (3.13)$$

which completes the proof. \square

4. Further consequences and applications

In this section, we will represent the various applications of our previous definition of a new Cauchy mean and monotonicity of this above defined a new Cauchy mean.

4.1. Tobey and Stolarsky-Tobey means

Let E_{n-1} represent the $(n - 1)$ -dimensional Euclidean simplex given by

$$E_{n-1} = \left\{ (u_1, u_2, \dots, u_{n-1}) : u_i \geq 0, 1 \leq i \leq n-1, \sum_{i=1}^{n-1} u_i \leq 1 \right\}, \quad (4.1)$$

and set $u_n = 1 - \sum_{i=1}^{n-1} u_i$. Moreover, with $u = (u_1, \dots, u_n)$, let $\mu(u)$ be a probability measure on E_{n-1} . The power mean of order p ($p \in \mathbb{R}$) of the positive n -tuple $x = (x_1, \dots, x_n) \in \mathbb{R}_+^n$, with the weights $u = (u_1, \dots, u_n)$, is defined by

$$\overline{M}_p(x, \mu) = \begin{cases} \left(\sum_{i=1}^n u_i x_i^p \right)^{1/p}, & p \neq 0; \\ \prod_{i=1}^n x_i^{u_i}, & p = 0. \end{cases} \quad (4.2)$$

Then, the Tobey mean $L_{p,r}(x; \mu)$ is defined as follows:

$$L_{p,r}(x; \mu) = M_r(\overline{M}_p(x, \mu); \mu), \quad (4.3)$$

where $M_r(g, \mu)$ denotes the integral power mean, in which Ω is now the $(n - 1)$ -dimensional Euclidean simplex E_{n-1} . We note that, since $\overline{M}_p(x, \mu)$ is a mean we have $\min\{x_i\} \leq \overline{M}_p(x, \mu) \leq \max\{x_i\}$. Now setting $f(x, \mu) = \overline{M}_p(x, \mu)$ in (2.4) we get

$$\Gamma_{p,r,l}^s(x, \mu) = \left(\frac{l(l-s) L_{p,r}^r(x, \mu) - L_{p,s}^r(x, \mu)}{r(r-s) L_{p,l}^l(x, \mu) - L_{p,s}^l(x, \mu)} \right)^{1/(r-l)}, \quad l \neq r \neq s, l, r \neq 0;$$

$$\Gamma_{p,r,0}^s(x, \mu) = \Gamma_{p,0,r}^s(x, \mu) = \left(\frac{s[L_{p,r}^r(x, \mu) - L_{p,s}^r(x, \mu)]}{r(r-s)[\log L_{p,s}(x, \mu) - \log L_{p,0}(x, \mu)]} \right)^{1/r}, \quad r \neq s, r, s \neq 0;$$

$$\Gamma_{p,s,l}^s(x, \mu) = \Gamma_{p,l,s}^s(x, \mu) = \left(\frac{l(l-s) \int \overline{M}_p(x, \mu)^s \log d\mu(u) - L_{p,s}^s(x, \mu) \log L_{p,s}(x, \mu)}{s [L_{p,l}^l(x, \mu) - L_{p,s}^l(x, \mu)]} \right)^{1/(s-l)},$$

$l \neq s, l, s \neq 0;$

$$\Gamma_{p,s,0}^s(x, \mu) = \Gamma_{p,0,s}^s(x, \mu) = \left(\frac{\int \overline{M}_p(x, \mu)^s \log \overline{M}_p(x, \mu) d\mu(u) - L_{p,s}^s(x, \mu) \log L_{p,s}(x, \mu)}{\log L_{p,s}(x, \mu) - \log L_{p,0}(x, \mu)} \right)^{1/s}, \quad s \neq 0;$$

$$\begin{aligned}
\Gamma_{p,r,l}^0(x,\mu) &= \left(\frac{l^2(L_{p,r}^r(x,\mu) - L_{p,0}^r(x,\mu))}{r^2(L_{p,l}^l(x,\mu) - L_{p,0}^l(x,\mu))} \right)^{1/(r-l)}, \quad l, r \neq 0; \\
\Gamma_{p,r,0}^0(x,\mu) &= \Gamma_{p,0,r}^0(x,\mu) = \left(\frac{2[L_{p,r}^r(x,\mu) - L_{p,0}^r(x,\mu)]}{r^2[M_2^2(\log \bar{M}_p(x,\mu), \mu) - M_1^2(\log \bar{M}_p(x,\mu), \mu)]} \right)^{1/r}, \quad r \neq 0; \\
\Gamma_{p,t,t}^s(x,\mu) &= \exp \left(-\frac{2t-s}{t(t-s)} + \frac{\int \bar{M}_p(x,\mu)^t \log \bar{M}_p(x,\mu) d\mu(u) - L_{p,s}^t(x,\mu) \log L_{p,s}(x,\mu)}{L_{p,t}^t(x,\mu) - L_{p,s}^t(x,\mu)} \right), \quad t \neq s; \\
\Gamma_{p,t,t}^0(x,\mu) &= \exp \left(-\frac{2}{t} + \frac{\int \bar{M}_p(x,\mu)^t \log \bar{M}_p(x,\mu) d\mu(u) - L_{p,0}^t(x,\mu) \log L_{p,0}(x,\mu)}{L_{p,t}^t(x,\mu) - L_{p,0}^t(x,\mu)} \right), \quad t \neq 0; \\
\Gamma_{p,0,0}^0(x,\mu) &= \exp \left(\frac{1 \int (\log \bar{M}_p(x,\mu))^3 d\mu(u) - (\log L_{p,0}(x,\mu))^3}{3 \int (\log \bar{M}_p(x,\mu))^2 d\mu(u) - (\log L_{p,0}(x,\mu))^2} \right), \\
\Gamma_{p,s,s}^s(x,\mu) &= \exp \left(-\frac{1}{s} + \frac{\int \bar{M}_p(x,\mu)^s (\log \bar{M}_p(x,\mu))^2 d\mu(u) - L_{p,s}^s(x,\mu) (\log L_{p,s}(x,\mu))^2}{2(\int \bar{M}_p(x,\mu)^s \log \bar{M}_p(x,\mu) d\mu(u) - (L_{p,s}^s(x,\mu) \log L_{p,s}(x,\mu)))} \right), \quad s \neq 0; \\
\Gamma_{p,0,0}^s(x,\mu) &= \exp \left(\frac{1}{s} + \frac{\int (\log \bar{M}_p(x,\mu))^2 d\mu(u) - (\log L_{p,s}(x,\mu))^2}{2(\int \log \bar{M}_p(x,\mu) d\mu(u) - \log M_s(x,\mu))} \right), \quad s \neq 0.
\end{aligned} \tag{4.4}$$

Theorem 4.1. Let $t, r, u, v \in \mathbb{R}$, such that, $t < v$, $r < u$. Then for (4.4), we have

$$\Gamma_{p,t,r}^s \leq \Gamma_{p,v,u}^s. \tag{4.5}$$

Proof. It is a simple consequence of Theorem 3.4. \square

Pečarić and Šimić (see [5, Definition 1]) introduced the Stolarsky-Tobey mean $\varepsilon_{p,q}(x,\mu)$ defined by

$$\varepsilon_{p,q}(x,\mu) = L_{p,q-p}(x,\nu) = M_{q-p}(\bar{M}_p(x,\mu); \mu), \tag{4.6}$$

where $L_{p,r}(x,\nu)$ is the Tobey mean already introduced above.

For the Stolarsky-Tobey mean and (2.4), we get the following:

$$\begin{aligned}
\Upsilon_{p,r,l}^s(x,\mu) &= \left(\frac{l(l-s) \varepsilon_{p,p+r}^r(x,\mu) - \varepsilon_{p,p+s}^r(x,\mu)}{r(r-s) \varepsilon_{p,p+l}^l(x,\mu) - \varepsilon_{p,p+s}^l(x,\mu)} \right)^{1/(r-l)}, \quad l \neq r \neq s, \quad l, r \neq 0; \\
\Upsilon_{p,r,0}^s(x,\mu) &= \Upsilon_{p,0,r}^s(x,\mu) = \left(\frac{s[\varepsilon_{p,p+r}^r(x,\mu) - \varepsilon_{p,p+s}^r(x,\mu)]}{r(r-s)[\log \varepsilon_{p,p+s}(x,\mu) - \log \varepsilon_{p,p}(x,\mu)]} \right)^{1/r}, \quad r \neq s, \quad r, s \neq 0;
\end{aligned}$$

$$\begin{aligned}
\Upsilon_{p,s,l}^s(x, \mu) &= \Upsilon_{p,l,s}^s(x, \mu) = \left(\frac{l(l-s) \int \overline{M}_p(x, \mu)^s \log d\mu(u) - \varepsilon_{p,p+s}^s(x, \mu) \log \varepsilon_{p,p+s}(x, \mu)}{\varepsilon_{p,p+l}^l(x, \mu) - \varepsilon_{p,p+s}^l(x, \mu)} \right)^{1/(s-l)}, \\
& \quad l \neq s, l, s \neq 0; \\
\Upsilon_{p,s,0}^s(x, \mu) &= \Upsilon_{p,0,s}^s(x, \mu) = \left(\frac{\int \overline{M}_p(x, \mu)^s \log \overline{M}_p(x, \mu) d\mu(u) - \varepsilon_{p,p+s}^s(x, \mu) \log \varepsilon_{p,p+s}(x, \mu)}{\log \varepsilon_{p,p+s}(x, \mu) - \log \varepsilon_{p,p}(x, \mu)} \right)^{1/s}, \\
& \quad s \neq 0; \\
\Upsilon_{p,r,l}^0(x, \mu) &= \left(\frac{l^2 (\varepsilon_{p,p+r}^r(x, \mu) - \varepsilon_{p,p}^r(x, \mu))}{r^2 (\varepsilon_{p,p+l}^l(x, \mu) - \varepsilon_{p,p}^l(x, \mu))} \right)^{1/(r-l)}, \quad l, r \neq 0; \\
\Upsilon_{p,r,0}^0(x, \mu) &= \Upsilon_{p,0,r}^0(x, \mu) = \left(\frac{2[\varepsilon_{p,p+r}^r(x, \mu) - \varepsilon_{p,p}^r(x, \mu)]}{r^2 [M_2^2(\log \overline{M}_p(x, \mu), \mu) - M_1^2(\log \overline{M}_p(x, \mu), \mu)]} \right)^{1/r}, \quad r \neq 0; \\
\Upsilon_{p,t,t}^s(x, \mu) &= \exp \left(-\frac{2t-s}{t(t-s)} + \frac{\int \overline{M}_p(x, \mu)^t \log \overline{M}_p(x, \mu) d\mu(u) - M_s^t \log \varepsilon_{p,p+s}(x, \mu)}{\varepsilon_{p,p+t}^t(x, \mu) - \varepsilon_{p,p+s}^t(x, \mu)} \right), \quad t \neq s; \\
\Upsilon_{p,t,t}^0(x, \mu) &= \exp \left(-\frac{2}{t} + \frac{\int \overline{M}_p(x, \mu)^t \log \overline{M}_p(x, \mu) d\mu(u) - \varepsilon_{p,p}^t(x, \mu) \log \varepsilon_{p,p}(x, \mu)}{\varepsilon_{p,p+t}^t(x, \mu) - \varepsilon_{p,p}^t(x, \mu)} \right), \quad t \neq 0; \\
\Upsilon_{p,0,0}^0(x, \mu) &= \exp \left(\frac{\frac{1}{3} \int (\log \overline{M}_p(x, \mu))^3 d\mu(u) - (\log \varepsilon_{p,p}(x, \mu))^3}{\int (\log \overline{M}_p(x, \mu))^2 d\mu(u) - (\log \varepsilon_{p,p}(x, \mu))^2} \right), \\
\Upsilon_{p,s,s}^s(x, \mu) &= \exp \left(-\frac{1}{s} + \frac{\int \overline{M}_p(x, \mu)^s (\log \overline{M}_p(x, \mu))^2 d\mu(u) - \varepsilon_{p,p+s}^s(x, \mu) (\log \varepsilon_{p,p+s}(x, \mu))^2}{2 \left(\int \overline{M}_p(x, \mu)^s \log \overline{M}_p(x, \mu) d\mu(u) - (\varepsilon_{p,p+s}^s(x, \mu) \log \varepsilon_{p,p+s}(x, \mu)) \right)} \right), \\
& \quad s \neq 0; \\
\Upsilon_{p,0,0}^s(x, \mu) &= \exp \left(\frac{1}{s} + \frac{\int (\log \overline{M}_p(x, \mu))^2 d\mu(u) - (\log \varepsilon_{p,p+s}(x, \mu))^2}{2 \left(\int \log \overline{M}_p(x, \mu) d\mu(u) - \log \varepsilon_{p,p+s}(x, \mu) \right)} \right), \quad s \neq 0.
\end{aligned} \tag{4.7}$$

Theorem 4.2. Let $t, r, u, v \in \mathbb{R}$, such that, $t < v$, $r < u$. Then for (4.7), we have

$$\Upsilon_{p,t,r}^s \leq \Upsilon_{p,v,u}^s. \tag{4.8}$$

Proof. It is a simple consequence of Theorem 3.4. □

4.2. The complete symmetric mean

The r th complete symmetric polynomial mean (the complete symmetric mean) of the positive real n -tuple x is defined by (see [6, pages 332,341])

$$Q_n^{[r]}(x) = \left(q_n^{[r]}(x) \right)^{1/r} = \left(\frac{c_n^{[r]}(x)}{\binom{n+r-1}{r}} \right)^{1/r}, \quad (4.9)$$

where $c_n^{[0]}(x) = 1$ and $c_n^{[r]}(x) = \sum_{j=1}^n \left(\prod_{i=1}^n x_i^{i_j} \right)$ and the sum is taken over all $\binom{n+r-1}{r}$ nonnegative integer n -tuples (i_1, \dots, i_n) with $\sum_{j=1}^n i_j = r$ ($r \neq 0$). The complete symmetric polynomial mean can also be written in an integral form as follows:

$$Q_n^{[r]} = \left(\int_{E_{n-1}} \left(\sum_{i=1}^n x_i u_i \right)^r d\mu(u) \right)^{1/r}, \quad (4.10)$$

where μ represents a probability measure such that $d\mu(u) = (n-1)! du_1 \cdots du_{n-1}$. We can see this as a special case of the integral power mean $M_r(f, \mu)$, where $f(u) = \sum_{i=1}^n x_i u_i$, μ is a probability measure as above, and Ω is the above defined $(n-1)$ -dimensional simplex E_{n-1} . Thus from (2.4), we have the following result:

$$\Theta_{n,r,l}^s(x, \mu) = \left(\frac{l(l-s)}{r(r-s)} \frac{(Q_n^{[r]})^r(x, \mu) - (Q_n^{[s]})^r(x, \mu)}{(Q_n^{[l]})^l(x, \mu) - (Q_n^{[s]})^l(x, \mu)} \right)^{1/(r-l)}, \quad l \neq r \neq s, \quad l, r \in \mathbb{N}. \quad (4.11)$$

A simple consequence of Theorem 3.4 is the following result.

Theorem 4.3. *Let $t, r, u, v \in \mathbb{N}$, such that, $t < v$, $r < u$. Then for (4.11), we have*

$$\Theta_{n,t,r}^s \leq \Theta_{n,v,u}^s. \quad (4.12)$$

4.3. Whiteley means

Let x be a positive real n -tuple, $s \in \mathbb{R}$ ($s \neq 0$) and $r \in \mathbb{N}$. Then, the s th function of degree r is defined by the following generating function (see [6, pages 341–344]):

$$\sum_{r=0}^{\infty} t_n^{[r,s]}(x) t^r = \begin{cases} \prod_{i=1}^n (1 + x_i t)^s, & s > 0, \\ \prod_{i=1}^n (1 - x_i t)^s, & s < 0. \end{cases} \quad (4.13)$$

The Whiteley mean is now defined by

$$\mathcal{W}_n^{[r,s]}(x) = \left(w_n^{[r,s]}(x) \right)^{1/r} = \begin{cases} \left(\frac{t_n^{[r,s]}(x)}{\binom{nr}{s}} \right)^{1/r}, & s > 0, \\ \left(\frac{t_n^{[r,s]}(x)}{(-1)^r \binom{nr}{s}} \right)^{1/r}, & s < 0. \end{cases} \quad (4.14)$$

For $s < 0$, the Whiteley mean can be further generalized if we slightly change the definition of $t_n^{[r,s]}(x)$ and define $h_n^{[r,\sigma]}(x)$ as follows:

$$\sum_{r=0}^{\infty} h_n^{[r,\sigma]}(x)t^r = \prod_{i=1}^n \frac{1}{(1-x_it)^{\sigma_i}}, \quad (4.15)$$

where $\sigma = (\sigma_1, \dots, \sigma_n)$; $\sigma \in \mathbb{R}_+$; $i = 1, \dots, n$. The following generalization of the Whiteley mean for $s < 0$ is defined by (see [7, Lemma 2.3])

$$\mathcal{H}_n^{[r,\sigma]}(x) = \left(\frac{h_n^{[r,\sigma]}(x)}{\left(\sum_{i=1}^n \sigma_i + r - 1 \right)} \right)^{1/r}. \quad (4.16)$$

If we denote by μ a measure on the simplex $\Delta_{n-1} = \{(u_1, \dots, u_n) : u_i \geq 0, i = 1, \dots, n-1, \sum_{i=1}^n u_i \leq 1\}$ such that

$$d\mu(u) = \frac{\Gamma(\sum_{i=1}^n \sigma_i)}{\prod_{i=1}^n \Gamma(\sigma_i)} \prod_{i=1}^n u_i^{\sigma_i-1} du_1 \cdots du_{n-1}, \quad (4.17)$$

where $u_n = 1 - \sum_{i=1}^{n-1} u_i$, then we have μ as a probability measure and we can also write the mean $\mathcal{H}_n^{[r,\sigma]}(x)$ in integral form as follows:

$$\mathcal{H}_n^{[r,\sigma]}(x) = \left(\int_{\Delta_{n-1}} \left(\sum_{i=1}^n x_i u_i \right)^r d\mu(u) \right)^{1/r}. \quad (4.18)$$

Finally, just as we did above in this investigation, we can develop the following analogous definition:

$$\mathfrak{H}_{n,r,l}^s(x, \mu) = \left(\frac{l(l-s) \left(\mathcal{H}_n^{[r,\sigma]} \right)^r(x, \mu) - \left(\mathcal{H}_n^{[s,\sigma]} \right)^r(x, \mu)}{r(r-s) \left(\mathcal{H}_n^{[l,\sigma]} \right)^l(x, \mu) - \left(\mathcal{H}_n^{[s,\sigma]} \right)^l(x, \mu)} \right)^{1/(r-l)}, \quad l \neq r \neq s, l, r \in \mathbb{N}. \quad (4.19)$$

A simple consequence of Theorem 3.4 is the following result.

Theorem 4.4. *Let $t, r, u, v \in \mathbb{N}$, such that $t < v$, $r < u$. Then for (4.19), we have*

$$\mathfrak{H}_{n,t,r}^s \leq \mathfrak{H}_{n,v,u}^s. \quad (4.20)$$

References

- [1] J. Pečarić, M. R. Lipanović, and H. M. Srivastava, "Some mean-value theorems of the Cauchy type," *Fractional Calculus & Applied Analysis*, vol. 9, no. 2, pp. 143–158, 2006.
- [2] J. Pečarić, F. Proschan, and Y. L. Tong, *Convex Functions, Partial Orderings, and Statistical Applications*, vol. 187 of *Mathematics in Science and Engineering*, Academic Press, Boston, Mass, USA, 1992.
- [3] M. Anwar and J. Pečarić, "On logarithmic convexity for differences of power means," to appear in *Mathematical Inequalities & Applications*.
- [4] S. Simić, "On logarithmic convexity for differences of power means," *Journal of Inequalities and Applications*, vol. 2007, Article ID 37359, 8 pages, 2007.
- [5] J. Pečarić and V. Šimić, "Stolarsky-Tobey mean in n variables," *Mathematical Inequalities & Applications*, vol. 2, no. 3, pp. 325–341, 1999.
- [6] P. S. Bullen, *Handbook of Means and Their Inequalities*, vol. 560 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2003.
- [7] J. Pečarić, I. Perić, and M. R. Lipanović, "Generalized Whiteley means and related inequalities," to appear in *Mathematical Inequalities & Applications*.