

Research Article

Certain Integral Operators on the Classes $\mathcal{M}(\beta_i)$ and $\mathcal{N}(\beta_i)$

Daniel Breaz

Department of Mathematics, 1st December 1918, University of Alba Iulia, 510009 Alba, Romania

Correspondence should be addressed to Daniel Breaz, dbreaz@uab.ro

Received 13 September 2007; Revised 21 October 2007; Accepted 2 January 2008

Recommended by Vijay Gupta

We consider the classes $\mathcal{M}(\beta_i)$ and $\mathcal{N}(\beta_i)$ of the analytic functions and two general integral operators. We prove some properties for these operators on these classes.

Copyright © 2008 Daniel Breaz. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let $\mathbf{U} = \{z \in \mathbf{C}, |z| < 1\}$ be the open unit disk and let \mathcal{A} denote the class of the functions $f(z)$ of the form

$$f(z) = z + a_2 z^2 + a_3 z^3 + \dots, \quad z \in \mathbf{U}, \quad (1.1)$$

which are analytic in the open disk \mathbf{U} .

Let $\mathcal{M}(\beta)$ be the subclass of \mathcal{A} , consisting of the functions $f(z)$, which satisfy the inequality

$$\operatorname{Re} \left\{ \frac{z f'(z)}{f(z)} \right\} < \beta, \quad z \in \mathbf{U}, \quad \beta > 1, \quad (1.2)$$

and let $\mathcal{N}(\beta)$ be the subclass of \mathcal{A} , consisting of functions $f(z)$, which satisfy the inequality

$$\operatorname{Re} \left\{ \frac{z f''(z)}{f'(z)} + 1 \right\} < \beta, \quad z \in \mathbf{U}. \quad (1.3)$$

These classes are studied by Uralegaddi et al. in [1], and Owa and Srivastava in [2].

Consider the integral operator F_n introduced by D. Breaz and N. Breaz in [3], having the form

$$F_n(z) = \int_0^z \left(\frac{f_1(t)}{t} \right)^{\alpha_1} \dots \left(\frac{f_n(t)}{t} \right)^{\alpha_n} dt, \quad (1.4)$$

where $f_i(z) \in \mathcal{A}$ and $\alpha_i > 0$, for all $i \in \{1, \dots, n\}$.

Remark 1.1. This operator extends the integral operator of Alexander given by $F(z) = \int_0^z (f(t)/t)dt$.

Also, we consider the next integral operator denoted by $F_{\alpha_1, \dots, \alpha_n}$ that was introduced by Breaz et al. in [4], having the form

$$F_{\alpha_1, \dots, \alpha_n}(z) = \int_0^z [f_1'(t)]^{\alpha_1} \cdots [f_n'(t)]^{\alpha_n} dt, \quad (1.5)$$

where $f_i(z) \in \mathcal{A}$ and $\alpha_i > 0$ for all $i \in \{1, \dots, n\}$.

It is easy to see that these integral operators are analytic operators.

2. Main results

Theorem 2.1. Let $f_i \in \mathcal{M}(\beta_i)$, for each $i = 1, 2, 3, \dots, n$ with $\beta_i > 1$. Then $F_n(z) \in \mathcal{N}(\mu)$ with $\mu = 1 + \sum_{i=1}^n \alpha_i(\beta_i - 1)$ and $\alpha_i > 0$, ($i = 1, 2, 3, \dots, n$).

Proof. After some calculi, we obtain that

$$\frac{zF_n''(z)}{F_n'(z)} = \sum_{i=1}^n \alpha_i \frac{zf_i'(z)}{f_i(z)} - \sum_{i=1}^n \alpha_i. \quad (2.1)$$

The relation (2.1) is equivalent to

$$\operatorname{Re}\left(\frac{zF_n''(z)}{F_n'(z)} + 1\right) = \sum_{i=1}^n \alpha_i \operatorname{Re}\left(\frac{zf_i'(z)}{f_i(z)}\right) - \sum_{i=1}^n \alpha_i + 1. \quad (2.2)$$

Since $f_i \in \mathcal{M}(\beta_i)$, we have

$$\operatorname{Re}\left(\frac{zF_n''(z)}{F_n'(z)} + 1\right) < \sum_{i=1}^n \alpha_i \beta_i - \sum_{i=1}^n \alpha_i + 1 = \sum_{i=1}^n \alpha_i(\beta_i - 1) + 1. \quad (2.3)$$

Because $\sum_{i=1}^n \alpha_i(\beta_i - 1) > 0$, we obtain that $F_n \in \mathcal{N}(\mu)$, where $\mu = 1 + \sum_{i=1}^n \alpha_i(\beta_i - 1)$. \square

Corollary 2.2. Let $f_i \in \mathcal{M}(\beta)$ for each $i = 1, 2, 3, \dots, n$ with $\beta > 1$. Then $F_n(z) \in \mathcal{N}(\gamma)$ with $\gamma = 1 + (\beta - 1)\sum_{i=1}^n \alpha_i$ and $\alpha_i > 0$, ($i = 1, 2, 3, \dots, n$).

Proof. In Theorem 2.1, we consider $\beta_1 = \beta_2 = \cdots = \beta_n = \beta$. \square

Corollary 2.3. Let $f \in \mathcal{M}(\beta)$ with $\beta > 1$. Then the integral operator $F(z) = \int_0^z (f(t)/t)^\alpha dt \in \mathcal{N}(\delta)$ with $\delta = \alpha(\beta - 1) + 1$ and $\alpha > 0$.

Proof. In Corollary 2.2, we consider $n = 1$ and $\alpha_1 = \alpha$. \square

Corollary 2.4. Let $f \in \mathcal{M}(\beta)$ with $\beta > 1$. Then the integral operator of Alexander $F(z) = \int_0^z (f(t)/t)dt \in \mathcal{N}(\beta)$.

Proof. We have

$$\frac{zF''(z)}{F'(z)} = \frac{zf'(z)}{f(z)} - 1. \quad (2.4)$$

From (2.4), we have

$$\operatorname{Re}\left(\frac{zF''(z)}{F'(z)} + 1\right) = \operatorname{Re}\frac{zf'(z)}{f(z)} < \beta. \quad (2.5)$$

So relation (2.5) implies that Alexander operator is in $\mathcal{N}(\beta)$. \square

Theorem 2.5. Let $f_i \in \mathcal{N}(\beta_i)$ for each $i = 1, 2, 3, \dots, n$, with $\beta_i > 1$. Then $F_{\alpha_1, \dots, \alpha_n}(z) \in \mathcal{N}(\rho)$ with $\rho = 1 + \sum_{i=1}^n \alpha_i(\beta_i - 1)$ and $\alpha_i > 0$, ($i = 1, 2, 3, \dots, n$).

Proof. After some calculi, we have

$$\frac{zF''_{\alpha_1, \dots, \alpha_n}(z)}{F'_{\alpha_1, \dots, \alpha_n}(z)} = \alpha_1 \frac{zf''_1(z)}{f'_1(z)} + \dots + \alpha_n \frac{zf''_n(z)}{f'_n(z)} \quad (2.6)$$

that is equivalent to

$$\frac{zF''_{\alpha_1, \dots, \alpha_n}(z)}{F'_{\alpha_1, \dots, \alpha_n}(z)} + 1 = \alpha_1 \left(\frac{zf''_1(z)}{f'_1(z)} + 1 \right) + \dots + \alpha_n \left(\frac{zf''_n(z)}{f'_n(z)} + 1 \right) - \sum_{i=1}^n \alpha_i + 1. \quad (2.7)$$

Since $f_i \in \mathcal{N}(\beta_i)$, for all $i \in \{1, \dots, n\}$, we have

$$\operatorname{Re}\left(\frac{zf''_n(z)}{f'_n(z)} + 1\right) < \beta_i. \quad (2.8)$$

So we obtain

$$\operatorname{Re}\left(\frac{zF''_{\alpha_1, \dots, \alpha_n}(z)}{F'_{\alpha_1, \dots, \alpha_n}(z)} + 1\right) < \sum_{i=1}^n \alpha_i \beta_i - \sum_{i=1}^n \alpha_i + 1 = \sum_{i=1}^n \alpha_i (\beta_i - 1) + 1 \quad (2.9)$$

which implies that $F_{\alpha_1, \dots, \alpha_n} \in \mathcal{N}(\rho)$, where $\rho = 1 + \sum_{i=1}^n \alpha_i(\beta_i - 1)$. \square

Corollary 2.6. Let $f_i \in \mathcal{N}(\beta)$ for each $i = 1, 2, 3, \dots, n$ with $\beta > 1$. Then $F_{\alpha_1, \dots, \alpha_n}(z) \in \mathcal{N}(\eta)$ with $\eta = 1 + \sum_{i=1}^n \alpha_i(\beta - 1)$ and $\alpha_i > 0$, ($i = 1, 2, 3, \dots, n$).

Proof. In Theorem 2.5, we consider $\beta_1 = \beta_2 = \dots = \beta_n = \beta$. \square

Corollary 2.7. Let $f \in \mathcal{N}(\beta)$ with $\beta > 1$. Then the integral operator

$$F_\alpha(z) = \int_0^z [f'(t)]^\alpha dt \quad (2.10)$$

is in the class $\mathcal{N}(\alpha(\beta - 1) + 1)$ and $\alpha > 0$.

Proof. We have

$$\frac{zF''_{\alpha}(z)}{F'_{\alpha}(z)} = \alpha \frac{zf''(z)}{f'(z)}. \quad (2.11)$$

From (2.11) we have

$$\operatorname{Re}\left(\frac{zF''_{\alpha}(z)}{F'_{\alpha}(z)} + 1\right) = \alpha \operatorname{Re}\left(\frac{zf''(z)}{f'(z)} + 1\right) + 1 - \alpha < \alpha\beta + 1 - \alpha = \alpha(\beta - 1) + 1. \quad (2.12)$$

So the relation (2.12) implies that the operator F_{α} is in $\mathcal{N}(\alpha(\beta - 1) + 1)$. \square

Example 2.8. Let $f(z) = (1/(2\beta - 1))\{1 - (1 - z)^{2\beta-1}\} \in \mathcal{N}(\beta)$. After some calculi, we obtain that

$$F_{\alpha}(z) = \int_0^z [f'(t)]^{\alpha} dt = \frac{1}{2\alpha(1 - \beta) - 1} (1 - z)^{2\alpha(\beta-1)+1} \in \mathcal{N}(\alpha(\beta - 1) + 1). \quad (2.13)$$

Acknowledgment

The paper is supported by Grant no. 2-CEX 06-11-10/25.07.2006.

References

- [1] B. A. Uralegaddi, M. D. Ganigi, and S. M. Sarangi, "Univalent functions with positive coefficients," *Tamkang Journal of Mathematics*, vol. 25, no. 3, pp. 225–230, 1994.
- [2] S. Owa and H. M. Srivastava, "Some generalized convolution properties associated with certain subclasses of analytic functions," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 3, no. 3, Article ID 42, 13 pages, 2002.
- [3] D. Breaz and N. Breaz, "Two integral operators," *Studia Universitatis Babeş-Bolyai, Mathematica*, vol. 47, no. 3, pp. 13–19, 2002.
- [4] D. Breaz, S. Owa, and N. Breaz, "A new integral univalent operator," in press.