

Research Article

A New System of Nonlinear Fuzzy Variational Inclusions Involving (A, η) -Accretive Mappings in Uniformly Smooth Banach Spaces

M. Alimohammady,^{1,2} J. Balooee,¹ Y. J. Cho,³ and M. Roohi¹

¹ Department of Mathematics, Faculty of Basic Sciences, University of Mazandaran, Babolsar 47416-1468, Iran

² Department of Mathematics, King's College London Strand, London WC2R 2LS, UK

³ Department of Mathematics Education and the RINS, Gyeongsang National University, Chinju 660-701, South Korea

Correspondence should be addressed to Y. J. Cho, yjcho@gnu.ac.kr

Received 4 October 2009; Accepted 4 November 2009

Recommended by Charles E. Chidume

A new system of nonlinear fuzzy variational inclusions involving (A, η) -accretive mappings in uniformly smooth Banach spaces is introduced and studied many fuzzy variational and variational inequality (inclusion) problems as special cases of this system. By using the resolvent operator technique associated with (A, η) -accretive operator due to Lan et al. and Nadler's fixed points theorem for set-valued mappings, an existence theorem of solutions for this system of fuzzy variational inclusions is proved. We also construct some new iterative algorithms for the solutions of this system of nonlinear fuzzy variational inclusions in uniformly smooth Banach spaces and discuss the convergence of the sequences generated by the algorithms in uniformly smooth Banach spaces. Our results extend, improve, and unify many known results in the recent literatures.

Copyright © 2009 M. Alimohammady et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Variational inequality was initially studied by Stampacchia [1] in 1964. In order to study many kinds of problems arising in industrial, physical, regional, economical, social, pure, and applied sciences, the classical variational inequality problems have been extended and generalized in many directions. Among these generalizations, variational inclusion introduced and studied by Hassouni and Moudafi [2] is of interest and importance. It provides us with a unified, natural, novel innovative, and general technique to study a wide class of the problems arising in different branches of mathematical and engineering sciences (see, e.g., [3–7]).

Next, the development of variational inequality is to design efficient iterative algorithms to compute approximate solutions for variational inequalities and their generalizations. Up to now, many authors have presented implementable and significant numerical methods such as projection method, and its variant forms, linear approximation, descent method, Newton's method and the method based on the auxiliary principle technique. In particular, the method based on the resolvent operator technique is a generalization of the projection method and has been widely used to solve variational inclusions.

Some new and interesting problems, which are called the *systems of variational inequality problems*, were introduced and studied. Pang [8], Cohen and Chaplais [9], Bianchi [10], and Ansari and Yao [11] considered some systems of scalar variational inequalities and Pang showed that the traffic equilibrium problem, the spatial equilibrium problem, the Nash equilibrium, and the general equilibrium programming problems can be modelled as variational inequalities. He decomposed the original variational inequality into a system of variational inequalities which are easy to solve and studied the convergence of such methods. Ansari et al. [12] introduced and studied a system of vector variational inequalities by a fixed point theorem. Allevi et al. [13] considered a system of generalized vector variational inequalities and established some existence results under relative pseudomonotonicity. Kassay and Kolumbán [14] introduced a system of variational inequalities and proved an existence theorem by the Ky Fan lemma. Kassay et al. [15] studied Minty and Stampacchia variational inequality systems with the help of the Kakutani-Fan-Glicksberg fixed point theorem. Peng [16, 17] Peng and Yang [18] introduced a system of quasivariational inequality problems and proved its existence theorem by maximal element theorems. Verma [19–23] introduced and studied some systems of variational inequalities and developed some iterative algorithms for approximating the solution for this system of generalized nonlinear quasivariational inequalities in Hilbert spaces. J. K. Kim and D. S. Kim [24] introduced a new system of generalized nonlinear quasivariational inequalities and obtained some existence and uniqueness results of solution for this system of generalized nonlinear quasivariational inequalities in Hilbert spaces. Cho et al. [25] introduced a new system of nonlinear variational inequalities and proved some existence and uniqueness theorems of solutions for the system of nonlinear variational inequalities in Hilbert spaces. As generalizations of system of variational inequalities, Agarwal et al. [26] introduced a system of generalized nonlinear mixed quasivariational inclusions and investigated the sensitivity of solutions for this system of generalized nonlinear mixed quasivariational inclusions in Hilbert spaces. Kazmi and Bhat [27] introduced a system of nonlinear variational-like inclusions and gave an iterative algorithm for finding its approximate solution. It is known that accretivity of the underlying operator plays indispensable roles in the theory of variational inequality and its generalizations.

In 2001, Huang and Fang [28] were the first to introduce generalized m -accretive mapping and give the definition of the resolvent operator for generalized m -accretive mappings in Banach spaces. They also proved some properties of the resolvent operator for generalized m -accretive mappings in Banach spaces. Subsequently, Fang and Huang [29], Yan et al. [30], Fang et al. [31], Lan et al. [32, 33], Fang and Huang [34], and Peng et al. [35] introduced and investigated many new systems of variational inclusions involving H -monotone operators and (H, η) -monotone operators in Hilbert spaces, generalized m -accretive mappings, H -accretive mappings and (H, η) -accretive mappings in Banach spaces, respectively.

In 2004, Verma in [36, 37] introduced new notions of A -monotone and (A, η) -monotone operators and studied some properties of A -monotone and (A, η) -monotone operators in Hilbert spaces. In [38], Lan et al. first introduced a new concept of (A, η) -accretive mappings, which generalizes the existing monotone or accretive operators and studied some properties of (A, η) -accretive mappings and defined resolvent operators associated with (A, η) -accretive mappings. They also investigated a class of variational inclusions using the resolvent operator associated with (A, η) -accretive mappings. Subsequently, Lan [39], by using the concept of (A, η) -accretive mappings and the new resolvent operator technique associated with (A, η) -accretive mappings, introduced and studied a system of general mixed quasivariational inclusions involving (A, η) -accretive mappings in Banach spaces and constructed a perturbed iterative algorithm with mixed errors for this system of nonlinear (A, η) -accretive variational inclusions in q -uniformly smooth Banach spaces.

On the other hand, the fuzzy set theory introduced by Zadeh [40] has emerged as an interesting and fascinating branch of pure and applied sciences. The application of the fuzzy set theory can be found in many branches of regional, physical, mathematical, and engineering sciences (see [41–45] and the references therein).

In 1989, Chang and Zhu [46] first introduced the classes of variational inequalities for fuzzy mappings. In subsequent years, several classes of variational inequalities, variational inclusions, and complementarity problems for fuzzy mappings were investigated by many authors, in particular, by Chang and Haung [47, 48], Lan et al. [49], Noor [50–52], Noor and Al-said [53], and many others.

Recently, Lan and Verma [54], by using the concept of (A, η) -accretive mappings, the resolvent operator technique associated with (A, η) -accretive mappings, introduced and studied a new class of nonlinear fuzzy variational inclusion systems with (A, η) -accretive mappings in Banach spaces and construct some new iterative algorithms to approximate the solutions of the nonlinear fuzzy variational inclusion systems.

Inspired and motivated by recent research works in these fields, in this paper, we introduce and study a new system of nonlinear fuzzy variational inclusions with (A, η) -accretive mappings in Banach spaces. By using the resolvent operator associated with (A, η) -mappings due to Lan et al. and Nadler's fixed points theorem, we construct some new iterative algorithms for approximating the solutions of this system of nonlinear fuzzy variational inclusions in Banach spaces and prove the existence of solutions and the convergence of the sequences generated by the algorithms in q -uniformly smooth Banach spaces. The results presented in this paper improve and extend the corresponding results of [29–35, 38, 39, 55–60] and many other recent works.

2. Preliminaries

Let X be a real Banach space with dual space X^* , $\langle \cdot, \cdot \rangle$ be the dual pair between X and X^* , 2^X denote the family of all nonempty subsets of X , and let $CB(X)$ denote the family of all nonempty closed bounded subsets of X . The *generalized duality mapping* $J_q : X \rightarrow 2^{X^*}$ is defined by

$$J_q(x) = \left\{ f^* \in X^* : \langle x, f^* \rangle = \|x\|^q, \|f^*\| = \|x\|^{q-1} \right\}, \quad \forall x \in X, \quad (2.1)$$

where $q > 1$ is a constant. In particular, J_2 is the usual normalized duality mapping.

It is known that, in general, $J_q(x) = \|x\|^{q-2}J_2(x)$ for all $x \neq 0$ and J_q is single valued if X^* is strictly convex. In the sequel, we always assume that X is a real Banach space such that J_q is single-valued. If X is a Hilbert space, then J_2 becomes the identity mapping on X .

The *modulus of smoothness* of X is the function $\rho_X : [0, \infty) \rightarrow [0, \infty)$ defined by

$$\rho_X(t) = \sup \left\{ \frac{1}{2}(\|x + y\| + \|x - y\|) - 1 : \|x\| \leq 1, \|y\| \leq t \right\}. \quad (2.2)$$

A Banach space X is said to be *uniformly smooth* if

$$\lim_{t \rightarrow 0} \frac{\rho_X(t)}{t} = 0. \quad (2.3)$$

X is called *q -uniformly smooth* if there exists a constant $c > 0$ such that

$$\rho_X(t) \leq ct^q, \quad \forall q > 1. \quad (2.4)$$

Note that J_q is single-valued if X is uniformly smooth. Concerned with the characteristic inequalities in q -uniformly smooth Banach spaces, Xu [61] proved the following result.

Lemma 2.1. *A real Banach space X is q -uniformly smooth if and only if there exists a constant $c_q > 0$ such that, for all $x, y \in X$,*

$$\|x + y\|^q \leq \|x\|^q + q\langle y, J_q(x) \rangle + c_q \|y\|^q. \quad (2.5)$$

Definition 2.2. A set-valued mapping $T : X \rightarrow 2^X$ is said to be ξ - \widehat{H} -Lipschitz continuous if there exists a constant $\xi > 0$ such that

$$\widehat{H}(T(x), T(y)) \leq \xi \|x - y\|, \quad \forall x, y \in X, \quad (2.6)$$

where $\widehat{H} : 2^X \times 2^X \rightarrow \mathbb{R} \cup \{+\infty\}$ is the Hausdorff pseudo-metric, that is,

$$\widehat{H}(A, B) = \max \left\{ \sup_{x \in A} d(x, B), \sup_{y \in B} d(y, A) \right\}, \quad \forall A, B \in 2^X, \quad (2.7)$$

where $d(u, K) = \inf_{v \in K} \|u - v\|$.

It should be pointed that if domain of \widehat{H} is restricted to closed bounded subsets $CB(X)$, then \widehat{H} is the Hausdorff metric.

Lemma 2.3 (see [62]). *Let (X, d) be a complete metric space and let $T : X \rightarrow CB(X)$ be a set-valued mapping satisfying*

$$\widehat{H}(T(x), T(y)) \leq kd(x, y), \quad \forall x, y \in X, \quad (2.8)$$

where $k \in (0, 1)$ is a constant. Then the mapping T has a fixed point in X .

Lemma 2.4 (see [62]). *Let (X, d) be a complete metric space and let $T : X \rightarrow CB(X)$ be a set-valued mapping. Then for any $\varepsilon > 0$ and any $x, y \in X$, $u \in T(x)$, there exists $v \in T(y)$ such that*

$$d(u, v) \leq (1 + \varepsilon)\widehat{H}(T(x), T(y)). \quad (2.9)$$

Definition 2.5. Let X be a q -uniformly smooth Banach space, $T, A : X \rightarrow X$ and let $\eta : X \times X \rightarrow X$ be single-valued mappings.

(i) T is said to be *accretive* if

$$\langle T(x) - T(y), J_q(x - y) \rangle \geq 0, \quad \forall x, y \in X; \quad (2.10)$$

(ii) T is said to be *strictly accretive* if T is accretive and

$$\langle T(x) - T(y), J_q(x - y) \rangle = 0 \quad (2.11)$$

if and only if $x = y$;

(iii) T is said to be *r -strongly accretive* if there exists a constant $r > 0$ such that

$$\langle T(x) - T(y), J_q(x - y) \rangle \geq r\|x - y\|^q, \quad \forall x, y \in X; \quad (2.12)$$

(iv) T is said to be *m -relaxed accretive* if there exists a constant $m > 0$ such that

$$\langle T(x) - T(y), J_q(x - y) \rangle \geq -m\|x - y\|^q, \quad \forall x, y \in X; \quad (2.13)$$

(v) T is said to be *(ζ, ς) -relaxed cocoercive* if there exist constants $\zeta, \varsigma > 0$ such that

$$\langle T(x) - T(y), J_q(x - y) \rangle \geq -\zeta\|T(x) - T(y)\|^q + \varsigma\|x - y\|^q, \quad \forall x, y \in X; \quad (2.14)$$

(vi) T is said to be *γ -Lipschitz continuous* if there exists a constant $\gamma > 0$ such that

$$\|T(x) - T(y)\| \leq \gamma\|x - y\|, \quad \forall x, y \in X; \quad (2.15)$$

(vii) η is said to be *τ -Lipschitz continuous* if there exists a constant τ such that

$$\|\eta(x, y)\| \leq \tau\|x - y\|, \quad \forall x, y \in X; \quad (2.16)$$

(viii) $\eta(\cdot, \cdot)$ is said to be ϵ -Lipschitz continuous in the first variable if there exists a constant $\epsilon > 0$ such that

$$\|\eta(x, u) - \eta(y, u)\| \leq \epsilon \|x - y\|, \quad \forall x, y, u \in X; \quad (2.17)$$

(ix) $\eta(\cdot, u)$ is said to be (ρ, ξ) -relaxed cocoercive with respect to A if there exist constants $\rho, \xi > 0$ such that

$$\langle \eta(x, u) - \eta(y, u), J_q(A(x) - A(y)) \rangle \geq -\rho \|\eta(x, u) - \eta(y, u)\|^q + \xi \|x - y\|^q, \quad \forall x, y, u \in X. \quad (2.18)$$

In a similar way to (viii) and (ix), we can define the Lipschitz continuity of the mapping $\eta(\cdot, \cdot)$ in the second variable and relaxed cocoercivity of $\eta(u, \cdot)$ with respect to A .

Definition 2.6. Let X be a q -uniformly smooth Banach space, $\eta : X \times X \rightarrow X$ and let $H, A : X \rightarrow X$ be three single-valued mappings. Set-valued mapping $M : X \rightarrow 2^X$ is said to be

(i) *accretive* if

$$\langle u - v, J_q(x - y) \rangle \geq 0, \quad \forall x, y \in X, u \in Mx, v \in My; \quad (2.19)$$

(ii) η -*accretive* if

$$\langle u - v, J_q(\eta(x, y)) \rangle \geq 0, \quad \forall x, y \in X, u \in Mx, v \in My; \quad (2.20)$$

(iii) *strictly η -accretive* if M is η -accretive and the equality holds if and only if $x = y$;

(iv) r -*strongly η -accretive* if there exists a constant $r > 0$ such that

$$\langle u - v, J_q(\eta(x, y)) \rangle \geq r \|x - y\|^q, \quad \forall x, y \in X, u \in Mx, v \in My; \quad (2.21)$$

(v) α -*relaxed η -accretive* if there exists a constant $\alpha > 0$ such that

$$\langle u - v, J_q(\eta(x, y)) \rangle \geq -\alpha \|x - y\|^q, \quad \forall x, y \in X, u \in Mx, v \in My; \quad (2.22)$$

(vi) m -*accretive* if M is accretive and $(I + \lambda M)(X) = X$ for all $\lambda > 0$, where I denotes the identity operator on X ;

(vii) *generalized m -accretive* if M is η -accretive and $(I + \lambda M)(X) = X$ for all $\lambda > 0$;

(viii) H -*accretive* if M is accretive and $(H + \lambda M)(X) = X$ for all $\lambda > 0$;

(ix) (H, η) -*accretive* if M is η -accretive and $(H + \lambda M)(X) = X$ for all $\lambda > 0$.

Remark 2.7. The following should be noticed.

(1) The class of generalized m -accretive operators was first introduced by Huang and Fang [28] and includes that of m -accretive operators as a special case. The class of

H -accretive operators was first introduced and studied by Fang and Huang [63] and also includes that of m -accretive operators as a special case.

- (2) When $X = \mathcal{H}$ is a Hilbert space, (i)–(ix) of Definition 2.6 reduce to the definitions of monotone operators, η -monotone operators, strictly η -monotone operators, strongly η -monotone operators, relaxed η -monotone operators, maximal monotone operators, maximal η -monotone operators, H -monotone operators, and (H, η) -monotone operators, respectively.

Definition 2.8. Let $A : X \rightarrow X$, $\eta : X \times X \rightarrow X$ be two single-valued mappings and let $M : X \rightarrow 2^X$ be a set-valued mapping. Then M is said to be (A, η) -accretive with constant m if M is m -relaxed η -accretive and $(A + \lambda M)(X) = X$ for all $\lambda > 0$.

Remark 2.9. For appropriate and suitable choices of m , A , η , and the space X , it is easy to see that Definition 2.8 includes a number of definitions of monotone operators and accretive operators (see [38]).

In [38], Lan et al. showed that $(A + \rho M)^{-1}$ is a single-valued operator if $M : X \rightarrow 2^X$ is an (A, η) -accretive mapping and $A : X \rightarrow X$ an r -strongly η -accretive mapping. Based on this fact, we can define the resolvent operator $R_{M, \rho}^{\eta, A}$ associated with an (A, η) -accretive mapping M as follows.

Definition 2.10. Let $A : X \rightarrow X$ be a strictly η -accretive mapping and let $M : X \rightarrow 2^X$ be an (A, η) -accretive mapping. The resolvent operator $R_{M, \rho}^{\eta, A} : X \rightarrow X$ associated with A and M is defined by

$$R_{M, \rho}^{\eta, A}(x) = (A + \rho M)^{-1}(x), \quad \forall x \in X. \quad (2.23)$$

Proposition 2.11 (see [38]). Let X be a q -uniformly smooth Banach space, let $\eta : X \times X \rightarrow X$ be τ -Lipschitz continuous, let $A : X \rightarrow X$ be a r -strongly η -accretive mapping and let $M : X \rightarrow 2^X$ be an (A, η) -accretive mapping with constant m . Then the resolvent operator $R_{M, \rho}^{\eta, A} : X \rightarrow X$ is $(\tau^{q-1}/(r - \rho m))$ -Lipschitz continuous, that is,

$$\|R_{M, \rho}^{\eta, A}(x) - R_{M, \rho}^{\eta, A}(y)\| \leq \frac{\tau^{q-1}}{r - \rho m} \|x - y\|, \quad \forall x, y \in X, \quad (2.24)$$

where $\rho \in (0, r/m)$ is a constant.

In what follows, we denote the collection of all fuzzy sets on X by $\mathfrak{F}(X) = \{A \mid A : X \rightarrow [0, 1]\}$. A mapping \mathcal{S} from X to $\mathfrak{F}(X)$ is called a *fuzzy mapping*. If $\mathcal{S} : X \rightarrow \mathfrak{F}(X)$ is a fuzzy mapping, then the set $\mathcal{S}(x)$ for any $x \in X$ is a fuzzy set on $\mathfrak{F}(X)$ (in the sequel we denote $\mathcal{S}(x)$ by \mathcal{S}_x) and $\mathcal{S}_x(y)$ for any $y \in X$ is the degree of membership of y in \mathcal{S}_x . For any $A \in \mathfrak{F}(X)$ and $\alpha \in [0, 1]$, the set

$$(A)_\alpha = \{x \in X : A(x) \geq \alpha\} \quad (2.25)$$

is called a α -cut set of A .

A fuzzy mapping $\mathcal{S} : X \rightarrow \mathfrak{F}(X)$ is said to satisfy the condition $(*)$ if there exists a function $a : X \rightarrow [0, 1]$ such that for each $x \in X$ the set

$$(\mathcal{S}_x)_{a(x)} := \{y \in X : \mathcal{S}_x(y) \geq a(x)\} \quad (2.26)$$

is a nonempty closed and bounded subset of X , that is, $(\mathcal{S}_x)_{a(x)} \in CB(X)$.

By using the fuzzy mapping \mathcal{S} satisfying the condition $(*)$ with corresponding function $a : X \rightarrow [0, 1]$, we can define a set-valued mapping S as follows:

$$S : X \rightarrow CB(X), \quad x \mapsto (\mathcal{S}_x)_{a(x)}. \quad (2.27)$$

In the sequel, S, T, L, D, G, W , and K are called *the set-valued mappings induced by the fuzzy mappings $\mathcal{S}, \mathcal{T}, \mathcal{L}, \mathcal{D}, \mathcal{G}, \mathcal{W}$, and \mathcal{K}* , respectively.

3. A New System of Fuzzy Variational Inclusions

In this section, we introduce some systems of fuzzy variational inclusions q -uniformly smooth Banach spaces X and their relations.

Let X_1 be a q_1 -uniformly smooth Banach space with $q_1 > 1$, let X_2 be a q_2 -uniformly smooth Banach space with $q_2 > 1$, let $E, P : X_1 \times X_2 \rightarrow X_1, F, Q : X_1 \times X_2 \rightarrow X_2, A_1 : X_1 \rightarrow X_1, A_2 : X_2 \rightarrow X_2, f, p, l : X_1 \rightarrow X_1, g, h, k : X_2 \rightarrow X_2, \eta_1 : X_1 \times X_1 \rightarrow X_1, \eta_2 : X_2 \times X_2 \rightarrow X_2$ be single-valued mappings, and let $\mathcal{S}, \mathcal{T}, \mathcal{L}, \mathcal{D} : X_1 \rightarrow \mathfrak{F}(X_1)$ and $\mathcal{G}, \mathcal{W}, \mathcal{K} : X_2 \rightarrow \mathfrak{F}(X_2)$ be fuzzy mappings. Further, suppose that $M : X_1 \times X_1 \rightarrow 2^{X_1}$ and $N : X_2 \times X_2 \rightarrow 2^{X_2}$ are any nonlinear operators such that for all $z \in X_1, M(\cdot, z) : X_1 \rightarrow 2^{X_1}$ is an (A_1, η_1) -accretive with $f(x) - y \in \text{dom}(M(\cdot, z))$ for all $x, y \in X_1$ and for all $t \in X_2, N(\cdot, t) : X_2 \rightarrow 2^{X_2}$ is an (A_2, η_2) -accretive with $g(u) \in \text{dom}(N(\cdot, t))$ for all $u \in X_2$. Now, for given mappings $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d} : X_1 \rightarrow [0, 1]$ and $\tilde{e}, \tilde{f}, \tilde{g} : X_2 \rightarrow [0, 1]$, we consider the following system.

System 3.1. For any given $a \in X_1, b \in X_2, \lambda_1 > 0, \lambda_2 > 0$, our problem is as follows:

Find $x, z, u, v, m \in X_1$ and $y, w, t, s \in X_2$ such that $\mathcal{S}_x(u) \geq \tilde{a}(x), \mathcal{T}_x(v) \geq \tilde{b}(x), \mathcal{L}_x(z) \geq \tilde{c}(x), \mathcal{D}_x(m) \geq \tilde{d}(x), \mathcal{G}_y(w) \geq \tilde{e}(y), \mathcal{W}_y(t) \geq \tilde{f}(y), \mathcal{K}_y(s) \geq \tilde{g}(y)$, and

$$\begin{cases} a \in E(p(x), w) + P(l(z), t) + \lambda_1 M(f(x) - v, x), \\ b \in F(u, h(y)) + Q(m, k(s)) + \lambda_2 N(g(y), y). \end{cases} \quad (3.1)$$

This system is called a *system of nonlinear fuzzy variational inclusions involving (A, η) -accretive mappings* in uniformly smooth Banach spaces.

Remark 3.2. For appropriate and suitable choices of $X_1, X_2, q_1, q_2, E, P, F, Q, A_1, A_2, f, g, h, k, l, p, \eta_1, \eta_2, \mathcal{S}, \mathcal{T}, \mathcal{L}, \mathcal{D}, \mathcal{G}, \mathcal{W}, \mathcal{K}, M, N, \tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}, \tilde{e}, \tilde{f}$, and \tilde{g} one can obtain many known and new classes of (fuzzy) variational inequalities and (fuzzy) variational inclusions as special cases of System 3.1.

Now, we consider some special cases of System 3.1.

System 3.3. Let $S, T, L, D : X_1 \rightarrow CB(X_1)$ and $G, W, K : X_2 \rightarrow CB(X_2)$ be classical set-valued mappings and let $M, N, f, g, E, P, F, Q, p, l, h, k$ be the mappings as in System 3.1. Now, by using $S, T, L, D, G, W,$ and $K,$ we define fuzzy mappings $\mathcal{S}, \mathcal{T}, \mathcal{L}, \mathfrak{D} : X_1 \rightarrow 2^{X_1}$ and $\mathcal{G}, \mathcal{W}, \mathcal{K} : X_2 \rightarrow 2^{X_2}$ as follows:

$$\begin{aligned} \mathcal{S}_x &= \chi_{S(x)}, & \mathcal{T}_x &= \chi_{T(x)}, & \mathcal{L}_x &= \chi_{L(x)} \\ \mathfrak{D}_x &= \chi_{D(x)}, & \mathcal{G}_x &= \chi_{G(x)}, & \mathcal{W}_x &= \chi_{W(x)}, & \mathcal{K}_x &= \chi_{K(x)}, \end{aligned} \tag{3.2}$$

where $\chi_{S(x)}, \chi_{T(x)}, \chi_{L(x)}, \chi_{D(x)}, \chi_{G(x)}, \chi_{W(x)},$ and $\chi_{K(x)}$ are the characteristic functions of the sets $S(x), T(x), L(x), D(x), G(x), W(x),$ and $K(x),$ respectively.

It is easy to see that $\mathcal{S}, \mathcal{T}, \mathcal{L},$ and \mathfrak{D} are fuzzy mappings satisfying the condition (*) with constant functions $\tilde{a}(x) = 1, \tilde{b}(x) = 1, \tilde{c}(x) = 1, \tilde{d}(x) = 1$ for all $x \in X_1,$ respectively, and $\mathcal{G}, \mathcal{W},$ and \mathcal{K} are fuzzy mappings satisfying the condition (*) with constant functions $\tilde{e}(y) = 1, \tilde{f}(y) = 1, \tilde{g}(y) = 1$ for all $y \in X_2,$ respectively. Also

$$\begin{aligned} (\mathcal{S})_{\tilde{a}(x)} &= (\chi_{S(x)})_1 = \{r \in X_1 : \chi_{S(x)}(r) = 1\} = S(x), \\ (\mathcal{T})_{\tilde{b}(x)} &= (\chi_{T(x)})_1 = \{r \in X_1 : \chi_{T(x)}(r) = 1\} = T(x), \\ (\mathcal{L})_{\tilde{c}(x)} &= (\chi_{L(x)})_1 = \{r \in X_1 : \chi_{L(x)}(r) = 1\} = L(x), \\ (\mathfrak{D})_{\tilde{d}(x)} &= (\chi_{D(x)})_1 = \{r \in X_1 : \chi_{D(x)}(r) = 1\} = D(x), \\ (\mathcal{G})_{\tilde{e}(y)} &= (\chi_{G(y)})_1 = \{t \in X_2 : \chi_{G(y)}(t) = 1\} = G(y), \\ (\mathcal{W})_{\tilde{f}(y)} &= (\chi_{W(y)})_1 = \{t \in X_2 : \chi_{W(y)}(t) = 1\} = W(y), \\ (\mathcal{K})_{\tilde{g}(y)} &= (\chi_{K(y)})_1 = \{t \in X_2 : \chi_{K(y)}(t) = 1\} = K(y). \end{aligned} \tag{3.3}$$

Then System 3.1 is equivalent to the following:

Find $x, z, u, v, m \in X_1, y, w, t, s \in X_2$ such that $u \in S(x), v \in T(x), z \in L(x), m \in D(x), w \in G(y), t \in W(y), s \in K(y),$ and

$$\begin{cases} a \in E(p(x), w) + P(l(z), t) + \lambda_1 M(f(x) - v, x), \\ b \in F(u, h(y)) + Q(m, k(s)) + \lambda_2 N(g(y), y). \end{cases} \tag{3.4}$$

System 3.3 is called a *system of nonlinear set-valued variational inclusions with (A, η) -accretive mappings.*

System 3.4. If $T : X_1 \rightarrow X_1$ is a single-valued mapping, then System 3.3 collapses to the following system of nonlinear variational inclusions:

Find $x, z, u, m \in X_1, y, w, t, s \in X_2$ such that $u \in S(x), z \in L(x), m \in D(x), w \in G(y), t \in W(y), s \in K(y),$ and

$$\begin{cases} a \in E(p(x), w) + P(l(z), t) + \lambda_1 M(f(x) - T(x), x), \\ b \in F(u, h(y)) + Q(m, k(s)) + \lambda_2 N(g(y), y). \end{cases} \tag{3.5}$$

System 3.5. If $X_i = \mathcal{H}_i$ ($i = 1, 2$) are two Hilbert spaces, $S : \mathcal{H}_1 \rightarrow \mathcal{H}_1$ and $G : \mathcal{H}_2 \rightarrow \mathcal{H}_2$ are two single-valued mappings, $p = h = l = k \equiv I$ (the identity mapping), $\lambda_1 = \lambda_2 = 1$, $T \equiv 0$ (the zero mapping), $a = b = 0$, $M(x, y) = M(x)$ for all $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_1$, $N(x, y) = N(x)$ for all $(x, y) \in \mathcal{H}_2 \times \mathcal{H}_2$, then System 3.4 reduces to the following system:

Find (x, y, z, m, t, s) such that $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$, $z \in L(x)$, $m \in D(x)$, $t \in W(y)$, $s \in K(y)$, and

$$\begin{cases} 0 \in E(x, y) + P(z, t) + M(f(x)), \\ 0 \in F(x, y) + Q(m, s) + N(g(y)). \end{cases} \quad (3.6)$$

System 3.5 was introduced and studied by Peng and Zhu in [59].

System 3.6. If $a = b = 0$, $\lambda_1 = \lambda_2 = 1$, and $P = Q \equiv 0$, then System 3.3 can be replaced by the following:

Find $u \in S(x)$, $v \in T(x)$ and $w \in G(y)$ such that

$$\begin{cases} 0 \in E(p(x), w) + M(f(x) - v, x), \\ 0 \in F(u, h(y)) + N(g(y), y), \end{cases} \quad (3.7)$$

which is studied by Lan and Verma [54].

System 3.7. If $T : X_1 \rightarrow X_1$ is a single-valued mapping, then System 3.6 collapses to the following system of nonlinear variational inclusions:

Find $x, u \in X_1$, $y, w \in X_2$ such that $u \in S(x)$, $w \in G(y)$ and

$$\begin{cases} 0 \in E(p(x), w) + M(f(x) - T(x), x), \\ 0 \in F(u, h(y)) + N(g(y), y), \end{cases} \quad (3.8)$$

which is studied by Lan and Verma [54].

System 3.8. If $p = h \equiv I$, $T \equiv 0$, $M(x, y) = M(x)$ for all $(x, y) \in X_1 \times X_1$, $N(x, y) = N(x)$ for all $(x, y) \in X_2 \times X_2$, $S : X_1 \rightarrow X_1$, and $G : X_2 \rightarrow X_2$ are identity mappings, then System 3.7 reduces to the following system:

Find $(x, y) \in X_1 \times X_2$ such that

$$\begin{cases} 0 \in E(x, G(y)) + M(f(x)), \\ 0 \in F(S(x), y) + N(g(y)). \end{cases} \quad (3.9)$$

System 3.8 is investigated by Jin [55] when S and G are the identity mappings.

System 3.9. If $f - T = g \equiv I$, $P = Q \equiv 0$, $\lambda_1 = \lambda_2 = 1$, $S : X_1 \rightarrow X_1$, and $G : X_2 \rightarrow X_2$ are two single-valued mappings, then System 3.4 is equivalent to the following:

Find $(x, y) \in X_1 \times X_2$ such that

$$\begin{cases} a \in E(p(x), G(y)) + M(x, x), \\ b \in F(S(x), h(y)) + N(y, y), \end{cases} \quad (3.10)$$

which is introduced and studied by Lan [39] when S and G are identity mappings.

System 3.10. When $p = h = S = G \equiv I$, $a = b = 0$, System 3.9 can be replaced to the following:

Find $(x, y) \in X_1 \times X_2$ such that

$$\begin{cases} 0 \in E(x, y) + M(x, x), \\ 0 \in F(x, y) + N(y, y), \end{cases} \quad (3.11)$$

which is studied by Jin [56].

System 3.11. If $X_i = \mathcal{H}_i$ ($i = 1, 2$) is two Hilbert spaces, $M(x, y) = M(x)$ for all $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_1$ and $N(x, y) = N(x)$ for all $(x, y) \in \mathcal{H}_2 \times \mathcal{H}_2$, then System 3.7 reduces to the following generalized system of set-valued variational inclusions:

Find $x, u \in \mathcal{H}_1, y, w \in \mathcal{H}_2$ such that $u \in S(x), w \in G(y)$, and

$$\begin{cases} 0 \in E(p(x), w) + M(f(x) - T(x)), \\ 0 \in F(u, h(y)) + N(g(y)), \end{cases} \quad (3.12)$$

which is studied by Lan et al. [57] when M, N are A -monotone mappings and there exists single-valued mapping $\psi : \mathcal{H}_1 \rightarrow \mathcal{H}_1$ such that $\psi(x) = f(x) - T(x)$ for all $x \in \mathcal{H}_1$.

System 3.12. If $g = p = h = f - T \equiv I$, then System 3.11 collapses to the following system of nonlinear variational inclusions:

Find $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2, u \in S(x), w \in G(y)$ such that

$$\begin{cases} 0 \in E(x, w) + M(x), \\ 0 \in F(u, y) + N(y), \end{cases} \quad (3.13)$$

which is considered by Huang and Fang [28].

System 3.13. If $S : \mathcal{H}_1 \rightarrow \mathcal{H}_1$ and $G : \mathcal{H}_2 \rightarrow \mathcal{H}_2$ are two single-valued mappings, then System 3.12 is equivalent to the following:

Find $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$ such that

$$\begin{cases} 0 \in E(x, G(y)) + M(x), \\ 0 \in F(S(x), y) + N(y), \end{cases} \quad (3.14)$$

which is investigated by Fang et al. [31] and Peng et al. [35], Fang and Huang [34], and Verma [36] with $S = G \equiv I$.

System 3.14. If $M(x) = \partial\varphi(x)$ and $N(y) = \partial\phi(y)$ for all $x \in \mathcal{A}_1$ and $y \in \mathcal{A}_2$, where $\varphi : \mathcal{A}_1 \rightarrow \mathbb{R} \cup \{+\infty\}$ and $\phi : \mathcal{A}_2 \rightarrow \mathbb{R} \cup \{+\infty\}$ are two proper, convex, and lower semicontinuous functionals, and $\partial\varphi$ and $\partial\phi$ denote subdifferential operators of φ and ϕ , respectively, then System 3.13 reduces to the following system:

Find $(x, y) \in \mathcal{A}_1 \times \mathcal{A}_2$ such that

$$\begin{cases} \langle E(x, G(y)), s - x \rangle + \varphi(s) - \varphi(x) \geq 0, & \forall s \in \mathcal{A}_1, \\ \langle F(S(x), y), t - y \rangle + \phi(t) - \phi(y) \geq 0, & \forall t \in \mathcal{A}_2, \end{cases} \quad (3.15)$$

which is called a *system of nonlinear mixed variational inequalities*. Some special cases of System 3.7 can be found in [22]. Further, if $S = G \equiv I$, then System 3.7 reduces to the system of nonlinear variational inequalities considered by Cho et al. [25].

System 3.15. If $M(x) = \partial\delta_{K_1}(x)$ and $N(y) = \partial\delta_{K_2}(y)$ for all $x \in K_1$ and $y \in K_2$, where K_1 and K_2 are nonempty closed convex subsets of \mathcal{A}_1 and \mathcal{A}_2 , respectively, and δ_{K_1} and δ_{K_2} denote indicator functions of K_1 and K_2 , respectively, then System 3.14 becomes the following problem:

Find $(x, y) \in K_1 \times K_2$ such that

$$\begin{cases} \langle E(x, G(y)), s - x \rangle \geq 0, & \forall s \in K_1, \\ \langle F(S(x), y), t - y \rangle \geq 0, & \forall t \in K_2, \end{cases} \quad (3.16)$$

which is the just system in [24] when S and G are singlevalued and $S = G \equiv I$.

System 3.16. If $\mathcal{A}_1 = \mathcal{A}_2 = \mathcal{A}$, $K_1 = K_2 = K$, $E(x, G(y)) = \rho_1 G(y) + x - y$, and $F(S(x), y) = \rho_2 S(x) + y - x$ for all $x, y \in \mathcal{A}$, where $\rho_1 > 0$ and $\rho_2 > 0$ are two constants, then System 3.15 is equivalent to the following:

Find an element $(x, y) \in K \times K$ such that

$$\begin{cases} \langle \rho_1 G(y) + x - y, s - x \rangle \geq 0, & \forall s \in K, \\ \langle \rho_2 S(x) + y - x, t - y \rangle \geq 0, & \forall t \in K, \end{cases} \quad (3.17)$$

which is the system of nonlinear variational inequalities considered by Verma [22] with $S = G$.

Remark 3.17. If $x = y$, $S = G$ and $\rho_1 = \rho_2$, then System 3.16 reduces to the following classical nonlinear variational inequality problem:

Find an element $x \in K$ such that

$$\langle S(x), z - x \rangle \geq 0, \quad \forall z \in K. \quad (3.18)$$

4. Existence Theorems

In this section, we prove the existence theorem for solutions of Systems 3.1. For our main results, we have the following lemma which offers a good approach to solve System 3.1.

Lemma 4.1. *Let $X_i, A_i, \eta_i, \lambda_i$ ($i = 1, 2$), $E, F, P, Q, S, \mathcal{T}, \mathcal{L}, \mathfrak{D}, \mathcal{G}, \mathcal{W}, \mathcal{K}, M, N, f, g, h, p, l, k, a$, and b be the same as in System 3.1. Then, for any given $x, z, u, v, m \in X_1$ and $y, w, t, s \in X_2$, $(x, y, z, t, m, s, u, v, w)$ is a solution of System 3.1 if and only if*

$$\begin{aligned} f(x) &= v + R_{M(\cdot, x), \rho_1}^{\eta_1, A_1} \left[A_1(f(x) - v) - \frac{\rho_1}{\lambda_1} (E(p(x), w) + P(l(z), t) - a) \right], \\ g(y) &= R_{N(\cdot, y), \rho_2}^{\eta_2, A_2} \left[A_2(g(y)) - \frac{\rho_2}{\lambda_2} (F(u, h(y)) + Q(m, k(s)) - b) \right], \end{aligned} \tag{4.1}$$

where $\rho_1 > 0$ and $\rho_2 > 0$ are two constants.

Proof. The conclusion follows directly from Definition 2.10 and some simple arguments. \square

From Lemma 4.1, we have the following.

Theorem 4.2. *Let X_1 and X_2 be the same as in Lemma 4.1, let $S, \mathcal{T}, \mathcal{L}, \mathfrak{D} : X_1 \rightarrow \mathfrak{F}(X_1)$ and $\mathcal{G}, \mathcal{W}, \mathcal{K} : X_2 \rightarrow \mathfrak{F}(X_2)$ be fuzzy mappings satisfying the condition (*) with the corresponding functions $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}, \tilde{e}, \tilde{f}$ and \tilde{g} , respectively, $S, T, L, D : X_1 \rightarrow CB(X_1)$, and let $G, W, K : X_2 \rightarrow CB(X_2)$ be ξ - \widehat{H}_1 -Lipschitz continuous, ζ - \widehat{H}_1 -Lipschitz continuous, γ - \widehat{H}_1 -Lipschitz continuous, ϖ - \widehat{H}_1 -Lipschitz continuous, ξ' - \widehat{H}_2 -Lipschitz continuous, ζ' - \widehat{H}_2 -Lipschitz continuous, and γ' - \widehat{H}_2 -Lipschitz continuous, respectively, where \widehat{H}_i is the Hausdorff pseudometric on 2^{X_i} for $i = 1, 2$. Assume that $\eta_i : X_i \times X_i \rightarrow X_i$ is τ_i -Lipschitz continuous, $A_i : X_i \rightarrow X_i$ is r_i -strongly η_i -accretive and β_i -Lipschitz continuous for $i = 1, 2$, $p, l : X_1 \rightarrow X_1$ are δ_1 -Lipschitz continuous, and δ_2 -Lipschitz continuous, respectively, $h, k : X_2 \rightarrow X_2$ are π_1 -Lipschitz continuous and π_2 -Lipschitz continuous, respectively, $f : X_1 \rightarrow X_1$ is (κ, e_1) -relaxed cocoercive, μ -Lipschitz continuous and $g : X_2 \rightarrow X_2$ is (σ, e_2) -relaxed cocoercive, e -Lipschitz continuous. Suppose that $M(\cdot, z) : X_1 \rightarrow 2^{X_1}$ is an (A_1, η_1) -accretive operator with constant m_1 for all $z \in X_1$ and $N(\cdot, t) : X_2 \rightarrow 2^{X_2}$ is an (A_2, η_2) -accretive operator with constant m_2 for all $t \in X_2$, and $E, P : X_1 \times X_2 \rightarrow X_1$ are two single-valued mappings such that $E(\cdot, y)$ and $P(\cdot, y)$ are ν_1 -Lipschitz continuous and ν_2 -Lipschitz continuous in the first variable, respectively, $E(x, \cdot), P(x, \cdot)$ are ι_1 -Lipschitz continuous ι_2 -Lipschitz continuous in the second variable, respectively, for all $(x, y) \in X_1 \times X_2$, and $E(p_1(\cdot), y)$ is (θ_1, s_1) -relaxed cocoercive with respect to f' , where $f' : X_1 \rightarrow X_1$ is defined by $f'(x) = A_1 \circ (f(x) - v) = A_1(f(x) - v)$ for all $x \in X_1$, $\tilde{b} : X_1 \rightarrow [0, 1]$ and $\mathcal{T}_x(v) \geq \tilde{b}(x)$. Further, suppose that $F, Q : X_1 \times X_2 \rightarrow X_2$ are two nonlinear mappings such that $F(\cdot, y), Q(\cdot, y)$ are ρ_1 -Lipschitz continuous and ρ_2 -Lipschitz continuous in the first variable, respectively, $F(x, \cdot)$ and $Q(x, \cdot)$ are ν_1 -Lipschitz continuous and ν_2 -Lipschitz continuous in the second variable, respectively, and $F(x, h(\cdot))$ is (θ_2, s_2) -relaxed cocoercive with respect to g' , where $g' : X_2 \rightarrow X_2$ is defined by $g'(x) = A_2 \circ g(x) = A_2(g(x))$ for all $x \in X_2$.*

In addition, if there exist constants $\rho_1 \in (0, r_1/m_1)$ and $\rho_2 \in (0, r_2/m_2)$ such that

$$\left\| R_{M(\cdot, x), \rho_1}^{\eta_1, A_1}(z) - R_{M(\cdot, y), \rho_1}^{\eta_1, A_1}(z) \right\| \leq \varsigma \|x - y\|, \quad \forall x, y, z \in X_1, \tag{4.2}$$

$$\left\| R_{N(\cdot, x), \rho_2}^{\eta_2, A_2}(z) - R_{N(\cdot, y), \rho_2}^{\eta_2, A_2}(z) \right\| \leq \mathfrak{D} \|x - y\|, \quad \forall x, y, z \in X_2, \tag{4.3}$$

$$\begin{aligned}
& \varsigma + \zeta + \sqrt[q_1]{1 - q_1 e_1 + (c_{q_1} + q_1 \kappa) \mu^{q_1}} < 1, \\
& \vartheta + \sqrt[q_2]{1 - q_2 e_2 + (c_{q_2} + q_2 \sigma) \epsilon^{q_2}} < 1, \\
& \sqrt[q_1]{\beta_1^{q_1} (\mu + \zeta)^{q_1} - q_1 \frac{\rho_1}{\lambda_1} (-\theta_1 \nu_1^{q_1} \delta_1^{q_1} + s_1) + \frac{c_{q_1} \rho_1^{q_1} \nu_1^{q_1} \delta_1^{q_1}}{\lambda_1^{q_1}}} < \frac{\tau_1^{1-q_1} \lambda_1}{\rho_1} (r_1 - \rho_1 m_1) \chi_1 - \nu_2 \delta_2 \gamma, \\
& \sqrt[q_2]{\beta_2^{q_2} \epsilon^{q_2} - q_2 \frac{\rho_2}{\lambda_2} (-\theta_2 \nu_1^{q_2} \pi_1^{q_2} + s_2) + \frac{c_{q_2} \rho_2^{q_2} \nu_1^{q_2} \pi_1^{q_2}}{\lambda_2^{q_2}}} < \frac{\tau_2^{1-q_2} \lambda_2}{\rho_2} (r_2 - \rho_2 m_2) \chi_2 - \nu_2 \pi_2 \gamma',
\end{aligned} \tag{4.4}$$

where

$$\begin{aligned}
\chi_1 &= 1 - \left(\varsigma + \zeta + \sqrt[q_1]{1 - q_1 e_1 + (c_{q_1} + q_1 \kappa) \mu^{q_1}} \right) - \frac{\rho_2 \tau_2^{q_2-1} (\rho_1 \xi + \rho_2 \varpi)}{\lambda_2 (r_2 - \rho_2 m_2)}, \\
\chi_2 &= 1 - \left(\vartheta + \sqrt[q_2]{1 - q_2 e_2 + (c_{q_2} + q_2 \sigma) \epsilon^{q_2}} \right) - \frac{\rho_1 \tau_1^{q_1-1} (l_1 \xi' + l_2 \zeta')}{\lambda_1 (r_1 - \rho_1 m_1)},
\end{aligned} \tag{4.5}$$

λ_1, λ_2 are the same as in System 3.1, and c_{q_1}, c_{q_2} are two constants guaranteed by Lemma 2.1, then System 3.1 admits a solution.

Proof. For any given $\rho_1 > 0$ and $\rho_2 > 0$, define mappings $\Phi_{\rho_1} : X_1 \times X_1 \times X_1 \times X_2 \times X_2 \rightarrow X_1$ and $\Psi_{\rho_2} : X_1 \times X_1 \times X_2 \times X_2 \rightarrow X_2$ as follows:

$$\begin{aligned}
& \Phi_{\rho_1}(x, z, v, t, w) \\
&= x - f(x) + v + R_{M(\cdot, x), \rho_1}^{q_1, A_1} \left[A_1(f(x) - v) - \frac{\rho_1}{\lambda_1} (E(p(x), w) + P(l(z), t) - a) \right], \\
& \Psi_{\rho_2}(u, m, s, y) \\
&= y - g(y) + R_{N(\cdot, y), \rho_2}^{q_2, A_2} \left[A_2(g(y)) - \frac{\rho_2}{\lambda_2} (F(u, h(y)) + Q(m, k(s)) - b) \right]
\end{aligned} \tag{4.6}$$

for all $(x, y, z, t, m, s, u, v, w) \in X_1 \times X_2 \times X_1 \times X_2 \times X_1 \times X_2 \times X_1 \times X_1 \times X_2$, where $a \in X_1$ and $b \in X_2$ are the same as in System 3.1, and let $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d} : X_1 \rightarrow [0, 1]$ and $\tilde{e}, \tilde{f}, \tilde{g} : X_2 \rightarrow [0, 1]$ be mappings such that $\mathcal{S}_x(u) \geq \tilde{a}(x)$, $\mathcal{T}_x(v) \geq \tilde{b}(x)$, $\mathcal{L}_x(z) \geq \tilde{c}(x)$, $\mathfrak{D}_x(m) \geq \tilde{d}(x)$, $\mathcal{G}_y(w) \geq \tilde{e}(y)$, $\mathcal{W}_y(t) \geq \tilde{f}(y)$, and $\mathcal{K}_y(s) \geq \tilde{g}(y)$.

Now, define a norm $\|\cdot\|_*$ on $X_1 \times X_2$ by

$$\|(u, v)\|_* = \|u\| + \|v\|, \quad \forall (u, v) \in X_1 \times X_2. \tag{4.7}$$

It is easy to see that $(X_1 \times X_2, \|\cdot\|_*)$ is a Banach space (see [34]). For any given $\rho_1 > 0$ and $\rho_2 > 0$, define a mapping $Q_{\rho_1, \rho_2} : X_1 \times X_2 \times X_1 \times X_1 \times X_1 \times X_1 \times X_2 \times X_2 \times X_2 \rightarrow X_1 \times X_2$ by

$$Q_{\rho_1, \rho_2}(x, y, z, u, v, m, s, t, w) = (\Phi_{\rho_1}(x, z, v, t, w), \Psi_{\rho_2}(u, m, s, y)) \tag{4.8}$$

for all $(x, y, z, u, v, m, s, t, w) \in X_1 \times X_2 \times X_1 \times X_1 \times X_1 \times X_2 \times X_2 \times X_2$ and let

$$\begin{aligned} \mathfrak{R}_{\rho_1, \rho_2}(x, y) = \{ & Q_{\rho_1, \rho_2}(x, y, z, u, v, m, s, t, w) : \\ & S_x(u) \geq \tilde{a}(x), \mathcal{T}_x(v) \geq \tilde{b}(x), \mathcal{L}_x(z) \geq \tilde{c}(x), \mathfrak{D}_x(m) \geq \tilde{d}(x), \mathcal{G}_y(w) \geq \tilde{e}(y), \\ & \mathcal{W}_y(t) \geq \tilde{f}(y), \mathcal{K}_y(s) \geq \tilde{g}(y), \text{ where } \tilde{a}, \tilde{b}, \tilde{c}, \tilde{d} : X_1 \rightarrow [0, 1], \\ & \tilde{e}, \tilde{f}, \tilde{g} : X_2 \rightarrow [0, 1] \} \end{aligned} \tag{4.9}$$

for all $(x, y) \in X_1 \times X_2$. Then, for any given $(x, y), (x', y') \in X_1 \times X_2$, $\varepsilon > 0$ and $Q_{\rho_1, \rho_2}(x, y, z, u, v, m, s, t, w) \in \mathfrak{R}_{\rho_1, \rho_2}(x, y)$, there exists $(z, u, v, m, s, t, w) \in X_1 \times X_1 \times X_1 \times X_1 \times X_2 \times X_2 \times X_2$ such that

$$\begin{aligned} S_x(u) \geq \tilde{a}(x), \quad \mathcal{T}_x(v) \geq \tilde{b}(x), \quad \mathcal{L}_x(z) \geq \tilde{c}(x), \quad \mathfrak{D}_x(m) \geq \tilde{d}(x), \\ \mathcal{G}_y(w) \geq \tilde{e}(y), \quad \mathcal{W}_y(t) \geq \tilde{f}(y), \quad \mathcal{K}_y(s) \geq \tilde{g}(y), \end{aligned} \tag{4.10}$$

where $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d} : X_1 \rightarrow [0, 1], \tilde{e}, \tilde{f}, \tilde{g} : X_2 \rightarrow [0, 1]$ and (4.6) holds. Since $S_x(u) \geq \tilde{a}(x), \mathcal{T}_x(v) \geq \tilde{b}(x), \mathcal{L}_x(z) \geq \tilde{c}(x), \mathfrak{D}_x(m) \geq \tilde{d}(x), \mathcal{G}_y(w) \geq \tilde{e}(y), \mathcal{W}_y(t) \geq \tilde{f}(y), \mathcal{K}_y(s) \geq \tilde{g}(y)$, that is, $u \in S(x) \in CB(X_1), v \in T(x) \in CB(X_1), z \in L(x) \in CB(X_1), m \in D(x) \in CB(X_1), w \in G(y) \in CB(X_2), t \in W(y) \in CB(X_2), s \in K(y) \in CB(X_2)$, it follows from Lemma 2.4 that there exist $u' \in S(x'), v' \in T(x'), z' \in L(x'), m' \in D(x'), w' \in G(y'), t' \in W(y'), s' \in K(y')$, that is, $S_{x'}(u') \geq \tilde{a}(x'), \mathcal{T}_{x'}(v') \geq \tilde{b}(x'), \mathcal{L}_{x'}(z') \geq \tilde{c}(x'), \mathfrak{D}_{x'}(m') \geq \tilde{d}(x'), \mathcal{G}_{y'}(w') \geq \tilde{e}(y'), \mathcal{W}_{y'}(t') \geq \tilde{f}(y'), \mathcal{K}_{y'}(s') \geq \tilde{g}(y')$ such that

$$\begin{aligned} \|u - u'\| &\leq (1 + \varepsilon)\widehat{H}_1(S(x), S(x')), & \|v - v'\| &\leq (1 + \varepsilon)\widehat{H}_1(T(x), T(x')), \\ \|z - z'\| &\leq (1 + \varepsilon)\widehat{H}_1(L(x), L(x')), & \|m - m'\| &\leq (1 + \varepsilon)\widehat{H}_1(D(x), D(x')), \\ \|w - w'\| &\leq (1 + \varepsilon)\widehat{H}_2(G(y), G(y')), & \|t - t'\| &\leq (1 + \varepsilon)\widehat{H}_2(W(y), W(y')), \\ \|s - s'\| &\leq (1 + \varepsilon)\widehat{H}_2(K(y), K(y')). \end{aligned} \tag{4.11}$$

Letting

$$\begin{aligned} & \Phi_{\rho_1}(x', z', v', t', w') \\ &= x' - f(x') + v' + R_{M(\cdot, x'), \rho_1}^{\eta_1, A_1} \left[A_1(f(x') - v') - \frac{\rho_1}{\lambda_1}(E(p(x'), w') + P(l(z'), t') - a) \right], \\ & \Psi_{\rho_2}(u', m', s', y') \\ &= y' - g(y') + R_{N(\cdot, y'), \rho_2}^{\eta_2, A_2} \left[A_2(g(y')) - \frac{\rho_2}{\lambda_2}(F(u', h(y')) + Q(m', k(s')) - b) \right], \end{aligned} \tag{4.12}$$

we have

$$(\Phi_{\rho_1}(x', z', v', t', w'), \Psi_{\rho_2}(u', m', s', y')) = Q_{\rho_1, \rho_2}(x', y', z', u', v', m', s', t', w'). \quad (4.13)$$

Now, it follows from (4.2) and Proposition 2.11 that

$$\begin{aligned} & \|\Phi_{\rho_1}(x, z, v, t, w) - \Phi_{\rho_1}(x', z', v', t', w')\| \\ & \leq \|x - x' - (f(x) - f(x'))\| + \|v - v'\| \\ & \quad + \left\| R_{M(\cdot, x), \rho_1}^{\eta_1, A_1} \left[A_1(f(x) - v) - \frac{\rho_1}{\lambda_1} (E(p(x), w) + P(l(z), t) - a)) \right] \right. \\ & \quad \left. - R_{M(\cdot, x'), \rho_1}^{\eta_1, A_1} \left[A_1(f(x') - v') - \frac{\rho_1}{\lambda_1} (E(p(x'), w') + P(l(z'), t') - a)) \right] \right\| \\ & \leq \|x - x' - (f(x) - f(x'))\| + \|v - v'\| \\ & \quad + \left\| R_{M(\cdot, x), \rho_1}^{\eta_1, A_1} \left[A_1(f(x) - v) - \frac{\rho_1}{\lambda_1} (E(p(x), w) + P(l(z), t) - a)) \right] \right. \\ & \quad \left. - R_{M(\cdot, x'), \rho_1}^{\eta_1, A_1} \left[A_1(f(x') - v') - \frac{\rho_1}{\lambda_1} (E(p(x'), w') + P(l(z'), t') - a)) \right] \right\| \\ & \quad + \left\| R_{M(\cdot, x), \rho_1}^{\eta_1, A_1} \left[A_1(f(x') - v') - \frac{\rho_1}{\lambda_1} (E(p(x'), w') + P(l(z'), t') - a)) \right] \right. \\ & \quad \left. - R_{M(\cdot, x'), \rho_1}^{\eta_1, A_1} \left[A_1(f(x') - v') - \frac{\rho_1}{\lambda_1} (E(p(x'), w') + P(l(z'), t') - a)) \right] \right\| \\ & \leq \|x - x' - (f(x) - f(x'))\| + \|v - v'\| + \varsigma \|x - x'\| \\ & \quad + \frac{\tau_1^{q_1-1}}{r_1 - \rho_1 m_1} \left\| \left\| A_1(f(x) - v) - \frac{\rho_1}{\lambda_1} (E(p(x), w) + P(l(z), t) - a)) \right. \right. \\ & \quad \left. \left. - \left(A_1(f(x') - v') - \frac{\rho_1}{\lambda_1} (E(p(x'), w') + P(l(z'), t') - a)) \right) \right\| \right\| \\ & \leq \|x - x' - (f(x) - f(x'))\| + \|v - v'\| + \varsigma \|x - x'\| \\ & \quad + \frac{\tau_1^{q_1-1}}{r_1 - \rho_1 m_1} \left\{ \frac{\rho_1}{\lambda_1} [\|E(p(x), w) - E(p(x), w')\| + \|P(l(z), t) - P(l(z), t')\| \right. \\ & \quad \left. + \|P(l(z), t') - P(l(z'), t')\|] \right. \\ & \quad \left. + \left\| A_1(f(x) - v) - A_1(f(x') - v') - \frac{\rho_1}{\lambda_1} (E(p(x), w') - E(p(x'), w')) \right\| \right\}. \quad (4.14) \end{aligned}$$

Thus, by Lemma 2.1, we have

$$\begin{aligned} & \|x - x' - (f(x) - f(x'))\|^{q_1} \\ & \leq \|x - x'\|^{q_1} - q_1 \langle f(x) - f(x'), J_{q_1}(x - x') \rangle + c_{q_1} \|f(x) - f(x')\|^{q_1}. \end{aligned} \quad (4.15)$$

Since f is (κ, e_1) -relaxed cocoercive and μ -Lipschitz continuous, we conclude that

$$\begin{aligned} \|x - x' - (f(x) - f(x'))\|^{q_1} & \leq \|x - x'\|^{q_1} - q_1 e_1 \|x - x'\|^{q_1} + (c_{q_1} + q_1 \kappa) \mu^{q_1} \|x - x'\|^{q_1} \\ & = (1 - q_1 e_1 + (c_{q_1} + q_1 \kappa) \mu^{q_1}) \|x - x'\|^{q_1}. \end{aligned} \quad (4.16)$$

By (4.11) and ζ - \widehat{H}_1 -Lipschitz continuity of T ,

$$\|v - v'\| \leq (1 + \varepsilon) \widehat{H}_1(T(x), T(x')) \leq \zeta(1 + \varepsilon) \|x - x'\|. \quad (4.17)$$

Since $E(x, \cdot)$ is ι_1 -Lipschitz continuous in the second variable and G is ξ' - \widehat{H}_2 -Lipschitz continuous, by (4.11), we have

$$\begin{aligned} \|E(p(x), w) - E(p(x), w')\| & \leq \iota_1 \|w - w'\| \leq \iota_1 (1 + \varepsilon) \widehat{H}_2(G(y), G(y')) \\ & \leq \iota_1 \xi' (1 + \varepsilon) \|y - y'\|. \end{aligned} \quad (4.18)$$

Since $P(x, \cdot)$ is ι_2 -Lipschitz continuous in the second variable, W is ζ' - \widehat{H}_2 -Lipschitz continuous, p_2 is δ_2 -Lipschitz continuous, $P(\cdot, y)$ is ν_2 -Lipschitz continuous in the first variable, and L is γ - \widehat{H}_1 -Lipschitz continuous, using (4.11), we deduce that

$$\begin{aligned} \|P(l(z), t) - P(l(z), t')\| & \leq \iota_2 \|t - t'\| \leq \iota_2 (1 + \varepsilon) \widehat{H}_2(W(y), W(y')) \\ & \leq \iota_2 \zeta' (1 + \varepsilon) \|y - y'\|, \end{aligned} \quad (4.19)$$

and also

$$\begin{aligned} \|P(l(z), t') - P(l(z'), t')\| & \leq \nu_2 \|l(z) - l(z')\| \leq \nu_2 \delta_2 \|z - z'\| \\ & \leq \nu_2 \delta_2 (1 + \varepsilon) \widehat{H}_1(L(x), L(x')) \\ & \leq \nu_2 \delta_2 \gamma (1 + \varepsilon) \|x - x'\|. \end{aligned} \quad (4.20)$$

Again, by Lemma 2.1, it follows that

$$\begin{aligned} & \left\| A_1(f(x) - v) - A_1(f(x') - v') - \frac{\rho_1}{\lambda_1}(E(p(x), w') - E(p(x'), w')) \right\|^{q_1} \\ & \leq \|A_1(f(x) - v) - A_1(f(x') - v')\|^{q_1} - q_1 \frac{\rho_1}{\lambda_1} \\ & \quad \times \langle E(p(x), w') - E(p(x'), w'), J_{q_1}(A_1(f(x) - v) - A_1(f(x') - v')) \rangle \\ & \quad + c_{q_1} \frac{\rho_1^{q_1}}{\lambda_1^{q_1}} \|E(p(x), w') - E(p(x'), w')\|^{q_1}. \end{aligned} \quad (4.21)$$

Since A_1 is β_1 -Lipschitz continuous, f is μ -Lipschitz continuous, and T is ζ -Lipschitz continuous, by (4.11), we get

$$\begin{aligned} \|A_1(f(x) - v) - A_1(f(x') - v')\| & \leq \beta_1 \|f(x) - f(x') - (v - v')\| \\ & \leq \beta_1 (\|f(x) - f(x')\| + \|v - v'\|) \\ & \leq \beta_1 (\mu + \zeta(1 + \varepsilon)) \|x - x'\|. \end{aligned} \quad (4.22)$$

Since $E(p(\cdot), y)$ is (θ_1, s_1) -relaxed cocoercive with respect to f' , where $f'(x) = A_1 \circ (f(x) - v) = A_1(f(x) - v)$, $E(\cdot, y)$ is ν_1 -Lipschitz continuous in the first variable and p is δ_1 -Lipschitz continuous, we have

$$\begin{aligned} & \langle E(p(x), w') - E(p(x'), w'), J_{q_1}(A_1(f(x) - v) - A_1(f(x') - v')) \rangle \\ & \leq -\theta_1 \|E(p(x), w') - E(p(x'), w')\|^{q_1} + s_1 \|x - x'\|^{q_1} \\ & \leq -\theta_1 \nu_1^{q_1} \|p(x) - p(x')\|^{q_1} + s_1 \|x - x'\|^{q_1} \\ & \leq (-\theta_1 \nu_1^{q_1} \delta_1^{q_1} + s_1) \|x - x'\|^{q_1}, \end{aligned} \quad (4.23)$$

and also

$$\|E(p(x), w') - E(p(x'), w')\| \leq \nu_1 \|p(x) - p(x')\| \leq \nu_1 \delta_1 \|x - x'\|. \quad (4.24)$$

Hence, using (4.21)–(4.24), we have

$$\begin{aligned} & \left\| A_1(f(x) - v) - A_1(f(x') - v') - \frac{\rho_1}{\lambda_1}(E(p(x), w') - E(p(x'), w')) \right\|^{q_1} \\ & \leq \left(\beta_1^{q_1} (\mu + \zeta(1 + \varepsilon))^{q_1} - q_1 \frac{\rho_1}{\lambda_1} (-\theta_1 \nu_1^{q_1} \delta_1^{q_1} + s_1) + \frac{c_{q_1} \rho_1^{q_1} \nu_1^{q_1} \delta_1^{q_1}}{\lambda_1^{q_1}} \right) \|x - x'\|^{q_1}, \end{aligned} \quad (4.25)$$

where c_{q_1} is the constant as in Lemma 2.1. Using (4.14)–(4.20), and (4.25), it follows that

$$\|\Phi_{\rho_1}(x, z, v, t, w) - \Phi_{\rho_1}(x', z', v', t', w')\| \leq \varphi_1(\varepsilon) \|x - x'\| + \phi_1(\varepsilon) \|y - y'\|, \quad (4.26)$$

where

$$\begin{aligned}\varphi_1(\varepsilon) &= \zeta + \zeta(1 + \varepsilon) + \sqrt[q_1]{1 - q_1 e_1 + (c_{q_1} + q_1 \kappa) \mu^{q_1}} + \frac{\rho_1 \tau_1^{q_1 - 1} (\nu_2 \delta_2 \gamma (1 + \varepsilon) + \psi_1(\varepsilon))}{\lambda_1 (r_1 - \rho_1 m_1)}, \\ \psi_1(\varepsilon) &= \sqrt[q_1]{\beta_1^{q_1} (\mu + \zeta(1 + \varepsilon))^{q_1} - q_1 \frac{\rho_1}{\lambda_1} (-\theta_1 \nu_1^{q_1} \delta_1^{q_1} + s_1) + \frac{c_{q_1} \rho_1^{q_1} \nu_1^{q_1} \delta_1^{q_1}}{\lambda_1^{q_1}}}, \\ \phi_1(\varepsilon) &= \frac{\rho_1 \tau_1^{q_1 - 1} (l_1 \xi' + l_2 \zeta')(1 + \varepsilon)}{\lambda_1 (r_1 - \rho_1 m_1)}.\end{aligned}\tag{4.27}$$

Similarly, for any $(u, m, s, y), (u', m', s', y') \in X_1 \times X_1 \times X_2 \times X_2$, it follows from (4.3) and Proposition 2.11 that

$$\begin{aligned}& \|\Psi_{\rho_2}(u, m, s, y) - \Psi_{\rho_2}(u', m', s', y')\| \\ & \leq \|y - y' - (g(y) - g(y'))\| \\ & \quad + \left\| R_{N(\cdot, y), \rho_2}^{\eta_2, A_2} \left[A_2(g(y)) - \frac{\rho_2}{\lambda_2} (F(u, h(y)) + Q(m, k(s)) - b) \right] \right. \\ & \quad \left. - R_{N(\cdot, y'), \rho_2}^{\eta_2, A_2} \left[A_2(g(y')) - \frac{\rho_2}{\lambda_2} (F(u', h(y')) + Q(m', k(s')) - b) \right] \right\| \\ & \leq \|y - y' - (g(y) - g(y'))\| \\ & \quad + \left\| R_{N(\cdot, y), \rho_2}^{\eta_2, A_2} \left[A_2(g(y)) - \frac{\rho_2}{\lambda_2} (F(u, h(y)) + Q(m, k(s)) - b) \right] \right. \\ & \quad \left. - R_{N(\cdot, y), \rho_2}^{\eta_2, A_2} \left[A_2(g(y')) - \frac{\rho_2}{\lambda_2} (F(u', h(y')) + Q(m', k(s')) - b) \right] \right\| \\ & \quad + \left\| R_{N(\cdot, y), \rho_2}^{\eta_2, A_2} \left[A_2(g(y')) - \frac{\rho_2}{\lambda_2} (F(u', h(y')) + Q(m', k(s')) - b) \right] \right. \\ & \quad \left. - R_{N(\cdot, y'), \rho_2}^{\eta_2, A_2} \left[A_2(g(y')) - \frac{\rho_2}{\lambda_2} (F(u', h(y')) + Q(m', k(s')) - b) \right] \right\| \\ & \leq \|y - y' - (g(y) - g(y'))\| + \vartheta \|y - y'\| \\ & \quad + \frac{\tau_2^{q_2 - 1}}{r_2 - \rho_2 m_2} \left\| A_2(g(y)) - \frac{\rho_2}{\lambda_2} (F(u, h(y)) + Q(m, k(s)) - b) \right. \\ & \quad \left. - \left(A_2(g(y')) - \frac{\rho_2}{\lambda_2} (F(u', h(y')) + Q(m', k(s')) - b) \right) \right\|\end{aligned}$$

$$\begin{aligned}
&\leq \|y - y' - (g(y) - g(y'))\| + \vartheta \|y - y'\| + \frac{\tau_2^{q_2-1}}{r_2 - \rho_2 m_2} \\
&\quad \times \left\{ \frac{\rho_2}{\lambda_2} (\|F(u, h(y)) - F(u', h(y))\| + \|Q(m, k(s)) - Q(m', k(s))\| \right. \\
&\quad \quad \left. + \|Q(m', k(s)) - Q(m', k(s'))\|) \right. \\
&\quad \left. + \left\| A_2(g(y)) - A_2(g(y')) - \frac{\rho_2}{\lambda_2} (F(u', h(y)) - F(u', h(y'))) \right\| \right\}. \tag{4.28}
\end{aligned}$$

Thus, by Lemma 2.1, we have

$$\begin{aligned}
&\|y - y' - (g(y) - g(y'))\|^{q_2} \\
&\leq \|y - y'\|^{q_2} - q_2 \langle g(y) - g(y'), J_{q_2}(y - y') \rangle + c_{q_2} \|g(y) - g(y')\|^{q_2}. \tag{4.29}
\end{aligned}$$

Since g is (σ, e_2) -relaxed cocoercive and ϵ -Lipschitz continuous, we have

$$\begin{aligned}
\|y - y' - (g(y) - g(y'))\|^{q_2} &\leq \|y - y'\|^{q_2} - q_2 e_2 \|y - y'\|^{q_2} + (c_{q_2} + q_2 \sigma) \epsilon^{q_2} \|y - y'\|^{q_2} \\
&= (1 - q_2 e_2 + (c_{q_2} + q_2 \sigma) \epsilon^{q_2}) \|y - y'\|^{q_2}. \tag{4.30}
\end{aligned}$$

Since $F(\cdot, y)$ is ρ_1 -Lipschitz continuous in the first variable and S is ξ - \widehat{H}_1 -Lipschitz continuous, by (4.11), we obtain

$$\begin{aligned}
\|F(u, h(y)) - F(u', h(y))\| &\leq \rho_1 \|u - u'\| \leq \rho_1 (1 + \epsilon) \widehat{H}_1(S(x), S(x')) \\
&\leq \rho_1 \xi (1 + \epsilon) \|x - x'\|. \tag{4.31}
\end{aligned}$$

Since $Q(x, \cdot)$ is v_2 -Lipschitz continuous in the second variable, k is π_2 -Lipschitz continuous, $Q(\cdot, y)$ is ρ_2 -Lipschitz continuous in the first variable, D is ϖ - \widehat{H}_1 -Lipschitz continuous, and K is γ' - \widehat{H}_2 -Lipschitz continuous, using (4.11), we conclude that

$$\begin{aligned}
\|Q(m, k(s)) - Q(m', k(s))\| &\leq \rho_2 \|m - m'\| \\
&\leq \rho_2 (1 + \epsilon) \widehat{H}_1(D(x), D(x')) \\
&\leq \rho_2 \varpi (1 + \epsilon) \|x - x'\|, \tag{4.32}
\end{aligned}$$

$$\begin{aligned}
\|Q(m', k(s)) - Q(m', k(s'))\| &\leq v_2 \|k(s) - k(s')\| \\
&\leq v_2 \pi_2 \|s - s'\| \\
&\leq v_2 \pi_2 (1 + \epsilon) \widehat{H}_2(K(y), K(y')) \\
&\leq v_2 \pi_2 \gamma' (1 + \epsilon) \|y - y'\|. \tag{4.33}
\end{aligned}$$

Again, by Lemma 2.1, it follows that

$$\begin{aligned} & \left\| A_2(g(y)) - A_2(g(y')) - \frac{\rho_2}{\lambda_2}(F(u', h(y)) - F(u', h(y'))) \right\|^{q_2} \\ & \leq \|A_2(g(y)) - A_2(g(y'))\|^{q_2} \\ & \quad - q_2 \frac{\rho_2}{\lambda_2} \langle F(u', h(y)) - F(u', h(y')), J_{q_2}(A_2(g(y)) - A_2(g(y'))) \rangle \\ & \quad + c_{q_2} \frac{\rho_2^{q_2}}{\lambda_2^{q_2}} \|F(u', h(y)) - F(u', h(y'))\|^{q_2}. \end{aligned} \tag{4.34}$$

Since A_2 is β_2 -Lipschitz continuous and g is ϵ -Lipschitz continuous, we have

$$\|A_2(g(y)) - A_2(g(y'))\| \leq \beta_2 \|g(y) - g(y')\| \leq \beta_2 \epsilon \|y - y'\|. \tag{4.35}$$

Since $F(u, h(\cdot))$ is (θ_2, s_2) -relaxed cocoercive with respect to $g' = A_2 \circ g$, $F(x, \cdot)$ is ν_1 -Lipschitz continuous in the second variable, and h is π_1 -Lipschitz continuous, we get

$$\begin{aligned} & \langle F(u', h(y)) - F(u', h(y')), J_{q_2}(A_2(g(y)) - A_2(g(y'))) \rangle \\ & \leq -\theta_2 \|F(u', h(y)) - F(u', h(y'))\|^{q_2} + s_2 \|y - y'\|^{q_2} \\ & \leq -\theta_2 \nu_1^{q_2} \|h(y) - h(y')\|^{q_2} + s_2 \|y - y'\|^{q_2} \\ & \leq (-\theta_2 \nu_1^{q_2} \pi_1^{q_2} + s_2) \|y - y'\|^{q_2}, \end{aligned} \tag{4.36}$$

$$\begin{aligned} & \|F(u', h(y)) - F(u', h(y'))\| \leq \nu_1 \|h(y) - h(y')\| \\ & \leq \nu_1 \pi_1 \|y - y'\|. \end{aligned} \tag{4.37}$$

Therefore, it follows from (4.34)–(4.37) that

$$\begin{aligned} & \left\| A_2(g(y)) - A_2(g(y')) - \frac{\rho_2}{\lambda_2}(F(u', h(y)) - F(u', h(y'))) \right\|^{q_2} \\ & \leq \left(\beta_2^{q_2} \epsilon^{q_2} - q_2 \frac{\rho_2}{\lambda_2} (-\theta_2 \nu_1^{q_2} \pi_1^{q_2} + s_2) + \frac{c_{q_2} \rho_2^{q_2} \nu_1^{q_2} \pi_1^{q_2}}{\lambda_2^{q_2}} \right) \|y - y'\|^{q_2}, \end{aligned} \tag{4.38}$$

where c_{q_2} is the constant as in Lemma 2.1. From (4.28)–(4.33), and (4.38), it follows that

$$\|\Psi_{\rho_2}(u, m, s, y) - \Psi_{\rho_2}(u', m', s', y')\| \leq \varphi_2(\epsilon) \|x - x'\| + \phi_2(\epsilon) \|y - y'\|, \tag{4.39}$$

where

$$\begin{aligned}\varphi_2(\varepsilon) &= \frac{\rho_2 \tau_2^{q_2-1} (\rho_1 \xi + \rho_2 \varpi)(1 + \varepsilon)}{\lambda_2 (r_2 - \rho_2 m_2)}, \\ \phi_2(\varepsilon) &= \vartheta + \sqrt[q_2]{1 - q_2 e_2 + (c_{q_2} + q_2 \sigma) \varepsilon^{q_2}} + \frac{\rho_2 \tau_2^{q_2-1} (v_2 \pi_2 \gamma' (1 + \varepsilon) + \psi_2)}{\lambda_2 (r_2 - \rho_2 m_2)}, \\ \psi_2 &= \sqrt[q_2]{\beta_2^{q_2} \varepsilon^{q_2} - q_2 \frac{\rho_2}{\lambda_2} (-\theta_2 v_1^{q_2} \pi_1^{q_2} + s_2) + \frac{c_{q_2} \rho_2^{q_2} v_1^{q_2} \pi_1^{q_2}}{\lambda_2^{q_2}}}.\end{aligned}\quad (4.40)$$

It follows from (4.26) and (4.39) that

$$\begin{aligned}\|\Phi_{\rho_1}(x, z, v, t, w) - \Phi_{\rho_1}(x', z', v', t', w')\| + \|\Psi_{\rho_2}(u, m, s, y) - \Psi_{\rho_2}(u', m', s', y')\| \\ \leq \omega(\varepsilon) (\|x - x'\| + \|y - y'\|),\end{aligned}\quad (4.41)$$

where $\omega(\varepsilon) = \max\{\varphi_1(\varepsilon) + \varphi_2(\varepsilon), \phi_1(\varepsilon) + \phi_2(\varepsilon)\}$. Using (4.8) and (4.41), we deduce that

$$\begin{aligned}\|Q_{\rho_1, \rho_2}(x, y, z, u, v, m, s, t, w) - Q_{\rho_1, \rho_2}(x', y', z', u', v', m', s', t', w')\|_* \\ \leq \omega(\varepsilon) \|(x, y) - (x', y')\|_*,\end{aligned}\quad (4.42)$$

that is,

$$\begin{aligned}\sup_{Q_{\rho_1, \rho_2}(x, y, z, u, v, m, s, t, w) \in \mathfrak{R}_{\rho_1, \rho_2}(x, y)} d(Q_{\rho_1, \rho_2}(x, y, z, u, v, m, s, t, w), \mathfrak{R}_{\rho_1, \rho_2}(x', y')) \\ \leq \omega(\varepsilon) \|(x, y) - (x', y')\|_*.\end{aligned}\quad (4.43)$$

Similarly, we have

$$\begin{aligned}\sup_{Q_{\rho_1, \rho_2}(x', y', z', u', v', m', s', t', w') \in \mathfrak{R}_{\rho_1, \rho_2}(x', y')} d(Q_{\rho_1, \rho_2}(x', y', z', u', v', m', s', t', w'), \mathfrak{R}_{\rho_1, \rho_2}(x, y)) \\ \leq \omega(\varepsilon) \|(x, y) - (x', y')\|_*.\end{aligned}\quad (4.44)$$

By (4.43), (4.44), and the definition of Hausdorff pseudo-metric, we have

$$\widehat{H}(\mathfrak{R}_{\rho_1, \rho_2}(x, y), \mathfrak{R}_{\rho_1, \rho_2}(x', y')) \leq \omega(\varepsilon) \|(x, y) - (x', y')\|_*, \quad \forall (x, y), (x', y') \in X_1 \times X_2. \quad (4.45)$$

Letting $\varepsilon \rightarrow 0$, one has

$$\widehat{H}(\mathfrak{R}_{\rho_1, \rho_2}(x, y), \mathfrak{R}_{\rho_1, \rho_2}(x', y')) \leq \omega \|(x, y) - (x', y')\|_*, \quad \forall (x, y), (x', y') \in X_1 \times X_2, \quad (4.46)$$

where

$$\omega = \max\{\varphi_1 + \varphi_2, \phi_1 + \phi_2\}, \tag{4.47}$$

$$\begin{aligned} \varphi_1 &= \varsigma + \zeta + \sqrt[q_1]{1 - q_1 e_1 + (c_{q_1} + q_1 \kappa) \mu^{q_1}} + \frac{\rho_1 \tau_1^{q_1 - 1} (\nu_2 \delta_2 \gamma + \psi_1)}{\lambda_1 (r_1 - \rho_1 m_1)}, \\ \varphi_1 &= \sqrt[q_1]{\beta_1^{q_1} (\mu + \zeta)^{q_1} - q_1 \frac{\rho_1}{\lambda_1} (-\theta_1 \nu_1^{q_1} \delta_1^{q_1} + s_1) + \frac{c_{q_1} \rho_1^{q_1} \nu_1^{q_1} \delta_1^{q_1}}{\lambda_1^{q_1}}}, \\ \phi_2 &= \vartheta + \sqrt[q_2]{1 - q_2 e_2 + (c_{q_2} + q_2 \sigma) \epsilon^{q_2}} + \frac{\rho_2 \tau_2^{q_2 - 1} (\nu_2 \pi_2 \gamma' + \psi_2)}{\lambda_2 (r_2 - \rho_2 m_2)}, \\ \varphi_2 &= \frac{\rho_2 \tau_2^{q_2 - 1} (\rho_1 \xi + \rho_2 \varpi)}{\lambda_2 (r_2 - \rho_2 m_2)}, \\ \phi_1 &= \frac{\rho_1 \tau_1^{q_1 - 1} (\iota_1 \xi' + \iota_2 \zeta')}{\lambda_1 (r_1 - \rho_1 m_1)} \end{aligned} \tag{4.48}$$

and φ_2 is the constant as in (4.40). From (4.4), we know that $0 \leq \omega < 1$ and so it follows from (4.46) that $\mathfrak{R}_{\rho_1, \rho_2} : X_1 \times X_2 \rightarrow X_1 \times X_2$ is a contractive mapping. Hence Lemma 2.3 implies that $\mathfrak{R}_{\rho_1, \rho_2}$ has a fixed point in $X_1 \times X_2$; that is, there exists a point $(x^*, y^*) \in X_1 \times X_2$ such that $(x^*, y^*) \in \mathfrak{R}_{\rho_1, \rho_2}(x^*, y^*)$. Now, it follows from (4.6), (4.8), and Lemma 4.1 that $(x^*, y^*, z^*, u^*, v^*, m^*, n^*, t^*, w^*)$ is a solution of System 3.1 and this is the desired result. This completes the proof. \square

By using Theorem 4.2, we can derive the following.

Theorem 4.3. *Let X_i, A_i, η_i ($i = 1, 2$), $S, \mathcal{T}, \mathcal{L}, \mathfrak{D}, G, \mathcal{W}, \mathcal{K}, S, T, L, D, G, W, K, M, N, E, P, F, Q, p, l, h, k, f'$, and g' be the same as in Theorem 4.2. Assume that $f : X_1 \rightarrow X_1$ is κ -strongly accretive μ -Lipschitz continuous and $g : X_2 \rightarrow X_2$ is σ -strongly accretive ϵ -Lipschitz continuous.*

Further, if there exist constants $\rho_1 \in (0, r_1/m_1)$ and $\rho_2 \in (0, r_2/m_2)$ such that (4.2) and (4.3) hold and

$$\begin{aligned} \varsigma + \zeta + \sqrt[q_1]{1 - q_1 \kappa + c_{q_1} \mu^{q_1}} &< 1, \\ \vartheta + \sqrt[q_2]{1 - q_2 \sigma + c_{q_2} \epsilon^{q_2}} &< 1, \\ \sqrt[q_1]{\beta_1^{q_1} (\mu + \zeta)^{q_1} - q_1 \frac{\rho_1}{\lambda_1} (-\theta_1 \nu_1^{q_1} \delta_1^{q_1} + s_1) + \frac{c_{q_1} \rho_1^{q_1} \nu_1^{q_1} \delta_1^{q_1}}{\lambda_1^{q_1}}} &< \frac{\tau_1^{1 - q_1} \lambda_1}{\rho_1} (r_1 - \rho_1 m_1) \chi_1 - \nu_2 \delta_2 \gamma, \\ \sqrt[q_2]{\beta_2^{q_2} \epsilon^{q_2} - q_2 \frac{\rho_2}{\lambda_2} (-\theta_2 \nu_1^{q_2} \pi_1^{q_2} + s_2) + \frac{c_{q_2} \rho_2^{q_2} \nu_1^{q_2} \pi_1^{q_2}}{\lambda_2^{q_2}}} &< \frac{\tau_2^{1 - q_2} \lambda_2}{\rho_2} (r_2 - \rho_2 m_2) \chi_2 - \nu_2 \pi_2 \gamma', \end{aligned} \tag{4.49}$$

where

$$\begin{aligned}
 X_1 &= 1 - \left(\varsigma + \zeta + \sqrt[q_1]{1 - q_1\kappa + c_{q_1}\mu^{q_1}} \right) - \frac{\rho_2\tau_2^{q_2-1}(\rho_1\xi + \rho_2\varpi)}{\lambda_2(r_2 - \rho_2m_2)}, \\
 X_2 &= 1 - \left(\vartheta + \sqrt[q_2]{1 - q_2\sigma + c_{q_2}\epsilon^{q_2}} \right) - \frac{\rho_1\tau_1^{q_1-1}(\iota_1\xi' + \iota_2\zeta')}{\lambda_1(r_1 - \rho_1m_1)},
 \end{aligned}
 \tag{4.50}$$

λ_1, λ_2 are the same as in System 3.1, and c_{q_1}, c_{q_2} are two constants guaranteed by Lemma 2.1, then System 3.1 admits a solution.

Theorem 4.4. Let X_i, A_i, η_i ($i = 1, 2$), $p, l, h, k, f, g, F, Q, M, N$, and P be the same as in Theorem 4.2. Assume that $T : X_1 \rightarrow X_1$ is ζ -Lipschitz continuous, and $S, L, D : X_1 \rightarrow CB(X_1)$ and $G, W, K : X_2 \rightarrow CB(X_2)$ are ξ - \widehat{H}_1 -Lipschitz continuous, γ - \widehat{H}_1 -Lipschitz continuous, ϖ - \widehat{H}_1 -Lipschitz continuous, ξ' - \widehat{H}_2 -Lipschitz continuous, ζ' - \widehat{H}_2 -Lipschitz continuous, and γ' - \widehat{H}_2 -Lipschitz continuous, respectively. Suppose that $E : X_1 \times X_2 \rightarrow X_1$ is a single-valued mapping such that $E(\cdot, y)$ is ν_1 -Lipschitz continuous in the first variable and $E(x, \cdot)$ is ι_1 -Lipschitz continuous in the second variable for all $(x, y) \in X_1 \times X_2$, and $E(p(\cdot), y)$ is a (θ_1, s_1) -relaxed cocercive mapping with respect to $f' = A_1 \circ (f - T)$ defined by $f'(x) = A_1 \circ (f(x) - T(x)) = A_1(f(x) - T(x))$ for all $x \in X_1$. If there exist constants $\rho_1 \in (0, r_1/m_1)$ and $\rho_2 \in (0, r_2/m_2)$ such that conditions (4.2)–(4.4) hold, then System 3.4 has a solution $(x^*, y^*, z^*, u^*, m^*, s^*, t^*, w^*)$.

Theorem 4.5. Let X_i, A_i, η_i ($i = 1, 2$), $p, l, h, k, E, F, P, Q, M, N, S, T, L, D, G, W$, and K be the same as in Theorem 4.4. Suppose that $f : X_1 \rightarrow X_1$ is κ -strongly accretive and μ -Lipschitz continuous and $g : X_2 \rightarrow X_2$ is σ -strongly accretive and ϵ -Lipschitz continuous. If there exist constants $\rho_1 \in (0, r_1/m_1)$ and $\rho_2 \in (0, r_2/m_2)$ such that conditions (4.2), (4.3), and (4.49) hold, then System 3.3 has a solution $(x^*, y^*, z^*, u^*, m^*, s^*, t^*, w^*)$.

5. Iterative Algorithm and Convergence

In this section, motivated by Theorems 4.2 and 4.4, Lemmas 4.1 and 2.4, we construct the following iterative algorithms for approximating solutions of Systems 3.1 and 3.3 and discuss the convergence analysis of the algorithms.

Algorithm 5.1. Let $X_i, A_i, \eta_i, \lambda_i$ ($i = 1, 2$), $E, P, F, Q, p, l, h, k, f, g, M, N, S, \tau, \mathcal{L}, \mathfrak{D}, G, \mathcal{W}, \mathcal{K}, S, T, L, D, G, W, K, a$ and b be the same as in System 3.1. For any given $(x_0, y_0) \in X_1 \times X_2$, $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d} : X_1 \rightarrow [0, 1]$ and $\tilde{e}, \tilde{f}, \tilde{g} : X_2 \rightarrow [0, 1]$ for all $n \geq 0$ and an element $(x, y, z, u, v, m, s, t, w) \in X_1 \times X_2 \times X_1 \times X_1 \times X_1 \times X_1 \times X_2 \times X_2 \times X_2$, define the iterative sequence $\{(x_n, y_n, z_n, u_n, v_n, m_n, s_n, t_n, w_n)\}_{n=0}^\infty$ by

$$\begin{aligned}
 x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n \left(x_n - f(x_n) + v_n + R_{M(\cdot, x_n), \rho_1}^{\eta_1, A_1}(\Theta_n) \right) + \alpha_n e_n + r_n, \\
 y_{n+1} &= (1 - \alpha_n)y_n + \alpha_n \left(y_n - g(y_n) + R_{N(\cdot, y_n), \rho_2}^{\eta_2, A_2}(\Omega_n) \right) + \alpha_n f_n + k_n, \\
 \mathcal{S}_{x_n}(u_n) &\geq \tilde{a}(x_n), \quad \|u_n - u\| \leq \left(1 + \frac{1}{1+n} \right) \widehat{H}_1(S(x_n), S(x)),
 \end{aligned}$$

$$\begin{aligned}
\mathcal{T}_{x_n}(v_n) &\geq \tilde{b}(x_n), & \|v_n - v\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_1(T(x_n), T(x)), \\
\mathcal{L}_{x_n}(z_n) &\geq \tilde{c}(x_n), & \|z_n - z\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_1(L(x_n), L(x)), \\
\mathcal{D}_{x_n}(m_n) &\geq \tilde{d}(x_n), & \|m_n - m\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_1(D(x_n), D(x)), \\
\mathcal{G}_{y_n}(w_n) &\geq \tilde{e}(y_n), & \|w_n - w\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_2(G(y_n), G(y)), \\
\mathcal{W}_{y_n}(t_n) &\geq \tilde{f}(y_n), & \|t_n - t\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_2(W(y_n), W(y)), \\
\mathcal{K}_{y_n}(s_n) &\geq \tilde{g}(y_n), & \|s_n - s\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_2(K(y_n), K(y)),
\end{aligned} \tag{5.1}$$

where

$$\begin{aligned}
\Theta_n &= A_1(f(x_n) - v_n) - \frac{\rho_1}{\lambda_1}(E(p(x_n), w_n) + P(l(z_n), t_n) - a), \\
\Omega_n &= A_2(g(y_n)) - \frac{\rho_2}{\lambda_2}(F(u_n, h(y_n)) + Q(m_n, k(s_n)) - b),
\end{aligned} \tag{5.2}$$

ρ_1 and ρ_2 are constants, $\{\alpha_n\}$ is a sequence in $[0, 1]$ with $\sum_{n=0}^{\infty} \alpha_n = \infty$, and $\{(e_n, f_n)\}_{n=0}^{\infty}$ and $\{(r_n, k_n)\}_{n=0}^{\infty}$ are two sequences in $X_1 \times X_2$ to take into account a possible inexact computation of the resolvent operator point satisfying the following conditions:

$$\begin{aligned}
e_n &= e'_n + e''_n, & f_n &= f'_n + f''_n, \\
\lim_{n \rightarrow \infty} \|(e'_n, f'_n)\|_* &= 0, \\
\sum_{n=0}^{\infty} \|(e''_n, f''_n)\|_* &< \infty, & \sum_{n=0}^{\infty} \|(r_n, k_n)\|_* &< \infty.
\end{aligned} \tag{5.3}$$

Algorithm 5.2. Assume that $X_i, A_i, \eta_i, \lambda_i$ ($i = 1, 2$), $E, P, F, Q, p, l, h, k, f, g, M, N, S, T, L, D, G, W, K, a$ and b are the same as in System 3.4. For any given $(x_0, y_0) \in X_1 \times X_2$, $n \geq 0$ and an element $(x, y, z, u, m, s, t, w) \in X_1 \times X_2 \times X_1 \times X_1 \times X_2 \times X_2 \times X_2$, define the iterative sequence $\{(x_n, y_n, z_n, u_n, m_n, s_n, t_n, w_n)\}_{n=0}^{\infty}$ by

$$\begin{aligned}
x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n \left(x_n - f(x_n) + T(x_n) + R_{M(\cdot, x_n), \rho_1}^{\eta_1, A_1}(\Theta'_n) \right) + \alpha_n e_n + r_n, \\
y_{n+1} &= (1 - \alpha_n)y_n + \alpha_n \left(y_n - g(y_n) + R_{N(\cdot, y_n), \rho_2}^{\eta_2, A_2}(\Omega_n) \right) + \alpha_n f_n + k_n, \\
u_n &\in S(x_n), & \|u_n - u\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_1(S(x_n), S(x)),
\end{aligned}$$

$$\begin{aligned}
z_n \in L(x_n), \quad \|z_n - z\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_1(L(x_n), L(x)), \\
m_n \in D(x_n), \quad \|m_n - m\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_1(D(x_n), D(x)), \\
w_n \in G(y_n), \quad \|w_n - w\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_2(G(y_n), G(y)), \\
t_n \in W(y_n), \quad \|t_n - t\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_2(W(y_n), W(y)), \\
s_n \in K(y_n), \quad \|s_n - s\| &\leq \left(1 + \frac{1}{1+n}\right) \widehat{H}_2(K(y_n), K(y)),
\end{aligned} \tag{5.4}$$

where $\Theta'_n = A_1(f(x_n) - T(x_n)) - (\rho_1/\lambda_1)(E(p(x_n), w_n) + P(l(z_n), t_n) - a)$, Ω_n , ρ_1 , ρ_2 , $\{\alpha_n\}$, $\{(e_n, f_n)\}_{n=0}^\infty$ and $\{(r_n, k_n)\}_{n=0}^\infty$ are the same as in Algorithm 5.1.

Remark 5.3. If $e_n = f_n = 0$ for all $n \geq 0$, $L = D = W = K \equiv 0$, $P = Q \equiv 0$, $a = b = 0$, and $\lambda_1 = \lambda_2 = 1$, then Algorithms 5.1 and 5.2 reduce to Algorithms 4.1 and 4.2 of [38]. In particular, when we choose suitable $\alpha_n, e_n, f_n, r_n, k_n, A_i, \eta_i$ ($i = 1, 2$), $E, P, F, Q, p, l, h, k, f, g, M, N, S, \mathcal{T}, \mathcal{L}, \mathfrak{D}, \mathcal{C}, \mathcal{W}, \mathcal{K}, S, T, L, D, G, W, K$, and the spaces X_1, X_2 , then Algorithms 5.1 and 5.2 can be degenerated to a number of algorithms involving many known algorithms due to classes of variational inequalities and variational inclusions (see, e.g., [38, 55, 56, 58–60] and the references therein).

Lemma 5.4. Let $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ be three nonnegative real sequences satisfying the following condition: there exists a natural number n_0 such that

$$a_{n+1} \leq (1 - t_n)a_n + b_n t_n + c_n, \quad \forall n \geq n_0, \tag{5.5}$$

where $t_n \in [0, 1]$, $\sum_{n=0}^\infty t_n = \infty$, $\lim_{n \rightarrow \infty} b_n = 0$, $\sum_{n=0}^\infty c_n < \infty$. Then $\lim_{n \rightarrow \infty} a_n = 0$.

Proof. The proof directly follows from Liu [64, Lemma 2]. \square

Theorem 5.5. Let X_i, A_i, η_i ($i = 1, 2$), $E, P, F, Q, p, l, h, k, f, g, M, N, S, \mathcal{T}, \mathcal{L}, \mathfrak{D}, \mathcal{C}, \mathcal{W}, \mathcal{K}, S, T, L, D, G, W$, and K be the same as in Theorem 4.2. Suppose that all the conditions of Theorem 4.2 hold. Then the iterative sequence $\{(x_n, y_n, z_n, u_n, v_n, m_n, s_n, t_n, w_n)\}_{n=0}^\infty$ generated by Algorithm 5.1 converges strongly to a solution $(x^*, y^*, z^*, u^*, v^*, m^*, s^*, t^*, w^*)$ of System 3.1.

Proof. It follows from Theorem 4.2 that System 3.1 has a solution $(x^*, y^*, z^*, u^*, v^*, m^*, s^*, t^*, w^*)$. Hence, by Lemma 4.1, we have

$$\begin{aligned}
f(x^*) &= v^* + R_{M(\cdot, x^*), \rho_1}^{\eta_1, A_1} \left[A_1(f(x^*) - v^*) - \frac{\rho_1}{\lambda_1} (E(p(x^*), w^*) + P(l(z^*), t^*) - a) \right], \\
g(y^*) &= R_{M(\cdot, y^*), \rho_2}^{\eta_2, A_2} \left[A_2(g(y^*)) - \frac{\rho_2}{\lambda_2} (F(u^*, h(y^*)) + Q(m^*, k(s^*)) - b) \right].
\end{aligned} \tag{5.6}$$

Using (5.1), (5.6), and our assumptions, it follows that

$$\begin{aligned}
& \|x_{n+1} - x^*\| \leq (1 - \alpha_n) \|x_n - x^*\| + \alpha_n \\
& \quad \times \left(\|x_n - x^* - (f(x_n) - f(x^*))\| + \|v_n - v^*\| \right. \\
& \quad \quad + \left\| R_{M(\cdot, x_n), \rho_1}^{\eta_1, A_1} \left[A_1(f(x_n) - v_n) - \frac{\rho_1}{\lambda_1} (E(p(x_n), w_n) + P(l(z_n), t_n) - a) \right] \right. \\
& \quad \quad \quad \left. \left. - R_{M(\cdot, x^*), \rho_1}^{\eta_1, A_1} \left[A_1(f(x^*) - v^*) - \frac{\rho_1}{\lambda_1} (E(p(x^*), w^*) + P(l(z^*), t^*) - a) \right] \right\| \right) \\
& \quad + \alpha_n \|e_n\| + \|r_n\| \\
& \leq (1 - \alpha_n) \|x_n - x^*\| + \alpha_n \\
& \quad \times \left(\|x_n - x^* - (f(x_n) - f(x^*))\| + \|v_n - v^*\| \right. \\
& \quad \quad + \left\| R_{M(\cdot, x_n), \rho_1}^{\eta_1, A_1} \left[A_1(f(x_n) - v_n) - \frac{\rho_1}{\lambda_1} (E(p(x_n), w_n) + P(l(z_n), t_n) - a) \right] \right. \\
& \quad \quad \quad \left. \left. - R_{M(\cdot, x_n), \rho_1}^{\eta_1, A_1} \left[A_1(f(x^*) - v^*) - \frac{\rho_1}{\lambda_1} (E(p(x^*), w^*) + P(l(z^*), t^*) - a) \right] \right\| \right) \\
& \quad \quad + \left\| R_{M(\cdot, x_n), \rho_1}^{\eta_1, A_1} \left[A_1(f(x^*) - v^*) - \frac{\rho_1}{\lambda_1} (E(p(x^*), w^*) + P(l(z^*), t^*) - a) \right] \right. \\
& \quad \quad \quad \left. \left. - R_{M(\cdot, x^*), \rho_1}^{\eta_1, A_1} \left[A_1(f(x^*) - v^*) - \frac{\rho_1}{\lambda_1} (E(p(x^*), w^*) + P(l(z^*), t^*) - a) \right] \right\| \right) \\
& \quad + \alpha_n (\|e'_n\| + \|e''_n\|) + \|r_n\| \\
& \leq (1 - \alpha_n) \|x_n - x^*\| + \alpha_n \\
& \quad \times \left\{ \|x_n - x^* - (f(x_n) - f(x^*))\| + \|v_n - v^*\| + \varsigma \|x_n - x^*\| + \frac{\tau_1^{q_1-1}}{r_1 - \rho_1 m_1} \right. \\
& \quad \quad \times \left(\frac{\rho_1}{\lambda_1} (\|E(p(x_n), w_n) - E(p(x_n), w^*)\| + \|P(l(z_n), t_n) - P(l(z_n), t^*)\|) \right. \\
& \quad \quad \quad \left. \left. + \|P(l(z_n), t^*) - P(l(z^*), t^*)\|) \right) \right. \\
& \quad \quad \quad \left. \left. + \left\| A_1(f(x_n) - v_n) - A_1(f(x^*) - v^*) - \frac{\rho_1}{\lambda_1} (E(p(x_n), w^*) - E(p(x^*), w^*)) \right\| \right\} \right) \\
& \quad + \alpha_n \|e'_n\| + \|e''_n\| + \|r_n\| \\
& \leq (1 - \alpha_n) \|x_n - x^*\| + \alpha_n (\varphi_1(n) \|x_n - x^*\| + \phi_1(n) \|y_n - y^*\|) + \alpha_n \|e'_n\| + \|e''_n\| + \|r_n\|, \tag{5.7}
\end{aligned}$$

where

$$\begin{aligned}
 \varphi_1(n) &= \varsigma + \zeta \left(1 + \frac{1}{1+n}\right) \\
 &\quad + \sqrt[q_1]{1 - q_1 e_1 + (c_{q_1} + q_1 \kappa) \mu^{q_1}} + \frac{\rho_1 \tau_1^{q_1-1} (\nu_2 \delta_2 \Upsilon (1 + 1/(1+n)) + \varphi_1(n))}{\lambda_1 (r_1 - \rho_1 m_1)}, \\
 \psi_1(n) &= \sqrt[q_1]{\beta_1^{q_1} \left(\mu + \zeta \left(1 + \frac{1}{1+n}\right)\right)^{q_1} - q_1 \frac{\rho_1}{\lambda_1} (-\theta_1 \nu_1^{q_1} \delta_1^{q_1} + s_1) + \frac{c_{q_1} \rho_1^{q_1} \nu_1^{q_1} \delta_1^{q_1}}{\lambda_1^{q_1}}}, \\
 \phi_1(n) &= \frac{\rho_1 \tau_1^{q_1-1} (\iota_1 \xi' + \iota_2 \zeta') (1 + 1/(1+n))}{\lambda_1 (r_1 - \rho_1 m_1)}.
 \end{aligned} \tag{5.8}$$

Similarly, we have

$$\begin{aligned}
 \|y_{n+1} - y^*\| &\leq (1 - \alpha_n) \|y_n - y^*\| + \alpha_n \\
 &\quad \times \left(\|y_n - y^* - (g(y_n) - g(y^*))\| \right. \\
 &\quad \left. + \left\| R_{N(\cdot, y_n), \rho_2}^{\eta_2, A_2} \left[A_2(g(y_n)) - \frac{\rho_2}{\lambda_2} (F(u_n, h(y_n)) + Q(m_n, k(s_n)) - b) \right] \right. \right. \\
 &\quad \left. \left. - R_{N(\cdot, y^*), \rho_2}^{\eta_2, A_2} \left[A_2(g(y^*)) - \frac{\rho_2}{\lambda_2} (F(u^*, h(y^*)) + Q(m^*, k(s^*)) - b) \right] \right\| \right) \\
 &\quad + \alpha_n \|f_n\| + \|k_n\| \\
 &\leq (1 - \alpha_n) \|y_n - y^*\| + \alpha_n \\
 &\quad \times \left(\|y_n - y^* - (g(y_n) - g(y^*))\| \right. \\
 &\quad \left. + \left\| R_{N(\cdot, y_n), \rho_2}^{\eta_2, A_2} \left[A_2(g(y_n)) - \frac{\rho_2}{\lambda_2} (F(u_n, h(y_n)) + Q(m_n, k(s_n)) - b) \right] \right. \right. \\
 &\quad \left. \left. - R_{N(\cdot, y_n), \rho_2}^{\eta_2, A_2} \left[A_2(g(y^*)) - \frac{\rho_2}{\lambda_2} (F(u^*, h(y^*)) + Q(m^*, k(s^*)) - b) \right] \right\| \right) \\
 &\quad + \left\| R_{N(\cdot, y^*), \rho_2}^{\eta_2, A_2} \left[A_2(g(y^*)) - \frac{\rho_2}{\lambda_2} (F(u^*, h(y^*)) + Q(m^*, k(s^*)) - b) \right] \right. \\
 &\quad \left. - R_{N(\cdot, y^*), \rho_2}^{\eta_2, A_2} \left[A_2(g(y^*)) - \frac{\rho_2}{\lambda_2} (F(u^*, h(y^*)) + Q(m^*, k(s^*)) - b) \right] \right\| \\
 &\quad + \alpha_n (\|f'_n\| + \|f''_n\|) + \|k_n\|
 \end{aligned}$$

$$\begin{aligned}
 &\leq (1 - \alpha_n) \|y_n - y^*\| + \alpha_n \\
 &\quad \times \left\{ \|y_n - y^* - (g(y_n) - g(y^*))\| + \vartheta \|y_n - y^*\| + \frac{\tau_2^{q_2-1}}{r_2 - \rho_2 m_2} \right. \\
 &\quad \times \left(\frac{\rho_2}{\lambda_2} (\|F(u_n, h(y_n)) - F(u^*, h(y_n))\| + \|Q(m_n, k(s_n)) - Q(m^*, k(s_n))\| \right. \\
 &\quad \quad \left. + \|Q(m^*, k(s_n)) - Q(m^*, k(s^*))\|) \right. \\
 &\quad \quad \left. \left. + \left\| A_2(g(y_n)) - A_2(g(y^*)) - \frac{\rho_2}{\lambda_2} (F(u^*, h(y_n)) - F(u^*, h(y^*))) \right\| \right) \right\} \\
 &\quad + \alpha_n \|f'_n\| + \|f''_n\| + \|k_n\| \\
 &\leq (1 - \alpha_n) \|y_n - y^*\| + \alpha_n (\varphi_2(n) \|x_n - x^*\| + \phi_2(n) \|y_n - y^*\|) \\
 &\quad + \alpha_n \|f'_n\| + \|f''_n\| + \|k_n\|,
 \end{aligned} \tag{5.9}$$

where

$$\begin{aligned}
 \phi_2(n) &= \vartheta + \sqrt[q_2]{1 - q_2 e_2 + (c_{q_2} + q_2 \sigma) \epsilon^{q_2}} + \frac{\rho_2 \tau_2^{q_2-1} (v_2 \pi_2 \gamma' (1 + 1/(1+n)) + \psi_2)}{\lambda_2 (r_2 - \rho_2 m_2)}, \\
 \varphi_2(n) &= \frac{\rho_2 \tau_2^{q_2-1} (\rho_1 \xi + \rho_2 \varpi) (1 + 1/(1+n))}{\lambda_2 (r_2 - \rho_2 m_2)},
 \end{aligned} \tag{5.10}$$

and ψ_2 is the same as (4.40). By (5.7) and (5.9), we obtain

$$\begin{aligned}
 \|(x_{n+1}, y_{n+1}) - (x^*, y^*)\|_* &= \|x_{n+1} - x^*\| + \|y_{n+1} - y^*\| \leq (1 - \alpha_n) \|(x_n, y_n) - (x^*, y^*)\|_* \\
 &\quad + \alpha_n \max\{\varphi_1(n) + \varphi_2(n), \phi_1(n) + \phi_2(n)\} \|(x_n, y_n) - (x^*, y^*)\|_* \\
 &\quad + \alpha_n \| (e'_n, f'_n) \|_* + \| (e''_n, f''_n) \|_* + \| (r_n, k_n) \|_* \\
 &= (1 - (1 - \omega(n)) \alpha_n) \|(x_n, y_n) - (x^*, y^*)\|_* \\
 &\quad + \alpha_n \| (e'_n, f'_n) \|_* + \| (e''_n, f''_n) \|_* + \| (r_n, k_n) \|_*,
 \end{aligned} \tag{5.11}$$

where

$$\omega(n) = \max\{\varphi_1(n) + \varphi_2(n), \phi_1(n) + \phi_2(n)\}. \tag{5.12}$$

Now, $\omega(n) \rightarrow \omega = \max\{\varphi_1 + \varphi_2, \phi_1 + \phi_2\}$ as $n \rightarrow \infty$, where $\varphi_1, \varphi_2, \phi_1$, and ϕ_2 are the constants as in (4.48).

Since $\widehat{\omega} = (1/2)(\omega + 1) \in (\omega, 1)$, deduce that there exists $n_0 \geq 1$ such that $\omega(n) < \widehat{\omega}$, for all $n \geq n_0$. Accordingly, it follows from (5.11) that for all $n \geq n_0$,

$$\begin{aligned} \|(x_{n+1}, y_{n+1}) - (x^*, y^*)\|_* &\leq (1 - (1 - \widehat{\omega})\alpha_n) \|(x_n, y_n) - (x^*, y^*)\|_* + \alpha_n \|(e'_n, f'_n)\|_* \\ &\quad + \|(e''_n, f''_n)\|_* + \|(r_n, k_n)\|_*. \end{aligned} \quad (5.13)$$

Letting

$$\begin{aligned} a_n &= \|(x_{n+1}, y_{n+1}) - (x^*, y^*)\|_*, & t_n &= (1 - \widehat{\omega})\alpha_n, \\ b_n &= \frac{\|(e'_n, f'_n)\|_*}{1 - \widehat{\omega}}, & c_n &= \|(e''_n, f''_n)\|_* + \|(r_n, k_n)\|_*, \end{aligned} \quad (5.14)$$

then (5.13) can be written as

$$a_{n+1} \leq (1 - t_n)a_n + b_n t_n + c_n, \quad \forall n \geq 0. \quad (5.15)$$

Therefore, it follows from Lemma 5.4 that $\lim_{n \rightarrow \infty} a_n = 0$ and so the sequence $\{(x_n, y_n, z_n, u_n, v_n, m_n, s_n, t_n, w_n)\}_{n=0}^{\infty}$ defined by Algorithm 5.1 converges strongly to a solution $(x^*, y^*, z^*, u^*, v^*, m^*, s^*, t^*, w^*)$ of System 3.1. \square

Theorem 5.6. Suppose that X_i, A_i, η_i ($i = 1, 2$), $E, P, F, Q, p, l, h, k, f, g, M, N, S, \mathcal{T}, \mathcal{L}, \mathcal{D}, \mathcal{G}, \mathcal{W}, \mathcal{K}, S, T, L, D, G, W$, and K are the same as in Theorem 4.3. Assume that all the conditions of Theorem 4.3 hold. Then the iterative sequence $\{(x_n, y_n, z_n, u_n, v_n, m_n, s_n, t_n, w_n)\}_{n=0}^{\infty}$ generated by Algorithm 5.1 converges strongly to the solution $(x^*, y^*, z^*, u^*, v^*, m^*, s^*, t^*, w^*)$ of System 3.1.

Theorem 5.7. Assume that X_i, A_i, η_i ($i = 1, 2$), $E, P, F, Q, p, l, h, k, f, g, M, N, S, T, L, D, G, W$, and K are the same as in Theorem 4.4. Suppose that all the conditions of Theorem 4.4 hold. Then the iterative sequence $\{(x_n, y_n, z_n, u_n, m_n, s_n, t_n, w_n)\}_{n=0}^{\infty}$ generated by Algorithm 5.2 converges strongly to a solution $(x^*, y^*, z^*, u^*, m^*, s^*, t^*, w^*)$ of System 3.3.

Theorem 5.8. Let X_i, A_i, η_i ($i = 1, 2$), $E, P, F, Q, p, l, h, k, f, g, M, N, S, T, L, D, G, W$, and K be the same as in Theorem 4.5. Suppose that all the conditions of Theorem 4.5 hold. Then the iterative sequence $\{(x_n, y_n, z_n, u_n, m_n, s_n, t_n, w_n)\}_{n=0}^{\infty}$ generated by Algorithm 5.2 converges strongly to a solution $(x^*, y^*, z^*, u^*, m^*, s^*, t^*, w^*)$ of System 3.3.

Remark 5.9. The following should be noticed.

- (1) Theorem 3.1 in [54] is a special case of the Theorems 4.2 and 4.3. Moreover, Theorems 4.4 and 4.5 improve and extend Theorem 3.2 [54].
- (2) In view of Remark 5.3, Theorems 5.5 and 5.6 improve and generalize Theorem 4.1 in [54]. Also, Theorems 5.7 and 5.8 are extensions of Theorem 4.2 in [54].

Remark 5.10. When M and N are (A, η) -monotone operators, A -accretive mappings, A -monotone operators, (H, η) -accretive mappings, (H, η) -monotone operators, or H -monotone operators, respectively, from Theorems 4.2–4.5 and 5.5–5.8, we can obtain the existence and convergence results of solutions for Systems 3.1 and 3.4. In brief, for a suitable and

appropriate choice of the mappings A_i, η_i ($i = 1, 2$), $E, P, F, Q, p, l, h, k, f, g, M, N, S, \mathcal{T}, \mathcal{L}, \mathcal{D}, \mathcal{G}, \mathcal{W}, \mathcal{K}, S, T, L, D, G, W, K$, and the spaces X_1, X_2 , Theorems 4.2–4.5 and 5.5–5.8 include many known results of the generalized variational inclusions as special cases (see [29–35, 38, 39, 55–60] and the references therein).

Acknowledgment

This paper was written while the first author visiting the KCL. This work was supported by the Korea Research Foundation Grant funded by the Korean Government (KRF-2008-313-C00050). Also, this paper is partially supported by the Research Center in Algebraic Hyperstructures and Fuzzy Mathematics, University of Mazandaran, Babolsar, Iran.

References

- [1] G. Stampacchia, "Formes bilinéaires coercitives sur les ensembles convexes," *Comptes Rendus de l'Académie des Sciences*, vol. 258, pp. 4413–4416, 1964.
- [2] A. Hassouni and A. Moudafi, "A perturbed algorithm for variational inclusions," *Journal of Mathematical Analysis and Applications*, vol. 185, no. 3, pp. 706–712, 1994.
- [3] S. Adly, "Perturbed algorithms and sensitivity analysis for a general class of variational inclusions," *Journal of Mathematical Analysis and Applications*, vol. 201, no. 2, pp. 609–630, 1996.
- [4] C. Baiocchi and A. Capelo, *Variational and Quasivariational Inequalities*, A Wiley-Interscience Publications, John Wiley & Sons, New York, NY, USA, 1984.
- [5] V. Barbu, *Nonlinear Semigroups and Differential Equations in Banach Spaces*, Noordhoff, Leyden, The Netherlands, 1979.
- [6] K. Deimling, *Nonlinear Functional Analysis*, Springer, Berlin, Germany, 1985.
- [7] X. P. Ding, "Perturbed proximal point algorithms for generalized quasivariational inclusions," *Journal of Mathematical Analysis and Applications*, vol. 210, no. 1, pp. 88–101, 1997.
- [8] J.-S. Pang, "Asymmetric variational inequality problems over product sets: applications and iterative methods," *Mathematical Programming*, vol. 31, no. 2, pp. 206–219, 1985.
- [9] G. Cohen and F. Chaplais, "Nested monotony for variational inequalities over product of spaces and convergence of iterative algorithms," *Journal of Optimization Theory and Applications*, vol. 59, no. 3, pp. 369–390, 1988.
- [10] M. Bianchi, "Pseudo P -monotone operators and variational inequalities," Report 6, Istituto di econometria e Matematica per le decisioni economiche, Università Cattolica del sacro Cuore, Milan, Italy, 1993.
- [11] Q. H. Ansari and J.-C. Yao, "A fixed point theorem and its applications to a system of variational inequalities," *Bulletin of the Australian Mathematical Society*, vol. 59, no. 3, pp. 433–442, 1999.
- [12] Q. H. Ansari, S. Schaible, and J. C. Yao, "System of vector equilibrium problems and its applications," *Journal of Optimization Theory and Applications*, vol. 107, no. 3, pp. 547–557, 2000.
- [13] E. Allevi, A. Gnudi, and I. V. Konnov, "Generalized vector variational inequalities over product sets," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 47, no. 1, pp. 573–582, 2001.
- [14] G. Kassay, J. Kolumbán, and Z. Páles, "On Nash stationary points," *Publicationes Mathematicae Debrecen*, vol. 54, no. 3-4, pp. 267–279, 1999.
- [15] G. Kassay, J. Kolumbán, and Z. Páles, "Factorization of Minty and Stampacchia variational inequality systems," *European Journal of Operational Research*, vol. 143, no. 2, pp. 377–389, 2002.
- [16] J.-W. Peng, "System of generalised set-valued quasi-variational-like inequalities," *Bulletin of the Australian Mathematical Society*, vol. 68, no. 3, pp. 501–515, 2003.
- [17] J.-W. Peng, "Set-valued variational inclusions with T -accretive operators in Banach spaces," *Applied Mathematics Letters*, vol. 19, no. 3, pp. 273–282, 2006.
- [18] J.-W. Peng and X. Yang, "On existence of a solution for the system of generalized vector quasi-equilibrium problems with upper semicontinuous set-valued maps," *International Journal of Mathematics and Mathematical Sciences*, vol. 2005, no. 15, pp. 2409–2420, 2005.

- [19] R. U. Verma, "On a new system of nonlinear variational inequalities and associated iterative algorithms," *Mathematical Sciences Research Hot-Line*, vol. 3, no. 8, pp. 65–68, 1999.
- [20] R. U. Verma, "Projection methods, algorithms, and a new system of nonlinear variational inequalities," *Computers & Mathematics with Applications*, vol. 41, no. 7-8, pp. 1025–1031, 2001.
- [21] R. U. Verma, "Iterative algorithms and a new system of nonlinear quasivariational inequalities," *Advances in Nonlinear Variational Inequalities*, vol. 4, no. 1, pp. 117–124, 2001.
- [22] R. U. Verma, "Generalized system for relaxed cocoercive variational inequalities and projection methods," *Journal of Optimization Theory and Applications*, vol. 121, no. 1, pp. 203–210, 2004.
- [23] R. U. Verma, "General convergence analysis for two-step projection methods and applications to variational problems," *Applied Mathematics Letters*, vol. 18, no. 11, pp. 1286–1292, 2005.
- [24] J. K. Kim and D. S. Kim, "A new system of generalized nonlinear mixed variational inequalities in Hilbert spaces," *Journal of Convex Analysis*, vol. 11, no. 1, pp. 235–243, 2004.
- [25] Y. J. Cho, Y. P. Fang, N. J. Huang, and H. J. Hwang, "Algorithms for systems of nonlinear variational inequalities," *Journal of the Korean Mathematical Society*, vol. 41, no. 3, pp. 489–499, 2004.
- [26] R. P. Agarwal, N.-J. Huang, and M.-Y. Tan, "Sensitivity analysis for a new system of generalized nonlinear mixed quasi-variational inclusions," *Applied Mathematics Letters*, vol. 17, no. 3, pp. 345–352, 2004.
- [27] K. R. Kazmi and M. I. Bhat, "Iterative algorithm for a system of nonlinear variational-like inclusions," *Computers & Mathematics with Applications*, vol. 48, no. 12, pp. 1929–1935, 2004.
- [28] N.-J. Huang and Y.-P. Fang, "Generalized m -accretive mappings in Banach spaces," *Journal of Sichuan University*, vol. 38, pp. 591–592, 2001.
- [29] Y.-P. Fang and N.-J. Huang, " H -monotone operators and system of variational inclusions," *Communications on Applied Nonlinear Analysis*, vol. 11, no. 1, pp. 93–101, 2004.
- [30] W.-Y. Yan, Y.-P. Fang, and N.-J. Huang, "A new system of set-valued variational inclusions with H -monotone operators," *Mathematical Inequalities & Applications*, vol. 8, no. 3, pp. 537–546, 2005.
- [31] Y.-P. Fang, N.-J. Huang, and H. B. Thompson, "A new system of variational inclusions with (H, η) -monotone operators in Hilbert spaces," *Computers & Mathematics with Applications*, vol. 49, no. 2-3, pp. 365–374, 2005.
- [32] H.-Y. Lan, N.-J. Huang, and Y. J. Cho, "New iterative approximation for a system of generalized nonlinear variational inclusions with set-valued mappings in Banach spaces," *Mathematical Inequalities & Applications*, vol. 9, no. 1, pp. 175–187, 2006.
- [33] H.-Y. Lan, Q.-K. Liu, and J. Li, "Iterative approximation for a system of nonlinear variational inclusions involving generalized m -accretive mappings," *Nonlinear Analysis Forum*, vol. 9, no. 1, pp. 33–42, 2004.
- [34] Y.-P. Fang and N.-J. Huang, "Iterative algorithm for a system of variational inclusions involving H -accretive operators in Banach spaces," *Acta Mathematica Hungarica*, vol. 108, no. 3, pp. 183–195, 2005.
- [35] J.-W. Peng, D.-L. Zhu, and X.-P. Zheng, "Existence of solutions and convergence of a multistep iterative algorithm for a system of variational inclusions with (H, η) -accretive operators," *Fixed Point Theory and Applications*, vol. 2007, Article ID 93678, 20 pages, 2007.
- [36] R. U. Verma, "A-monotonicity and applications to nonlinear variational inclusion problems," *Journal of Applied Mathematics and Stochastic Analysis*, vol. 2004, no. 2, pp. 193–195, 2004.
- [37] R. U. Verma, "Sensitivity analysis for generalized strongly monotone variational inclusions based on the (A, η) -resolvent operator technique," *Applied Mathematics Letters*, vol. 19, no. 12, pp. 1409–1413, 2006.
- [38] H.-Y. Lan, Y. J. Cho, and R. U. Verma, "Nonlinear relaxed cocoercive variational inclusions involving (A, η) -accretive mappings in Banach spaces," *Computers & Mathematics with Applications*, vol. 51, no. 9-10, pp. 1529–1538, 2006.
- [39] H.-Y. Lan, "Stability of iterative processes with errors for a system of nonlinear (A, η) -accretive variational inclusions in Banach spaces," *Computers & Mathematics with Applications*, vol. 56, no. 1, pp. 290–303, 2008.
- [40] L. A. Zadeh, "Fuzzy sets," *Information and Computation*, vol. 8, pp. 338–353, 1965.
- [41] M. Alimohammady and M. Roohi, "Fuzzy minimal structure and fuzzy minimal vector spaces," *Chaos, Solitons & Fractals*, vol. 27, no. 3, pp. 599–605, 2006.
- [42] J.-P. Aubin, *Mathematical Methods of Game and Economic Theory*, vol. 7 of *Studies in Mathematics and Its Applications*, North-Holland, Amsterdam, The Netherlands, 1979.
- [43] D. Dubois and H. Prade, *Fuzzy Sets and Systems, Theory and Applications*, vol. 144 of *Mathematics in Science and Engineering*, Academic Press, New York, NY, USA, 1980.

- [44] M. A. Noor, "Two-step approximation schemes for multivalued quasi variational inclusions," *Nonlinear Functional Analysis and Applications*, vol. 7, no. 1, pp. 1–14, 2002.
- [45] H. I. Zimmermann, *Fuzzy Set Theory and Its Applications*, Kluwer Academic Publishers, Boston, Mass, USA, 1988.
- [46] S.-S. Chang and Y. G. Zhu, "On variational inequalities for fuzzy mappings," *Fuzzy Sets and Systems*, vol. 32, no. 3, pp. 359–367, 1989.
- [47] S.-S. Chang and N. J. Huang, "Generalized complementarity problems for fuzzy mappings," *Fuzzy Sets and Systems*, vol. 55, no. 2, pp. 227–234, 1993.
- [48] S.-S. Chang and N.-J. Huang, "Generalized quasi-complementarity problems for a pair of fuzzy mappings," *Journal of Fuzzy Mathematics*, vol. 4, no. 2, pp. 343–354, 1996.
- [49] H.-Y. Lan, Z.-Q. He, and J. Li, "Generalized nonlinear fuzzy quasi-variational-like inclusions involving maximal η -monotone mappings," *Nonlinear Analysis Forum*, vol. 8, no. 1, pp. 43–54, 2003.
- [50] M. A. Noor, "Variational inequalities for fuzzy mappings—I," *Fuzzy Sets and Systems*, vol. 55, no. 3, pp. 309–312, 1993.
- [51] M. A. Noor, "Variational inequalities for fuzzy mappings—II," *Fuzzy Sets and Systems*, vol. 97, no. 1, pp. 101–107, 1998.
- [52] M. A. Noor, "Variational inequalities for fuzzy mappings—III," *Fuzzy Sets and Systems*, vol. 110, no. 1, pp. 101–108, 2000.
- [53] M. A. Noor and E. A. Al-Said, "Quasi variational inequalities for fuzzy mappings," *Journal of Chinese Fuzzy Systems Association*, vol. 3, no. 2, pp. 89–96, 1997.
- [54] H.-Y. Lan and R. U. Verma, "Iterative algorithms for nonlinear fuzzy variational inclusion systems with (A, η) -accretive mappings in Banach spaces," *Advances in Nonlinear Variational Inequalities*, vol. 11, no. 1, pp. 15–30, 2008.
- [55] M.-M. Jin, "A new system of general nonlinear variational inclusions involving (A, η) -accretive mappings in Banach spaces," *Mathematical Inequalities & Applications*, vol. 11, no. 4, pp. 783–794, 2008.
- [56] M.-M. Jin, "Convergence and stability of iterative algorithm for a new system of (A, η) -accretive mapping inclusions in Banach spaces," *Computers & Mathematics with Applications*, vol. 56, no. 9, pp. 2305–2311, 2008.
- [57] H.-Y. Lan, J. H. Kim, and Y. J. Cho, "On a new system of nonlinear A -monotone multivalued variational inclusions," *Journal of Mathematical Analysis and Applications*, vol. 327, no. 1, pp. 481–493, 2007.
- [58] J. Lou, X.-F. He, and Z. He, "Iterative methods for solving a system of variational inclusions involving H - η -monotone operators in Banach spaces," *Computers & Mathematics with Applications*, vol. 55, no. 8, pp. 1832–1841, 2008.
- [59] J.-W. Peng and D. Zhu, "A new system of generalized mixed quasi-variational inclusions with (H, η) -monotone operators," *Journal of Mathematical Analysis and Applications*, vol. 327, no. 1, pp. 175–187, 2007.
- [60] J.-W. Peng and D. L. Zhu, "A system of variational inclusions with P - η -accretive operators," *Journal of Computational and Applied Mathematics*, vol. 216, no. 1, pp. 198–209, 2008.
- [61] H. K. Xu, "Inequalities in Banach spaces with applications," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 16, no. 12, pp. 1127–1138, 1991.
- [62] S. B. Nadler Jr., "Multi-valued contraction mappings," *Pacific Journal of Mathematics*, vol. 30, pp. 475–488, 1969.
- [63] Y.-P. Fang and N.-J. Huang, " H -accretive operators and resolvent operator technique for solving variational inclusions in Banach spaces," *Applied Mathematics Letters*, vol. 17, no. 6, pp. 647–653, 2004.
- [64] L. S. Liu, "Ishikawa and Mann iterative process with errors for nonlinear strongly accretive mappings in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 194, no. 1, pp. 114–125, 1995.