

Research Article

Iterative Schemes for Generalized Equilibrium Problem and Two Maximal Monotone Operators

L. C. Zeng,^{1,2} Y. C. Lin,³ and J. C. Yao⁴

¹ Department of Mathematics, Shanghai Normal University, Shanghai 200234, China

² Science Computing Key Laboratory of Shanghai Universities, Shanghai 200234, China

³ Department of Occupational Safety and Health, China Medical University, Taichung 404, Taiwan

⁴ Department of Applied Mathematics, National Sun Yat-sen University, Kaohsiung 804, Taiwan

Correspondence should be addressed to J. C. Yao, yaojc@math.nsysu.edu.tw

Received 20 July 2009; Accepted 27 October 2009

Recommended by Yeol Je Cho

The purpose of this paper is to introduce and study two new hybrid proximal-point algorithms for finding a common element of the set of solutions to a generalized equilibrium problem and the sets of zeros of two maximal monotone operators in a uniformly smooth and uniformly convex Banach space. We established strong and weak convergence theorems for these two modified hybrid proximal-point algorithms, respectively.

Copyright © 2009 L. C. Zeng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let X be a real Banach space and X^* its dual space. The normalized duality mapping $J : X \rightarrow 2^{X^*}$ is defined as

$$J(x) := \left\{ x^* \in X^* : \langle x^*, x \rangle = \|x\|^2 = \|x^*\|^2 \right\}, \quad \forall x \in X, \quad (1.1)$$

where $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing. Recall that if X is a smooth Banach space then J is singlevalued. Throughout this paper, we will still denote by J the single-valued normalized duality mapping. Let C be a nonempty closed convex subset of X , f a bifunction from $C \times C$ to \mathbb{R} , and $A : C \rightarrow X^*$ a nonlinear mapping. The generalized equilibrium problem is to find $\hat{x} \in C$ such that

$$f(\hat{x}, y) + \langle A\hat{x}, y - \hat{x} \rangle \geq 0, \quad \forall y \in C. \quad (1.2)$$

The set of solutions of (1.2) is denoted by EP . Problem (1.2) and similar problems have been extensively studied; see, for example, [1–11]. Whenever $A = 0$, problem (1.2) reduces to the equilibrium problem of finding $\hat{x} \in C$ such that

$$f(\hat{x}, y) \geq 0, \quad \forall y \in C. \quad (1.3)$$

The set of solutions of (1.3) is denoted by $EP(f)$. Whenever $f = 0$, problem (1.2) reduces to the variational inequality problem of finding $\hat{x} \in C$ such that

$$\langle A\hat{x}, y - \hat{x} \rangle \geq 0, \quad \forall y \in C. \quad (1.4)$$

Whenever $X = H$ a Hilbert space, problem (1.2) was very recently introduced and considered by Kamimura and Takahashi [12]. Problem (1.2) is very general in the sense that it includes, as special cases, optimization problems, variational inequalities, minimax problems, the Nash equilibrium problem in noncooperative games, and others; see, for example, [13, 14]. A mapping $S : C \rightarrow X$ is called nonexpansive if $\|Sx - Sy\| \leq \|x - y\|$ for all $x, y \in C$. Denote by $F(S)$ the set of fixed points of S , that is, $F(S) = \{x \in C : Sx = x\}$. Iterative schemes for finding common elements of EP and fixed points set of nonexpansive mappings have been studied recently; see, for example, [12, 15–17] and the references therein.

On the other hand, a classical method of solving $0 \in Tx$ in a Hilbert space H is the proximal point algorithm which generates, for any starting point $x_0 \in H$, a sequence $\{x_n\}$ in H by the iterative scheme

$$x_{n+1} = J_{r_n} x_n, \quad n = 0, 1, 2, \dots, \quad (1.5)$$

where $\{r_n\}$ is a sequence in $(0, \infty)$, $J_r = (I + rT)^{-1}$ for each $r > 0$ is the resolvent operator for T , and I is the identity operator on H . This algorithm was first introduced by Martinet [14] and generally studied by Rockafellar [18] in the framework of a Hilbert space H . Later many authors studied (1.5) and its variants in a Hilbert space H or in a Banach space X ; see, for example, [13, 19–23] and the references therein.

Let X be a uniformly smooth and uniformly convex Banach space and let C be a nonempty closed convex subset of X . Let f be a bifunction from $C \times C$ to \mathbb{R} satisfying the following conditions (A1)–(A4) which were imposed in [24]:

- (A1) $f(x, x) = 0$ for all $x \in C$;
- (A2) f is monotone, that is, $f(x, y) + f(y, x) \leq 0$, for all $x, y \in C$;
- (A3) for all $x, y, z \in C$, $\limsup_{t \downarrow 0} f(tz + (1-t)x, y) \leq f(x, y)$;
- (A4) for all $x \in C$, $f(x, \cdot)$ is convex and lower semicontinuous.

Let $T : X \rightarrow 2^{X^*}$ be a maximal monotone operator such that

- (A5) $T^{-1}0 \cap EP(f) \neq \emptyset$.

The purpose of this paper is to introduce and study two new iterative algorithms for finding a common element of the set EP of solutions for the generalized equilibrium problem (1.2) and the set $T^{-1}0 \cap \tilde{T}^{-1}0$ for maximal monotone operators T, \tilde{T} in a uniformly smooth and uniformly convex Banach space X . First, motivated by Kamimura and Takahashi

[12, Theorem 3.1], Ceng et al. [16, Theorem 3.1], and Zhang [17, Theorem 3.1], we introduce a sequence $\{x_n\}$ that, under some appropriate conditions, is strongly convergent to $\Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP}x_0$ in Section 3. Second, inspired by Kamimura and Takahashi [12, Theorem 3.1], Ceng et al. [16, Theorem 4.1], and Zhang [17, Theorem 3.1], we define a sequence weakly convergent to an element $z \in T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$, where $z = \lim_{n \rightarrow \infty} \Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP}x_n$ in Section 4. Our results represent a generalization of known results in the literature, including Takahashi and Zembayashi [15], Kamimura and Takahashi [12], Li and Song [22], Ceng and Yao [25], and Ceng et al. [16]. In particular, compared with Theorems 3.1 and 4.1 in [16], our results (i.e., Theorems 3.2 and 4.2 in this paper) extend the problem of finding an element of $T^{-1}0 \cap EP(f)$ to the one of finding an element of $T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$. Meantime, the algorithms in this paper are very different from those in [16] (because of considering the complexity involving the problem of finding an element of $T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$).

2. Preliminaries

In the sequel, we denote the strong convergence, weak convergence and weak* convergence of a sequence $\{x_n\}$ to a point $x \in X$ by $x_n \rightarrow x$, $x_n \rightharpoonup x$ and $x_n \xrightarrow{*} x$, respectively.

A Banach space X is said to be strictly convex, if $\|x + y\|/2 < 1$ for all $x, y \in U = \{z \in X : \|z\| = 1\}$ with $x \neq y$. X is said to be uniformly convex if for each $\epsilon \in (0, 2]$ there exists $\delta > 0$ such that $\|x + y\|/2 \leq 1 - \delta$ for all $x, y \in U$ with $\|x - y\| \geq \epsilon$. Recall that each uniformly convex Banach space has the Kadec-Klee property, that is,

$$\left. \begin{array}{l} x_n \rightarrow x \\ \|x_n\| \rightarrow \|x\| \end{array} \right\} \implies x_n \xrightarrow{*} x \quad (2.1)$$

The proof of the main results of Sections 3 and 4 will be based on the following assumption.

Assumption A. Let X be a uniformly smooth and uniformly convex Banach space and let C be a nonempty closed convex subset of X . Let f be a bifunction from $C \times C$ to R satisfying the same conditions (A1)–(A4) as in Section 1. Let $T, \tilde{T} : X \rightarrow 2^{X^*}$ be two maximal monotone operators such that

$$(A5)' \quad T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP \neq \emptyset.$$

Recall that if C is a nonempty closed convex subset of a Hilbert space H , then the metric projection $P_C : H \rightarrow C$ of H onto C is nonexpansive. This fact actually characterizes Hilbert spaces and hence, it is not available in more general Banach spaces. In this connection, Alber [26] recently introduced a generalized projection operator Π_C in a Banach space X which is an analogue of the metric projection in Hilbert spaces. Consider the functional defined as in [26] by

$$\phi(x, y) = \|x\|^2 - 2\langle x, Jy \rangle + \|y\|^2, \quad \forall x, y \in X. \quad (2.2)$$

It is clear that in a Hilbert space H , (2.2) reduces to $\phi(x, y) = \|x - y\|^2$, $\forall x, y \in H$.

The generalized projection $\Pi_C : X \rightarrow C$ is a mapping that assigns to an arbitrary point $x \in X$ the minimum point of the functional $\phi(y, x)$; that is, $\Pi_C x = \bar{x}$, where \bar{x} is the solution to the minimization problem

$$\phi(\bar{x}, x) = \min_{y \in C} \phi(y, x). \quad (2.3)$$

The existence and uniqueness of the operator Π_C follows from the properties of the functional $\phi(x, y)$ and strict monotonicity of the mapping J (see, e.g., [27]). In a Hilbert space, $\Pi_C = P_C$. From [26], in a smooth strictly convex and reflexive Banach space X , we have

$$(\|y\| - \|x\|)^2 \leq \phi(y, x) \leq (\|y\| + \|x\|)^2, \quad \forall x, y \in X. \quad (2.4)$$

Moreover, by the property of subdifferential of convex functions, we easily get the following inequality:

$$\phi(x, y) \leq \phi(x, J^{-1}(Jy + Jz)) - 2\langle y - x, Jz \rangle, \quad \forall x, y, z \in X. \quad (2.5)$$

Let S be a mapping from C into itself. A point p in C is called an asymptotically fixed point of S if C contains a sequence $\{x_n\}$ which converges weakly to p such that $Sx_n - x_n \rightarrow 0$ [28]. The set of asymptotically fixed points of S will be denoted by $\hat{F}(S)$. A mapping C from S into itself is called relatively nonexpansive if $\hat{F}(S) = F(S)$ and $\phi(p, Sx) \leq \phi(p, x)$, for all $x \in C$ and $p \in F(S)$ [15].

Observe that, if X is a reflexive strictly convex and smooth Banach space, then for any $x, y \in X$, $\phi(x, y) = 0$ if and only if $x = y$. To this end, it is sufficient to show that if $\phi(x, y) = 0$ then $x = y$. Actually, from (2.4), we have $\|x\| = \|y\|$ which implies that $\langle x, Jy \rangle = \|x\|^2 = \|y\|^2$. From the definition of J , we have $Jx = Jy$ and therefore, $x = y$; see [29] for more details.

We need the following lemmas for the proof of our main results.

Lemma 2.1 (Kamimura and Takahashi [12]). *Let X be a smooth and uniformly convex Banach space and let $\{x_n\}$ and $\{y_n\}$ be two sequences of X . If $\phi(x_n, y_n) \rightarrow 0$ and either $\{x_n\}$ or $\{y_n\}$ is bounded, then $x_n - y_n \rightarrow 0$.*

Lemma 2.2 (Alber [26], Kamimura and Takahashi [12]). *Let C be a nonempty closed convex subset of a smooth strictly convex and reflexive Banach space X . Let $x \in X$ and let $z \in C$. Then*

$$z = \Pi_C x \iff \langle y - z, Jx - Jz \rangle \leq 0, \quad \forall y \in C. \quad (2.6)$$

Lemma 2.3 (Alber [26], Kamimura and Takahashi [12]). *Let C be a nonempty closed convex subset of a smooth strictly convex and reflexive Banach space X . Then*

$$\phi(x, \Pi_C y) + \phi(\Pi_C y, y) \leq \phi(x, y), \quad \forall x \in C, y \in X. \quad (2.7)$$

Lemma 2.4 (Rockafellar [18]). *Let X be a reflexive strictly convex and smooth Banach space and let $T : X \rightarrow 2^{X^*}$ be a multivalued operator. Then there hold the following hold:*

- (i) $T^{-1}0$ is closed and convex if T is maximal monotone such that $T^{-1}0 \neq \emptyset$;
- (ii) T is maximal monotone if and only if T is monotone with $R(J + rT) = X^*$ for all $r > 0$.

Lemma 2.5 (Xu [30]). *Let X be a uniformly convex Banach space and let $r > 0$. Then there exists a strictly increasing, continuous, and convex function $g : [0, 2r] \rightarrow \mathbb{R}$ such that $g(0) = 0$ and*

$$\|tx + (1-t)y\|^2 \leq t\|x\|^2 + (1-t)\|y\|^2 - t(1-t)g(\|x-y\|), \quad (2.8)$$

for all $x, y \in B_r$ and $t \in [0, 1]$, where $B_r = \{z \in X : \|z\| \leq r\}$.

Lemma 2.6 (Kamimura and Takahashi [12]). *Let X be a smooth and uniformly convex Banach space and let $r > 0$. Then there exists a strictly increasing, continuous, and convex function $g : [0, 2r] \rightarrow \mathbb{R}$ such that $g(0) = 0$ and*

$$g(\|x-y\|) \leq \phi(x, y), \quad \forall x, y \in B_r. \quad (2.9)$$

The following result is due to Blum and Oettli [24].

Lemma 2.7 (Blum and Oettli [24]). *Let C be a nonempty closed convex subset of a smooth strictly convex and reflexive Banach space X , let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4), and let $r > 0$ and $x \in X$. Then, there exists $z \in C$ such that*

$$f(z, y) + \frac{1}{r} \langle y - z, Jz - Jx \rangle \geq 0, \quad \forall y \in C. \quad (2.10)$$

Motivated by Combettes and Hirstoaga [31] in a Hilbert space, Takahashi and Zembayashi [15] established the following lemma.

Lemma 2.8 (Takahashi and Zembayashi [15]). *Let C be a nonempty closed convex subset of a smooth strictly convex and reflexive Banach space X , and let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4). For $r > 0$ and $x \in X$, define a mapping $T_r : X \rightarrow C$ as follows:*

$$T_r(x) = \left\{ z \in C : f(z, y) + \frac{1}{r} \langle y - z, Jz - Jx \rangle \geq 0, \forall y \in C \right\} \quad (2.11)$$

for all $x \in X$. Then, the following hold:

- (i) T_r is singlevalued;
- (ii) T_r is a firmly nonexpansive-type mapping, that is, for all $x, y \in X$,

$$\langle T_r x - T_r y, J T_r x - J T_r y \rangle \leq \langle T_r x - T_r y, J x - J y \rangle; \quad (2.12)$$

- (iii) $F(T_r) = \widehat{F}(T_r) = EP(f)$;
- (iv) $EP(f)$ is closed and convex.

Using Lemma 2.8, one has the following result.

Lemma 2.9 (Takahashi and Zembayashi [15]). *Let C be a nonempty closed convex subset of a smooth strictly convex and reflexive Banach space X , let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4), and let $r > 0$. Then, for $x \in X$ and $q \in F(T_r)$,*

$$\phi(q, T_r x) + \phi(T_r x, x) \leq \phi(q, x). \quad (2.13)$$

Utilizing Lemmas 2.7, 2.8 and 2.9 as previously mentioned, Zhang [17] derived the following result.

Proposition 2.10 (Zhang [21, Lemma]). *Let X be a smooth strictly convex and reflexive Banach space and let C be a nonempty closed convex subset of X . Let $A : C \rightarrow X^*$ be an α -inverse-strongly monotone mapping, let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4), and let $r > 0$. Then the following hold:*

for $x \in X$, there exists $u \in C$ such that

$$f(u, y) + \langle Au, y - u \rangle + \frac{1}{r} \langle y - u, Ju - Jx \rangle \geq 0, \quad \forall y \in C, \quad (2.14)$$

if X is additionally uniformly smooth and $K_r : X \rightarrow C$ is defined as

$$K_r(x) = \left\{ u \in C : f(u, y) + \langle Au, y - u \rangle + \frac{1}{r} \langle y - u, Ju - Jx \rangle \geq 0, \forall y \in C \right\}, \quad \forall x \in X, \quad (2.15)$$

then the mapping K_r has the following properties:

- (i) K_r is singlevalued,
- (ii) K_r is a firmly nonexpansive-type mapping, that is,

$$\langle K_r x - K_r y, JK_r x - JK_r y \rangle \leq \langle K_r x - K_r y, Jx - Jy \rangle, \quad \forall x, y \in X, \quad (2.16)$$

- (iii) $F(K_r) = \widehat{F}(K_r) = EP$,
- (iv) EP is a closed convex subset of C ,
- (v) $\phi(p, K_r x) + \phi(K_r x, x) \leq \phi(p, x)$, for all $p \in F(K_r)$.

Let $T, \tilde{T} : X \rightarrow 2^{X^*}$ be two maximal monotone operators in a smooth Banach space X . We denote the resolvent operators of T and \tilde{T} by $J_r = (J + rT)^{-1}J$ and $\tilde{J}_r = (J + r\tilde{T})^{-1}J$ for each $r > 0$, respectively. Then $J_r : X \rightarrow D(T)$ and $\tilde{J}_r : X \rightarrow D(\tilde{T})$ are two single-valued mappings. Also, $T^{-1}0 = F(J_r)$ and $\tilde{T}^{-1}0 = F(\tilde{J}_r)$ for each $r > 0$, where $F(J_r)$ and $F(\tilde{J}_r)$ are the sets of fixed points of J_r and \tilde{J}_r , respectively. For each $r > 0$, the Yosida approximations of T and \tilde{T} are defined by $A_r = (J - JJ_r)/r$ and $\tilde{A}_r = (J - J\tilde{J}_r)/r$, respectively. It is known that

$$A_r x \in T(J_r x), \quad \tilde{A}_r x \in \tilde{T}(\tilde{J}_r x), \quad \forall r > 0, x \in X. \quad (2.17)$$

Lemma 2.11 (Kohsaka and Takahashi [13]). *Let X be a reflexive strictly convex and smooth Banach space and let $T : X \rightarrow 2^{X^*}$ be a maximal monotone operator with $T^{-1}0 \neq \emptyset$. Then*

$$\phi(z, J_r x) + \phi(J_r x, x) \leq \phi(z, x), \quad \forall r > 0, z \in T^{-1}0, x \in X. \quad (2.18)$$

Lemma 2.12 (Tan and Xu [32]). *Let $\{a_n\}$ and $\{b_n\}$ be two sequences of nonnegative real numbers satisfying the inequality: $a_{n+1} \leq a_n + b_n$ for all $n \geq 0$. If $\sum_{n=0}^{\infty} b_n < \infty$, then $\lim_{n \rightarrow \infty} a_n$ exists.*

3. Strong Convergence Theorem

In this section, we prove a strong convergence theorem for finding a common element of the set of solutions for a generalized equilibrium problem and the set $T^{-1}0 \cap \tilde{T}^{-1}0$ for two maximal monotone operators T and \tilde{T} .

Lemma 3.1. *Let X be a reflexive strictly convex and smooth Banach space and let $T : X \rightarrow 2^{X^*}$ be a maximal monotone operator. Then for each $r \in (0, \infty)$, the following holds:*

$$\langle Ju - Jv, J_r u - J_r v \rangle \geq \langle JJ_r u - JJ_r v, J_r u - J_r v \rangle, \quad \forall u, v \in X, \quad (3.1)$$

where $J_r = (J + rT)^{-1}J$ and J is the duality mapping on X . In particular, whenever $X = H$ a real Hilbert space, J_r is a nonexpansive mapping on H .

Proof. Since for each $u, v \in X$

$$J_r u = (J + rT)^{-1}Ju, \quad J_r v = (J + rT)^{-1}Jv, \quad (3.2)$$

we have that

$$\frac{1}{r} \cdot (Ju - JJ_r u) \in TJ_r u, \quad \frac{1}{r} \cdot (Jv - JJ_r v) \in TJ_r v. \quad (3.3)$$

Thus, from the monotonicity of T it follows that

$$\left\langle \frac{1}{r} \cdot (Ju - JJ_r u) - \frac{1}{r} \cdot (Jv - JJ_r v), J_r u - J_r v \right\rangle \geq 0, \quad (3.4)$$

and hence

$$\langle Ju - Jv, J_r u - J_r v \rangle \geq \langle JJ_r u - JJ_r v, J_r u - J_r v \rangle. \quad (3.5)$$

□

Theorem 3.2. *Suppose that Assumption A is fulfilled and let $x_0 \in X$ be chosen arbitrarily. Consider the sequence*

$$x_{n+1} = \Pi_{H_n \cap W_n} x_0, \quad n = 0, 1, 2, \dots, \quad (3.6)$$

where

$$\begin{aligned}
 H_n &= \left\{ z \in C : \phi(z, K_{r_n} y_n) \leq \left[\alpha_n + \tilde{\beta}_n - \alpha_n \tilde{\beta}_n - \tilde{\alpha}_n \tilde{\beta}_n + \alpha_n \tilde{\alpha}_n \tilde{\beta}_n \right] \phi(z, x_0) \right. \\
 &\quad \left. + \left[(1 - \alpha_n)(1 - \tilde{\alpha}_n)(1 - \tilde{\beta}_n) + \tilde{\alpha}_n(1 - \alpha_n) \right] \phi(z, x_n) \right\}, \\
 W_n &= \{ z \in C : \langle x_n - z, Jx_0 - Jx_n \rangle \geq 0 \}, \\
 \tilde{x}_n &= J^{-1}(\alpha_n Jx_0 + (1 - \alpha_n)(\beta_n Jx_n + (1 - \beta_n)JJ_{r_n}x_n)), \\
 y_n &= J^{-1}(\tilde{\alpha}_n J\tilde{x}_n + (1 - \tilde{\alpha}_n)J\tilde{J}_{r_n}J^{-1}(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{x}_n)),
 \end{aligned} \tag{3.7}$$

K_r is defined by (2.15), $\{\alpha_n\}, \{\beta_n\}, \{\tilde{\alpha}_n\}, \{\tilde{\beta}_n\} \subset [0, 1]$ satisfy

$$\lim_{n \rightarrow \infty} \alpha_n = 0, \quad \lim_{n \rightarrow \infty} \tilde{\beta}_n = 0, \quad \liminf_{n \rightarrow \infty} \beta_n(1 - \beta_n) > 0, \quad \liminf_{n \rightarrow \infty} \tilde{\alpha}_n(1 - \tilde{\alpha}_n) > 0, \tag{3.8}$$

and $\{r_n\} \subset (0, \infty)$ satisfies $\liminf_{n \rightarrow \infty} r_n > 0$. Then, the sequence $\{x_n\}$ converges strongly to $\Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP} x_0$ provided $\|\tilde{J}_{r_n} v_n - \tilde{J}_{r_n} \tilde{x}_n\| \rightarrow 0$ for any sequence $\{v_n\} \subset X$ with $\|v_n - \tilde{x}_n\| \rightarrow 0$, where $\Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP}$ is the generalized projection of X onto $T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$.

Remark 3.3. In Theorem 3.2, if $X = H$ a real Hilbert space, then $\{\tilde{J}_{r_n}\}$ is a sequence of nonexpansive mappings on H . This implies that as $n \rightarrow \infty$,

$$\left\| \tilde{J}_{r_n} v_n - \tilde{J}_{r_n} \tilde{x}_n \right\| \leq \|v_n - \tilde{x}_n\| \rightarrow 0. \tag{3.9}$$

In this case, we can remove the requirement that $\|\tilde{J}_{r_n} v_n - \tilde{J}_{r_n} \tilde{x}_n\| \rightarrow 0$ for any sequence $\{v_n\} \subset X$ with $\|v_n - \tilde{x}_n\| \rightarrow 0$.

Proof of Theorem 3.2. For the sake of simplicity, we define

$$u_n := K_{r_n} y_n, \quad z_n := J^{-1}(\beta_n Jx_n + (1 - \beta_n)JJ_{r_n}x_n), \quad \tilde{z}_n := \tilde{J}_{r_n} J^{-1}(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{x}_n), \tag{3.10}$$

so that

$$\tilde{x}_n = J^{-1}(\alpha_n Jx_0 + (1 - \alpha_n)Jz_n), \quad y_n = J^{-1}(\tilde{\alpha}_n J\tilde{x}_n + (1 - \tilde{\alpha}_n)J\tilde{z}_n). \tag{3.11}$$

We divide the proof into several steps.

Step 1. We claim that $H_n \cap W_n$ is closed and convex for each $n \geq 0$.

Indeed, it is obvious that H_n is closed and W_n is closed and convex for each $n \geq 0$. Let us show that H_n is convex. For $z_1, z_2 \in H_n$ and $t \in (0, 1)$, put $z = tz_1 + (1 - t)z_2$. It is sufficient

to show that $z \in H_n$. We first write $\gamma_n = \alpha_n + \tilde{\beta}_n - \alpha_n\tilde{\beta}_n - \tilde{\alpha}_n\tilde{\beta}_n + \alpha_n\tilde{\alpha}_n\tilde{\beta}_n$ for each $n \geq 0$. Next, we prove that

$$\phi(z, u_n) \leq \gamma_n\phi(z, x_0) + (1 - \gamma_n)\phi(z, x_n) \quad (3.12)$$

is equivalent to

$$2\gamma_n\langle z, Jx_0 \rangle + 2(1 - \gamma_n)\langle z, Jx_n \rangle - 2\langle z, Ju_n \rangle \leq \gamma_n\|x_0\|^2 + (1 - \gamma_n)\|x_n\|^2 - \|u_n\|^2. \quad (3.13)$$

Indeed, from (2.4) we deduce that the following equations hold:

$$\begin{aligned} \phi(z, x_0) &= \|z\|^2 - 2\langle z, Jx_0 \rangle + \|x_0\|^2, \\ \phi(z, x_n) &= \|z\|^2 - 2\langle z, Jx_n \rangle + \|x_n\|^2, \\ \phi(z, u_n) &= \|z\|^2 - 2\langle z, Ju_n \rangle + \|u_n\|^2, \end{aligned} \quad (3.14)$$

which combined with (3.12) yield that (3.12) is equivalent to (3.13). Thus we have

$$\begin{aligned} &2\gamma_n\langle z, Jx_0 \rangle + 2(1 - \gamma_n)\langle z, Jx_n \rangle - 2\langle z, Ju_n \rangle \\ &= 2\gamma_n\langle tz_1 + (1 - t)z_2, Jx_0 \rangle \\ &\quad + 2(1 - \gamma_n)\langle tz_1 + (1 - t)z_2, Jx_n \rangle \\ &\quad - 2\langle tz_1 + (1 - t)z_2, Ju_n \rangle \\ &= 2t\gamma_n\langle z_1, Jx_0 \rangle + 2(1 - t)\gamma_n\langle z_2, Jx_0 \rangle + 2(1 - \gamma_n)t\langle z_1, Jx_n \rangle \\ &\quad + 2(1 - \gamma_n)(1 - t)\langle z_2, Jx_n \rangle - 2t\langle z_1, Ju_n \rangle - 2(1 - t)\langle z_2, Ju_n \rangle \\ &\leq \gamma_n\|x_0\|^2 + (1 - \gamma_n)\|x_n\|^2 - \|u_n\|^2. \end{aligned} \quad (3.15)$$

This implies that $z \in H_n$. Therefore, H_n is closed and convex.

Step 2. We claim that $T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP \subset H_n \cap W_n$ for each $n \geq 0$ and that $\{x_n\}$ is well defined. Indeed, take $w \in T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$ arbitrarily. Note that $u_n = K_{r_n}y_n$ is equivalent to

$$u_n \in C \quad \text{such that} \quad f(u_n, y) + \langle Au_n, y - u_n \rangle + \frac{1}{r_n}\langle y - u_n, Ju_n - Jy_n \rangle \geq 0, \quad \forall y \in C. \quad (3.16)$$

Then from Lemma 2.11 we obtain

$$\begin{aligned}
 \phi(w, z_n) &= \phi\left(w, J^{-1}(\beta_n Jx_n + (1 - \beta_n)JJ_{r_n}x_n)\right) \\
 &= \|w\|^2 - 2\langle w, \beta_n Jx_n + (1 - \beta_n)JJ_{r_n}x_n \rangle + \|\beta_n Jx_n + (1 - \beta_n)JJ_{r_n}x_n\|^2 \\
 &\leq \|w\|^2 - 2\beta_n\langle w, Jx_n \rangle - 2(1 - \beta_n)\langle w, JJ_{r_n}x_n \rangle + \beta_n\|x_n\|^2 + (1 - \beta_n)\|J_{r_n}x_n\|^2 \quad (3.17) \\
 &= \beta_n\phi(w, x_n) + (1 - \beta_n)\phi(w, J_{r_n}x_n) \\
 &\leq \beta_n\phi(w, x_n) + (1 - \beta_n)\phi(w, x_n) = \phi(w, x_n),
 \end{aligned}$$

$$\begin{aligned}
 \phi(w, \tilde{x}_n) &= \phi\left(w, J^{-1}(\alpha_n Jx_0 + (1 - \alpha_n)Jz_n)\right) \\
 &= \|w\|^2 - 2\langle w, \alpha_n Jx_0 + (1 - \alpha_n)Jz_n \rangle + \|\alpha_n Jx_0 + (1 - \alpha_n)Jz_n\|^2 \\
 &\leq \|w\|^2 - 2\alpha_n\langle w, Jx_0 \rangle - 2(1 - \alpha_n)\langle w, Jz_n \rangle + \alpha_n\|x_0\|^2 + (1 - \alpha_n)\|z_n\|^2 \quad (3.18) \\
 &= \alpha_n\phi(w, x_0) + (1 - \alpha_n)\phi(w, z_n) \\
 &\leq \alpha_n\phi(w, x_0) + (1 - \alpha_n)\phi(w, x_n).
 \end{aligned}$$

Moreover, we have

$$\begin{aligned}
 \phi(w, \tilde{z}_n) &= \phi\left(w, \tilde{J}_{r_n}J^{-1}(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{x}_n)\right) \\
 &\leq \phi\left(w, J^{-1}(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{x}_n)\right) \\
 &= \|w\|^2 - 2\langle w, \tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{x}_n \rangle + \|\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{x}_n\|^2 \\
 &\leq \|w\|^2 - 2\tilde{\beta}_n\langle w, Jx_0 \rangle - 2(1 - \tilde{\beta}_n)\langle w, J\tilde{x}_n \rangle + \tilde{\beta}_n\|x_0\|^2 + (1 - \tilde{\beta}_n)\|\tilde{x}_n\|^2 \\
 &= \tilde{\beta}_n\phi(w, x_0) + (1 - \tilde{\beta}_n)\phi(w, \tilde{x}_n) \\
 &\leq \tilde{\beta}_n\phi(w, x_0) + (1 - \tilde{\beta}_n)[\alpha_n\phi(w, x_0) + (1 - \alpha_n)\phi(w, x_n)] \\
 &= [\tilde{\beta}_n + (1 - \tilde{\beta}_n)\alpha_n]\phi(w, x_0) + (1 - \tilde{\beta}_n)(1 - \alpha_n)\phi(w, x_n), \\
 \phi(w, y_n) &= \phi\left(w, J^{-1}(\tilde{\alpha}_n J\tilde{x}_n + (1 - \tilde{\alpha}_n)J\tilde{z}_n)\right) \\
 &\leq \|w\|^2 - 2\tilde{\alpha}_n\langle w, J\tilde{x}_n \rangle - 2(1 - \tilde{\alpha}_n)\langle w, J\tilde{z}_n \rangle + \tilde{\alpha}_n\|\tilde{x}_n\|^2 + (1 - \tilde{\alpha}_n)\|\tilde{z}_n\|^2
 \end{aligned}$$

$$\begin{aligned}
&= \tilde{\alpha}_n \phi(w, \tilde{x}_n) + (1 - \tilde{\alpha}_n) \phi(w, \tilde{z}_n) \\
&\leq \tilde{\alpha}_n [\alpha_n \phi(w, x_0) + (1 - \alpha_n) \phi(w, x_n)] \\
&\quad + (1 - \tilde{\alpha}_n) \left\{ [\tilde{\beta}_n + (1 - \tilde{\beta}_n) \alpha_n] \phi(w, x_0) + (1 - \tilde{\beta}_n) (1 - \alpha_n) \phi(w, x_n) \right\} \\
&= [\alpha_n + \tilde{\beta}_n - \alpha_n \tilde{\beta}_n - \tilde{\alpha}_n \tilde{\beta}_n + \alpha_n \tilde{\alpha}_n \tilde{\beta}_n] \phi(w, x_0) \\
&\quad + [(1 - \alpha_n) (1 - \tilde{\alpha}_n) (1 - \tilde{\beta}_n) + \tilde{\alpha}_n (1 - \alpha_n)] \phi(w, x_n) \\
&= \gamma_n \phi(w, x_0) + (1 - \gamma_n) \phi(w, x_n), \tag{3.19}
\end{aligned}$$

where $\gamma_n = \alpha_n + \tilde{\beta}_n - \alpha_n \tilde{\beta}_n - \tilde{\alpha}_n \tilde{\beta}_n + \alpha_n \tilde{\alpha}_n \tilde{\beta}_n$. So $w \in H_n$ for all $n \geq 0$. Now, let us show that

$$T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP \subset W_n \quad \forall n \geq 0. \tag{3.20}$$

We prove this by induction. For $n = 0$, we have $T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP \subset C = W_0$. Assume that $T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP \subset W_n$. Since x_{n+1} is the projection of x_0 onto $H_n \cap W_n$, by Lemma 2.2 we have

$$\langle x_{n+1} - z, Jx_0 - Jx_{n+1} \rangle \geq 0, \quad \forall z \in H_n \cap W_n. \tag{3.21}$$

As $T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP \subset H_n \cap W_n$ by the induction assumption, the last inequality holds, in particular, for all $z \in T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$. This, together with the definition of W_{n+1} implies that $T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP \subset W_{n+1}$. Hence (3.20) holds for all $n \geq 0$. So, $T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP \subset H_n \cap W_n$ for all $n \geq 0$. This implies that the sequence $\{x_n\}$ is well defined.

Step 3. We claim that $\{x_n\}$ is bounded and that $\phi(x_{n+1}, x_n) \rightarrow 0$ as $n \rightarrow \infty$.

Indeed, it follows from the definition of W_n that $x_n = \Pi_{W_n} x_0$. Since $x_n = \Pi_{W_n} x_0$ and $x_{n+1} = \Pi_{H_n \cap W_n} x_0 \in W_n$, so $\phi(x_n, x_0) \leq \phi(x_{n+1}, x_0)$ for all $n \geq 0$; that is, $\{\phi(x_n, x_0)\}$ is nondecreasing. It follows from $x_n = \Pi_{W_n} x_0$ and Lemma 2.3 that

$$\phi(x_n, x_0) = \phi(\Pi_{W_n} x_0, x_0) \leq \phi(p, x_0) - \phi(p, x_n) \leq \phi(p, x_0) \tag{3.22}$$

for each $p \in T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP \subset W_n$ for each $n \geq 0$. Therefore, $\{\phi(x_n, x_0)\}$ is bounded which implies that the limit of $\{\phi(x_n, x_0)\}$ exists. Since

$$(\|x_n\| - \|x_0\|)^2 \leq \phi(x_n, x_0) \leq (\|x_n\| + \|x_0\|)^2, \quad \forall n \geq 0, \tag{3.23}$$

so $\{x_n\}$ is bounded. From Lemma 2.3, we have

$$\begin{aligned}\phi(x_{n+1}, x_n) &= \phi(x_{n+1}, \Pi_{W_n} x_0) \leq \phi(x_{n+1}, x_0) - \phi(\Pi_{W_n} x_0, x_0) \\ &= \phi(x_{n+1}, x_0) - \phi(x_n, x_0),\end{aligned}\tag{3.24}$$

for each $n \geq 0$. This implies that

$$\lim_{n \rightarrow \infty} \phi(x_{n+1}, x_n) = 0.\tag{3.25}$$

Step 4. We claim that $\lim_{n \rightarrow \infty} \|x_n - u_n\| = 0$, $\lim_{n \rightarrow \infty} \|x_n - J_{r_n} x_n\| = 0$, and $\lim_{n \rightarrow \infty} \|\tilde{x}_n - \tilde{J}_{r_n} \tilde{x}_n\| = 0$.

Indeed, from $x_{n+1} = \Pi_{H_n \cap W_n} x_0 \in H_n$, we have

$$\begin{aligned}\phi(x_{n+1}, u_n) &\leq \left[\alpha_n + \tilde{\beta}_n - \alpha_n \tilde{\beta}_n - \tilde{\alpha}_n \tilde{\beta}_n + \alpha_n \tilde{\alpha}_n \tilde{\beta}_n \right] \phi(x_{n+1}, x_0) \\ &\quad + \left[(1 - \alpha_n)(1 - \tilde{\alpha}_n)(1 - \tilde{\beta}_n) + \tilde{\alpha}_n(1 - \alpha_n) \right] \phi(x_{n+1}, x_n)\end{aligned}\tag{3.26}$$

for all $n \geq 0$. Therefore, from $\alpha_n \rightarrow 0$, $\tilde{\beta}_n \rightarrow 0$ and $\phi(x_{n+1}, x_n) \rightarrow 0$, it follows that $\lim_{n \rightarrow \infty} \phi(x_{n+1}, u_n) = 0$. Since $\lim_{n \rightarrow \infty} \phi(x_{n+1}, x_n) = \lim_{n \rightarrow \infty} \phi(x_{n+1}, u_n) = 0$ and X is uniformly convex and smooth, we have from Lemma 2.1 that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} \|x_{n+1} - u_n\| = 0,\tag{3.27}$$

and therefore, $\lim_{n \rightarrow \infty} \|x_n - u_n\| = 0$. Since J is uniformly norm-to-norm continuous on bounded subsets of X and $x_n - u_n \rightarrow 0$, then $\lim_{n \rightarrow \infty} \|Jx_n - Ju_n\| = 0$.

Let us set $\Omega := T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$. Then, according to Lemma 2.4 and Proposition 2.10, we know that Ω is a nonempty closed convex subset of X such that $\Omega \subset C$. Fix $u \in \Omega$ arbitrarily. As in the proof of Step 2 we can show that $\phi(u, z_n) \leq \phi(u, x_n)$, $\phi(u, \tilde{x}_n) \leq \alpha_n \phi(u, x_0) + (1 - \alpha_n) \phi(u, x_n)$, $\phi(u, \tilde{z}_n) \leq \tilde{\beta}_n \phi(u, x_0) + (1 - \tilde{\beta}_n) \phi(u, \tilde{x}_n)$, $\phi(u, y_n) \leq \tilde{\alpha}_n \phi(u, \tilde{x}_n) + (1 - \tilde{\alpha}_n) \phi(u, \tilde{z}_n)$, and $\phi(u, u_n) \leq \tilde{\alpha}_n \phi(u, \tilde{x}_n) + (1 - \tilde{\alpha}_n) \phi(u, \tilde{z}_n)$. Hence it follows from the boundedness of $\{x_n\}$ that $\{z_n\}$, $\{\tilde{x}_n\}$, $\{\tilde{z}_n\}$, $\{y_n\}$, and $\{u_n\}$ are also bounded. Let $r = \sup\{\|x_n\|, \|\tilde{x}_n\|, \|J_{r_n} x_n\|, \|\tilde{z}_n\| : n \geq 0\}$. Since X is a uniformly smooth Banach space, we know that X^* is a uniformly convex Banach space. Therefore, by Lemma 2.5 there exists a continuous, strictly increasing, and convex function g with $g(0) = 0$ such that

$$\|\alpha x^* + (1 - \alpha)y^*\|^2 \leq \alpha \|x^*\|^2 + (1 - \alpha) \|y^*\|^2 - \alpha(1 - \alpha)g(\|x^* - y^*\|),\tag{3.28}$$

for $x^*, y^* \in B_r^*$ and $\alpha \in [0, 1]$. So, we have that

$$\begin{aligned}
 \phi(u, z_n) &= \phi\left(u, J^{-1}(\beta_n Jx_n + (1 - \beta_n) JJ_{r_n} x_n)\right) \\
 &= \|u\|^2 - 2\langle u, \beta_n Jx_n + (1 - \beta_n) JJ_{r_n} x_n \rangle + \|\beta_n Jx_n + (1 - \beta_n) JJ_{r_n} x_n\|^2 \\
 &\leq \|u\|^2 - 2\beta_n \langle u, Jx_n \rangle - 2(1 - \beta_n) \langle u, JJ_{r_n} x_n \rangle \\
 &\quad + \beta_n \|x_n\|^2 + (1 - \beta_n) \|J_{r_n} x_n\|^2 - \beta_n(1 - \beta_n) g(\|Jx_n - JJ_{r_n} x_n\|) \\
 &= \beta_n \phi(u, x_n) + (1 - \beta_n) \phi(u, J_{r_n} x_n) - \beta_n(1 - \beta_n) g(\|Jx_n - JJ_{r_n} x_n\|) \\
 &\leq \beta_n \phi(u, x_n) + (1 - \beta_n) \phi(u, x_n) - \beta_n(1 - \beta_n) g(\|Jx_n - JJ_{r_n} x_n\|) \\
 &= \phi(u, x_n) - \beta_n(1 - \beta_n) g(\|Jx_n - JJ_{r_n} x_n\|),
 \end{aligned} \tag{3.29}$$

$$\begin{aligned}
 \phi(u, \tilde{z}_n) &= \phi\left(u, \tilde{J}_{r_n} J^{-1}(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n) J\tilde{x}_n)\right) \\
 &\leq \phi\left(u, J^{-1}(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n) J\tilde{x}_n)\right) \\
 &\leq \|u\|^2 - 2\langle u, \tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n) J\tilde{x}_n \rangle + \tilde{\beta}_n \|x_0\|^2 + (1 - \tilde{\beta}_n) \|\tilde{x}_n\|^2 \\
 &= \tilde{\beta}_n \phi(u, x_0) + (1 - \tilde{\beta}_n) \phi(u, \tilde{x}_n) \\
 &\leq \tilde{\beta}_n \phi(u, x_0) + (1 - \tilde{\beta}_n) [\phi(u, x_n) - \beta_n(1 - \beta_n) g(\|Jx_n - JJ_{r_n} x_n\|)] \\
 &= \tilde{\beta}_n \phi(u, x_0) + (1 - \tilde{\beta}_n) \phi(u, x_n) - (1 - \tilde{\beta}_n) \beta_n(1 - \beta_n) g(\|Jx_n - JJ_{r_n} x_n\|),
 \end{aligned} \tag{3.30}$$

and hence

$$\begin{aligned}
 \phi(u, \tilde{x}_n) &= \phi\left(u, J^{-1}(\alpha_n Jx_0 + (1 - \alpha_n) Jz_n)\right) \\
 &= \|u\|^2 - 2\langle u, \alpha_n Jx_0 + (1 - \alpha_n) Jz_n \rangle + \|\alpha_n Jx_0 + (1 - \alpha_n) Jz_n\|^2 \\
 &\leq \|u\|^2 - 2\alpha_n \langle u, Jx_0 \rangle - 2(1 - \alpha_n) \langle u, Jz_n \rangle + \alpha_n \|x_0\|^2 + (1 - \alpha_n) \|z_n\|^2 \\
 &= \alpha_n \phi(u, x_0) + (1 - \alpha_n) \phi(u, z_n) \\
 &\leq \alpha_n \phi(u, x_0) + (1 - \alpha_n) [\phi(u, x_n) - \beta_n(1 - \beta_n) g(\|Jx_n - JJ_{r_n} x_n\|)] \\
 &= \alpha_n \phi(u, x_0) + (1 - \alpha_n) \phi(u, x_n) - (1 - \alpha_n) \beta_n(1 - \beta_n) g(\|Jx_n - JJ_{r_n} x_n\|),
 \end{aligned} \tag{3.31}$$

$$\phi(u, u_n) = \phi(u, K_{r_n} y_n) \leq \phi(u, y_n) \quad (\text{using Proposition 2.10})$$

$$= \phi\left(u, J^{-1}(\tilde{\alpha}_n J\tilde{x}_n + (1 - \tilde{\alpha}_n) J\tilde{z}_n)\right)$$

$$\begin{aligned}
&= \|u\|^2 - 2\langle u, \tilde{\alpha}_n J\tilde{x}_n + (1 - \tilde{\alpha}_n)J\tilde{z}_n \rangle + \|\tilde{\alpha}_n J\tilde{x}_n + (1 - \tilde{\alpha}_n)J\tilde{z}_n\|^2 \\
&\leq \|u\|^2 - 2\tilde{\alpha}_n \langle u, J\tilde{x}_n \rangle - 2(1 - \tilde{\alpha}_n) \langle u, J\tilde{z}_n \rangle + \tilde{\alpha}_n \|\tilde{x}_n\|^2 + (1 - \tilde{\alpha}_n) \|\tilde{z}_n\|^2 \\
&\quad - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{x}_n - J\tilde{z}_n\|) \\
&= \tilde{\alpha}_n \phi(u, \tilde{x}_n) + (1 - \tilde{\alpha}_n) \phi(u, \tilde{z}_n) - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{x}_n - J\tilde{z}_n\|) \\
&\leq \tilde{\alpha}_n [\alpha_n \phi(u, x_0) + (1 - \alpha_n) \phi(u, x_n) - (1 - \alpha_n)\beta_n(1 - \beta_n)g(\|Jx_n - JJ_{r_n}x_n\|)] \\
&\quad + (1 - \tilde{\alpha}_n) [\tilde{\beta}_n \phi(u, x_0) + (1 - \tilde{\beta}_n) \phi(u, x_n) \\
&\quad - (1 - \tilde{\beta}_n)\beta_n(1 - \beta_n)g(\|Jx_n - JJ_{r_n}x_n\|)] - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{x}_n - J\tilde{z}_n\|) \\
&\leq (\alpha_n + \tilde{\beta}_n) \phi(u, x_0) + [\tilde{\alpha}_n(1 - \alpha_n) + (1 - \tilde{\alpha}_n)(1 - \tilde{\beta}_n)] \phi(u, x_n) \\
&\quad - [\tilde{\alpha}_n(1 - \alpha_n) + (1 - \tilde{\alpha}_n)(1 - \tilde{\beta}_n)] \beta_n(1 - \beta_n)g(\|Jx_n - JJ_{r_n}x_n\|) \\
&\quad - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{x}_n - J\tilde{z}_n\|) \\
&= (\alpha_n + \tilde{\beta}_n) \phi(u, x_0) + [1 - \alpha_n \tilde{\alpha}_n - \tilde{\beta}_n(1 - \tilde{\alpha}_n)] \phi(u, x_n) \\
&\quad - [1 - \alpha_n \tilde{\alpha}_n - \tilde{\beta}_n(1 - \tilde{\alpha}_n)] \beta_n(1 - \beta_n)g(\|Jx_n - JJ_{r_n}x_n\|) \\
&\quad - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{x}_n - J\tilde{z}_n\|) \\
&\leq (\alpha_n + \tilde{\beta}_n) \phi(u, x_0) + \phi(u, x_n) \\
&\quad - [1 - \alpha_n \tilde{\alpha}_n - \tilde{\beta}_n(1 - \tilde{\alpha}_n)] \beta_n(1 - \beta_n)g(\|Jx_n - JJ_{r_n}x_n\|) \\
&\quad - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{x}_n - J\tilde{z}_n\|), \tag{3.32}
\end{aligned}$$

for all $n \geq 0$. Consequently we have

$$\begin{aligned}
&[1 - \alpha_n \tilde{\alpha}_n - \tilde{\beta}_n(1 - \tilde{\alpha}_n)] \beta_n(1 - \beta_n)g(\|Jx_n - JJ_{r_n}x_n\|) + \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{x}_n - J\tilde{z}_n\|) \\
&\leq (\alpha_n + \tilde{\beta}_n) \phi(u, x_0) + \phi(u, x_n) - \phi(u, u_n) \\
&= (\alpha_n + \tilde{\beta}_n) \phi(u, x_0) + \|x_n\|^2 - \|u_n\|^2 - 2\langle u, Jx_n - Ju_n \rangle \\
&\leq (\alpha_n + \tilde{\beta}_n) \phi(u, x_0) + \left| \|x_n\|^2 - \|u_n\|^2 \right| + 2|\langle u, Jx_n - Ju_n \rangle| \\
&\leq (\alpha_n + \tilde{\beta}_n) \phi(u, x_0) + \|x_n\| - \|u_n\|(\|x_n\| + \|u_n\|) + 2\|u\| \|Jx_n - Ju_n\| \\
&\leq (\alpha_n + \tilde{\beta}_n) \phi(u, x_0) + \|x_n - u_n\|(\|x_n\| + \|u_n\|) + 2\|u\| \|Jx_n - Ju_n\|. \tag{3.33}
\end{aligned}$$

Since $x_n - u_n \rightarrow 0$ and J is uniformly norm-to-norm continuous on bounded subsets of X , we obtain $Jx_n - Ju_n \rightarrow 0$. From $\liminf_{n \rightarrow \infty} \beta_n(1 - \beta_n) > 0$, $\liminf_{n \rightarrow \infty} \tilde{\alpha}_n(1 - \tilde{\alpha}_n) > 0$, and $\lim_{n \rightarrow \infty} \alpha_n = \lim_{n \rightarrow \infty} \tilde{\beta}_n = 0$ we have

$$\lim_{n \rightarrow \infty} g(\|Jx_n - JJ_{r_n}x_n\|) = \lim_{n \rightarrow \infty} g(\|J\tilde{x}_n - J\tilde{z}_n\|) = 0. \quad (3.34)$$

Therefore, from the properties of g we get

$$\begin{aligned} \lim_{n \rightarrow \infty} \|Jx_n - JJ_{r_n}x_n\| &= \lim_{n \rightarrow \infty} \|x_n - J_{r_n}x_n\| = 0, \\ \lim_{n \rightarrow \infty} \|J\tilde{x}_n - J\tilde{z}_n\| &= \lim_{n \rightarrow \infty} \|\tilde{x}_n - \tilde{z}_n\| = 0, \end{aligned} \quad (3.35)$$

recalling that J^{-1} is uniformly norm-to-norm continuous on bounded subsets of X^* . Next let us show that

$$\lim_{n \rightarrow \infty} \|J\tilde{x}_n - J\tilde{J}_{r_n}\tilde{x}_n\| = \lim_{n \rightarrow \infty} \|\tilde{x}_n - \tilde{J}_{r_n}\tilde{x}_n\| = 0. \quad (3.36)$$

Observe first that

$$\begin{aligned} \phi(u_n, x_n) - \phi(x_{n+1}, u_n) &= \|x_n\|^2 - \|x_{n+1}\|^2 - 2\langle u_n, Jx_n \rangle + 2\langle x_{n+1}, Ju_n \rangle \\ &= (\|x_n\| - \|x_{n+1}\|)(\|x_n\| + \|x_{n+1}\|) + 2\langle x_{n+1} - u_n, Jx_n \rangle + 2\langle x_{n+1}, Ju_n - Jx_n \rangle \\ &\leq \|x_n - x_{n+1}\|(\|x_n\| + \|x_{n+1}\|) + 2\|x_{n+1} - u_n\|\|x_n\| + 2\|x_{n+1}\|\|Ju_n - Jx_n\|. \end{aligned} \quad (3.37)$$

Since $\phi(x_{n+1}, u_n) \rightarrow 0$, $\|x_{n+1} - x_n\| \rightarrow 0$, $\|x_{n+1} - u_n\| \rightarrow 0$, $\|Ju_n - Jx_n\| \rightarrow 0$, and $\{x_n\}$ is bounded, so it follows that $\phi(u_n, x_n) \rightarrow 0$. Also, observe that

$$\begin{aligned} \phi(u_n, J_{r_n}x_n) - \phi(u_n, x_n) &= \|J_{r_n}x_n\|^2 - \|x_n\|^2 + 2\langle u_n, Jx_n - JJ_{r_n}x_n \rangle \\ &= (\|J_{r_n}x_n\| - \|x_n\|)(\|J_{r_n}x_n\| + \|x_n\|) + 2\langle u_n, Jx_n - JJ_{r_n}x_n \rangle \\ &\leq \|J_{r_n}x_n - x_n\|(\|J_{r_n}x_n\| + \|x_n\|) + 2\|u_n\|\|Jx_n - JJ_{r_n}x_n\|. \end{aligned} \quad (3.38)$$

Since $\phi(u_n, x_n) \rightarrow 0$, $\|J_{r_n}x_n - x_n\| \rightarrow 0$, $\|Jx_n - JJ_{r_n}x_n\| \rightarrow 0$ and the sequences $\{x_n\}$, $\{u_n\}$, $\{J_{r_n}x_n\}$ are bounded, so it follows that $\phi(u_n, J_{r_n}x_n) \rightarrow 0$. Meantime, observe that

$$\begin{aligned}\phi(u_n, z_n) &= \phi\left(u_n, J^{-1}(\beta_n Jx_n + (1 - \beta_n)JJ_{r_n}x_n)\right) \\ &= \|u_n\|^2 - 2\langle u_n, \beta_n Jx_n + (1 - \beta_n)JJ_{r_n}x_n \rangle + \|\beta_n Jx_n + (1 - \beta_n)JJ_{r_n}x_n\|^2 \\ &\leq \|u_n\|^2 - 2\beta_n\langle u_n, Jx_n \rangle - 2(1 - \beta_n)\langle u_n, JJ_{r_n}x_n \rangle + \beta_n\|x_n\|^2 + (1 - \beta_n)\|J_{r_n}x_n\|^2 \\ &= \beta_n\phi(u_n, x_n) + (1 - \beta_n)\phi(u_n, J_{r_n}x_n) \\ &\leq \phi(u_n, x_n) + \phi(u_n, J_{r_n}x_n),\end{aligned}\tag{3.39}$$

and hence

$$\begin{aligned}\phi(u_n, \tilde{x}_n) &= \phi\left(u_n, J^{-1}(\alpha_n Jx_0 + (1 - \alpha_n)Jz_n)\right) \\ &= \|u_n\|^2 - 2\langle u_n, \alpha_n Jx_0 + (1 - \alpha_n)Jz_n \rangle + \|\alpha_n Jx_0 + (1 - \alpha_n)Jz_n\|^2 \\ &\leq \|u_n\|^2 - 2\alpha_n\langle u_n, Jx_0 \rangle - 2(1 - \alpha_n)\langle u_n, Jz_n \rangle + \alpha_n\|x_0\|^2 + (1 - \alpha_n)\|z_n\|^2 \\ &= \alpha_n\phi(u_n, x_0) + (1 - \alpha_n)\phi(u_n, z_n) \\ &\leq \alpha_n\phi(u_n, x_0) + \phi(u_n, z_n) \\ &\leq \alpha_n\phi(u_n, x_0) + \phi(u_n, x_n) + \phi(u_n, J_{r_n}x_n).\end{aligned}\tag{3.40}$$

Since $\alpha_n \rightarrow 0$, $\phi(u_n, x_n) \rightarrow 0$ and $\phi(u_n, J_{r_n}x_n) \rightarrow 0$, it follows from the boundedness of $\{u_n\}$ that $\phi(u_n, \tilde{x}_n) \rightarrow 0$. Thus, in terms of Lemma 2.1, we have that $\|u_n - \tilde{x}_n\| \rightarrow 0$ and so $\|x_n - \tilde{x}_n\| \rightarrow 0$. Furthermore, since $\|\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{x}_n - J\tilde{x}_n\| = \tilde{\beta}_n\|Jx_0 - J\tilde{x}_n\| \rightarrow 0$, from the uniform norm-to-norm continuity of J^{-1} on bounded subsets of X^* , we obtain

$$\left\|J^{-1}\left(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{x}_n\right) - \tilde{x}_n\right\| \rightarrow 0.\tag{3.41}$$

Observe that

$$\begin{aligned}\|\tilde{x}_n - \tilde{J}_{r_n}\tilde{x}_n\| &\leq \|\tilde{x}_n - \tilde{z}_n\| + \|\tilde{z}_n - \tilde{J}_{r_n}\tilde{x}_n\| \\ &= \|\tilde{x}_n - \tilde{z}_n\| + \|\tilde{J}_{r_n}J^{-1}\left(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{x}_n\right) - \tilde{J}_{r_n}\tilde{x}_n\|.\end{aligned}\tag{3.42}$$

Thus, from (3.35) it follows that $\|\tilde{x}_n - \tilde{J}_{r_n}\tilde{x}_n\| \rightarrow 0$. Since J is uniformly norm-to-norm continuous on bounded subsets of X , it follows that $\|J\tilde{x}_n - J\tilde{J}_{r_n}\tilde{x}_n\| \rightarrow 0$.

Step 5. We claim that $\omega_w(\{x_n\}) \subset T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$, where

$$\omega_w(\{x_n\}) := \{\hat{x} \in C : x_{n_k} \rightarrow \hat{x} \text{ for some subsequence } \{n_k\} \subset \{n\} \text{ with } n_k \uparrow \infty\}.\tag{3.43}$$

Indeed, since $\{x_n\}$ is bounded and X is reflexive, we know that $\omega_w(\{x_n\}) \neq \emptyset$. Take $\hat{x} \in \omega_w(\{x_n\})$ arbitrarily. Then there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $x_{n_k} \rightharpoonup \hat{x}$. Hence it follows from $x_n - \tilde{x}_n \rightarrow 0$, $x_n - J_{r_n}x_n \rightarrow 0$, and $\tilde{x}_n - \tilde{J}_{r_n}\tilde{x}_n \rightarrow 0$ that $\{\tilde{x}_{n_k}\}$, $\{J_{r_{n_k}}x_{n_k}\}$, and $\{\tilde{J}_{r_{n_k}}\tilde{x}_{n_k}\}$ converge weakly to the same point \hat{x} . On the other hand, from (3.35), (3.36), and $\liminf_{n \rightarrow \infty} r_n > 0$ we obtain that

$$\lim_{n \rightarrow \infty} \|A_{r_n}x_n\| = \lim_{n \rightarrow \infty} \frac{1}{r_n} \|Jx_n - JJ_{r_n}x_n\| = 0, \tag{3.44}$$

$$\lim_{n \rightarrow \infty} \|\tilde{A}_{r_n}\tilde{x}_n\| = \lim_{n \rightarrow \infty} \frac{1}{r_n} \|J\tilde{x}_n - J\tilde{J}_{r_n}\tilde{x}_n\| = 0. \tag{3.45}$$

If $z^* \in Tz$ and $\tilde{z}^* \in \tilde{T}\tilde{z}$, then it follows from (2.17) and the monotonicity of the operators T, \tilde{T} that for all $k \geq 1$

$$\langle z - J_{r_{n_k}}x_{n_k}, z^* - A_{r_{n_k}}x_{n_k} \rangle \geq 0, \quad \langle \tilde{z} - \tilde{J}_{r_{n_k}}\tilde{x}_{n_k}, \tilde{z}^* - \tilde{A}_{r_{n_k}}\tilde{x}_{n_k} \rangle \geq 0. \tag{3.46}$$

Letting $k \rightarrow \infty$, we have that $\langle z - \hat{x}, z^* \rangle \geq 0$ and $\langle \tilde{z} - \hat{x}, \tilde{z}^* \rangle \geq 0$. Then the maximality of the operators T, \tilde{T} implies that $\hat{x} \in T^{-1}0$ and $\hat{x} \in \tilde{T}^{-1}0$. Next, let us show that $\hat{x} \in EP$. Since we have by (3.32)

$$\phi(u, y_n) \leq (\alpha_n + \tilde{\beta}_n)\phi(u, x_0) + \phi(u, x_n), \tag{3.47}$$

from $u_n = K_{r_n}y_n$ and Proposition 2.10 it follows that

$$\begin{aligned} \phi(u_n, y_n) &= \phi(K_{r_n}y_n, y_n) \\ &\leq \phi(u, y_n) - \phi(u, K_{r_n}y_n) \\ &\leq (\alpha_n + \tilde{\beta}_n)\phi(u, x_0) + \phi(u, x_n) - \phi(u, u_n). \end{aligned} \tag{3.48}$$

Also, since

$$\begin{aligned} |\phi(u, x_n) - \phi(u, u_n)| &= \left| \|x_n\|^2 - \|u_n\|^2 + 2\langle u, Ju_n - Jx_n \rangle \right| \\ &\leq \| \|x_n\| - \|u_n\| \| (\|x_n\| + \|u_n\|) + 2\|u\| \|Ju_n - Jx_n\| \\ &\leq \|x_n - u_n\| (\|x_n\| + \|u_n\|) + 2\|u\| \|Ju_n - Jx_n\|, \end{aligned} \tag{3.49}$$

so we get

$$\lim_{n \rightarrow \infty} (\phi(u, x_n) - \phi(u, u_n)) = 0. \tag{3.50}$$

Thus from (3.47), $\alpha_n \rightarrow 0$, $\tilde{\beta}_n \rightarrow 0$, and $\phi(u, x_n) - \phi(u, u_n) \rightarrow 0$, we have $\lim_{n \rightarrow \infty} \phi(u_n, y_n) = 0$. Since X is uniformly convex and smooth, we conclude from Lemma 2.1 that

$$\lim_{n \rightarrow \infty} \|u_n - y_n\| = 0. \quad (3.51)$$

From $x_{n_k} \rightharpoonup \hat{x}$, $x_n - u_n \rightarrow 0$, and (3.51), we have $y_{n_k} \rightharpoonup \hat{x}$ and $u_{n_k} \rightharpoonup \hat{x}$. Since J is uniformly norm-to-norm continuous on bounded subsets of X , from (3.51) we derive

$$\lim_{n \rightarrow \infty} \|Ju_n - Jy_n\| = 0. \quad (3.52)$$

From $\liminf_{n \rightarrow \infty} r_n > 0$, it follows that

$$\lim_{n \rightarrow \infty} \frac{\|Ju_n - Jy_n\|}{r_n} = 0. \quad (3.53)$$

By the definition of $u_n := K_{r_n}y_n$, we have

$$F(u_n, y) + \frac{1}{r_n} \langle y - u_n, Ju_n - Jy_n \rangle \geq 0, \quad \forall y \in C, \quad (3.54)$$

where

$$F(u_n, y) = f(u_n, y) + \langle Au_n, y - u_n \rangle. \quad (3.55)$$

Replacing n by n_k , we have from (A2) that

$$\frac{1}{r_{n_k}} \langle y - u_{n_k}, Ju_{n_k} - Jy_{n_k} \rangle \geq -F(u_{n_k}, y) \geq F(y, u_{n_k}), \quad \forall y \in C. \quad (3.56)$$

Since $y \mapsto f(x, y) + \langle Ax, y - x \rangle$ is convex and lower semicontinuous, it is also weakly lower semicontinuous. Letting $n_k \rightarrow \infty$ in the last inequality, from (3.53) and (A4) we have

$$F(y, \hat{x}) \leq 0, \quad \forall y \in C. \quad (3.57)$$

For t with $0 < t \leq 1$ and $y \in C$, let $y_t = ty + (1-t)\hat{x}$. Since $y \in C$ and $\hat{x} \in C$, $y_t \in C$ and hence $F(y_t, \hat{x}) \leq 0$. So, from (A1) we have

$$0 = F(y_t, y_t) \leq tF(y_t, y) + (1-t)F(y_t, \hat{x}) \leq tF(y_t, y). \quad (3.58)$$

Dividing by t , we have

$$F(y_t, y) \geq 0, \quad \forall y \in C. \quad (3.59)$$

Letting $t \downarrow 0$, from (A3) it follows that

$$F(\hat{x}, y) \geq 0, \quad \forall y \in C. \tag{3.60}$$

Thus $\hat{x} \in EP$. Therefore, we obtain that $\omega_w(\{x_n\}) \subset T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$ by the arbitrariness of \hat{x} .

Step 6. We claim that $\{x_n\}$ converges strongly to $w = \Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP}x_0$.

Indeed, from $x_{n+1} = \Pi_{H_n \cap W_n}x_0$ and $w \in T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP \subset H_n \cap W_n$, It follows that

$$\phi(x_{n+1}, x_0) \leq \phi(w, x_0). \tag{3.61}$$

Since the norm is weakly lower semicontinuous, we have

$$\begin{aligned} \phi(\hat{x}, x_0) &= \|\hat{x}\|^2 - 2\langle \hat{x}, Jx_0 \rangle + \|x_0\|^2 \leq \liminf_{k \rightarrow \infty} (\|x_{n_k}\|^2 - 2\langle x_{n_k}, Jx_0 \rangle + \|x_0\|^2) \\ &= \liminf_{k \rightarrow \infty} \phi(x_{n_k}, x_0) \leq \limsup_{k \rightarrow \infty} \phi(x_{n_k}, x_0) \leq \phi(w, x_0). \end{aligned} \tag{3.62}$$

From the definition of $\Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP}$, we have $\hat{x} = w$. Hence $\lim_{k \rightarrow \infty} \phi(x_{n_k}, x_0) = \phi(w, x_0)$ and

$$\begin{aligned} 0 &= \lim_{k \rightarrow \infty} (\phi(x_{n_k}, x_0) - \phi(w, x_0)) \\ &= \lim_{k \rightarrow \infty} (\|x_{n_k}\|^2 - \|w\|^2 - 2\langle x_{n_k} - w, Jx_0 \rangle) \\ &= \lim_{k \rightarrow \infty} (\|x_{n_k}\|^2 - \|w\|^2), \end{aligned} \tag{3.63}$$

which implies that $\lim_{k \rightarrow \infty} \|x_{n_k}\| = \|w\|$. Since X has the Kadec-Klee property, then $x_{n_k} \rightarrow w = \Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP}x_0$. Therefore, $\{x_n\}$ converges strongly to $\Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP}x_0$.

Remark 3.4. In Theorem 3.2, put $A \equiv 0$, $\tilde{T} \equiv 0$, and $\tilde{\beta}_n = 0$, $\forall n \geq 0$. Then, for all $\alpha, r \in (0, \infty)$ and $x, y \in C$, we have that

$$\begin{aligned} \langle Ax - Ay, x - y \rangle &\geq a\|Ax - Ay\|^2, \\ K_r(x) &= \left\{ u \in C : f(u, y) + \langle Au, y - u \rangle + \frac{1}{r}\langle y - u, Ju - Jx \rangle \geq 0, \forall y \in C \right\} \\ &= \left\{ u \in C : f(u, y) + \frac{1}{r}\langle y - u, Ju - Jx \rangle \geq 0, \forall y \in C \right\} = T_r(x). \end{aligned} \tag{3.64}$$

Moreover, the following hold:

$$\begin{aligned}
 H_n &= \left\{ z \in C : \phi(z, K_{r_n} y_n) \leq \left[\alpha_n + \tilde{\beta}_n - \alpha_n \tilde{\beta}_n - \tilde{\alpha}_n \tilde{\beta}_n + \alpha_n \tilde{\alpha}_n \tilde{\beta}_n \right] \phi(z, x_0) \right. \\
 &\quad \left. + \left[(1 - \alpha_n)(1 - \tilde{\alpha}_n)(1 - \tilde{\beta}_n) + \tilde{\alpha}_n(1 - \alpha_n) \right] \phi(z, x_n) \right\}, \\
 &= \{ z \in C : \phi(z, T_{r_n} y_n) \leq \alpha_n \phi(z, x_0) + (1 - \alpha_n) \phi(z, x_n) \}, \\
 y_n &= J^{-1} \left(\tilde{\alpha}_n J \tilde{x}_n + (1 - \tilde{\alpha}_n) J J_{r_n} J^{-1} \left(\tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J \tilde{x}_n \right) \right) \\
 &= J^{-1} \left(\tilde{\alpha}_n J \tilde{x}_n + (1 - \tilde{\alpha}_n) (0 J x_0 + (1 - 0) J \tilde{x}_n) \right) \\
 &= J^{-1} \left(\tilde{\alpha}_n J \tilde{x}_n + (1 - \tilde{\alpha}_n) J \tilde{x}_n \right) \\
 &= J^{-1} J \tilde{x}_n = \tilde{x}_n,
 \end{aligned} \tag{3.65}$$

and hence

$$y_n = \tilde{x}_n = J^{-1} (\alpha_n J x_0 + (1 - \alpha_n) (\beta_n J x_n + (1 - \beta_n) J J_{r_n} x_n)). \tag{3.66}$$

In this case, the previous Theorem 3.2 reduces to [20, Theorem 3.1]. \square

4. Weak Convergence Theorem

In this section, we present the following algorithm for finding a common element of the set of solutions for a generalized equilibrium problem and the set $T^{-1}0 \cap \tilde{T}^{-1}0$ for two maximal monotone operators T and \tilde{T} .

Let $x_0 \in X$ be chosen arbitrarily and consider the sequence $\{x_n\}$ generated by

$$\begin{aligned}
 \tilde{x}_n &= J^{-1} (\alpha_n J x_0 + (1 - \alpha_n) (\beta_n J K_{r_n} x_n + (1 - \beta_n) J J_{r_n} K_{r_n} x_n)), \\
 x_{n+1} &= J^{-1} \left(\tilde{\alpha}_n J K_{r_n} \tilde{x}_n + (1 - \tilde{\alpha}_n) J J_{r_n} J^{-1} \left(\tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J K_{r_n} \tilde{x}_n \right) \right), \quad n = 0, 1, 2, \dots,
 \end{aligned} \tag{4.1}$$

where $\{\alpha_n\}, \{\beta_n\}, \{\tilde{\alpha}_n\}, \{\tilde{\beta}_n\} \subset [0, 1]$, $\{r_n\} \subset (0, \infty)$, and $K_r, r > 0$, is defined by (2.15).

Before proving a weak convergence theorem, we need the following proposition.

Proposition 4.1. *Suppose that Assumption A is fulfilled and let $\{x_n\}$ be a sequence defined by (4.1), where $\{\alpha_n\}, \{\beta_n\}, \{\tilde{\alpha}_n\}, \{\tilde{\beta}_n\} \subset [0, 1]$ satisfy the following conditions:*

$$\sum_{n=0}^{\infty} \alpha_n < \infty, \quad \sum_{n=0}^{\infty} \tilde{\beta}_n < \infty, \quad \liminf_{n \rightarrow \infty} \beta_n (1 - \beta_n) > 0, \quad \liminf_{n \rightarrow \infty} \tilde{\alpha}_n (1 - \tilde{\alpha}_n) > 0. \tag{4.2}$$

Then, $\{\Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP} x_n\}$ converges strongly to $z \in T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$, where $\Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP}$ is the generalized projection of X onto $T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$.

Proof. We set $\Omega := T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$ and

$$\begin{aligned} u_n &:= K_{r_n} x_n, & y_n &:= J^{-1}(\beta_n J u_n + (1 - \beta_n) J J_{r_n} u_n), \\ \tilde{u}_n &:= K_{r_n} \tilde{x}_n, & \tilde{y}_n &:= \tilde{J}_{r_n} J^{-1}(\tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J \tilde{u}_n), \end{aligned} \quad (4.3)$$

so that

$$\begin{aligned} \tilde{x}_n &= J^{-1}(\alpha_n J x_0 + (1 - \alpha_n) J y_n), \\ x_{n+1} &= J^{-1}(\tilde{\alpha}_n J \tilde{u}_n + (1 - \tilde{\alpha}_n) J \tilde{y}_n), \quad n = 0, 1, 2, \dots \end{aligned} \quad (4.4)$$

Then, in terms of Lemma 2.4 and Proposition 2.10, Ω is a nonempty closed convex subset of X such that $\Omega \subset C$. We first prove that $\{x_n\}$ is bounded. Fix $u \in \Omega$. Note that by the first and third of (4.3), $u_n, \tilde{u}_n \in C$, and

$$\begin{aligned} F(u_n, y) + \frac{1}{r_n} \langle y - u_n, J u_n - J x_n \rangle &\geq 0, \quad \forall y \in C, \\ F(\tilde{u}_n, y) + \frac{1}{r_n} \langle y - \tilde{u}_n, J \tilde{u}_n - J \tilde{x}_n \rangle &\geq 0, \quad \forall y \in C. \end{aligned} \quad (4.5)$$

Here, each K_{r_n} is relatively nonexpansive. Then from Proposition 2.10 we obtain

$$\begin{aligned} \phi(u, y_n) &= \phi\left(u, J^{-1}(\beta_n J u_n + (1 - \beta_n) J J_{r_n} u_n)\right) \\ &= \|u\|^2 - 2\langle u, \beta_n J u_n + (1 - \beta_n) J J_{r_n} u_n \rangle + \|\beta_n J u_n + (1 - \beta_n) J J_{r_n} u_n\|^2 \\ &\leq \|u\|^2 - 2\beta_n \langle u, J u_n \rangle - 2(1 - \beta_n) \langle u, J J_{r_n} u_n \rangle + \beta_n \|u_n\|^2 + (1 - \beta_n) \|J_{r_n} u_n\|^2 \\ &= \beta_n \phi(u, u_n) + (1 - \beta_n) \phi(u, J_{r_n} u_n) \\ &\leq \beta_n \phi(u, u_n) + (1 - \beta_n) \phi(u, u_n) \\ &= \phi(u, u_n) = \phi(u, K_{r_n} x_n) \leq \phi(u, x_n), \end{aligned} \quad (4.6)$$

$$\begin{aligned}
\phi(u, \tilde{y}_n) &= \phi\left(u, \tilde{J}_n J^{-1}\left(\tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J \tilde{u}_n\right)\right) \\
&\leq \phi\left(u, J^{-1}\left(\tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J \tilde{u}_n\right)\right) \\
&= \|u\|^2 - 2\langle u, \tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J \tilde{u}_n \rangle + \left\| \tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J \tilde{u}_n \right\|^2 \\
&\leq \|u\|^2 - 2\tilde{\beta}_n \langle u, J x_0 \rangle - 2(1 - \tilde{\beta}_n) \langle u, J \tilde{u}_n \rangle + \tilde{\beta}_n \|x_0\|^2 + (1 - \tilde{\beta}_n) \|\tilde{u}_n\|^2 \\
&= \tilde{\beta}_n \phi(u, x_0) + (1 - \tilde{\beta}_n) \phi(u, \tilde{u}_n) \\
&\leq \tilde{\beta}_n \phi(u, x_0) + \phi(u, \tilde{u}_n) \\
&= \tilde{\beta}_n \phi(u, x_0) + \phi(u, K_{r_n} \tilde{x}_n) \\
&\leq \tilde{\beta}_n \phi(u, x_0) + \phi(u, \tilde{x}_n),
\end{aligned} \tag{4.7}$$

and hence by Proposition 2.10, we have

$$\begin{aligned}
\phi(u, \tilde{x}_n) &= \phi\left(u, J^{-1}\left(\alpha_n J x_0 + (1 - \alpha_n) J y_n\right)\right) \\
&= \|u\|^2 - 2\langle u, \alpha_n J x_0 + (1 - \alpha_n) J y_n \rangle + \left\| \alpha_n J x_0 + (1 - \alpha_n) J y_n \right\|^2 \\
&\leq \|u\|^2 - 2\alpha_n \langle u, J x_0 \rangle - 2(1 - \alpha_n) \langle u, J y_n \rangle + \alpha_n \|x_0\|^2 + (1 - \alpha_n) \|y_n\|^2 \\
&= \alpha_n \phi(u, x_0) + (1 - \alpha_n) \phi(u, y_n) \\
&\leq \alpha_n \phi(u, x_0) + \phi(u, y_n) \\
&\leq \phi(u, x_n) + \alpha_n \phi(u, x_0), \\
\phi(u, x_{n+1}) &= \phi\left(u, J^{-1}\left(\tilde{\alpha}_n J \tilde{u}_n + (1 - \tilde{\alpha}_n) J \tilde{y}_n\right)\right) \\
&= \|u\|^2 - 2\langle u, \tilde{\alpha}_n J \tilde{u}_n + (1 - \tilde{\alpha}_n) J \tilde{y}_n \rangle + \left\| \tilde{\alpha}_n J \tilde{u}_n + (1 - \tilde{\alpha}_n) J \tilde{y}_n \right\|^2 \\
&\leq \|u\|^2 - 2\tilde{\alpha}_n \langle u, J \tilde{u}_n \rangle - 2(1 - \tilde{\alpha}_n) \langle u, J \tilde{y}_n \rangle + \tilde{\alpha}_n \|\tilde{u}_n\|^2 + (1 - \tilde{\alpha}_n) \|\tilde{y}_n\|^2 \\
&= \tilde{\alpha}_n \phi(u, \tilde{u}_n) + (1 - \tilde{\alpha}_n) \phi(u, \tilde{y}_n) \\
&\leq \tilde{\alpha}_n \phi(u, \tilde{x}_n) + (1 - \tilde{\alpha}_n) \left[\tilde{\beta}_n \phi(u, x_0) + \phi(u, \tilde{x}_n) \right] \\
&\leq \phi(u, \tilde{x}_n) + \tilde{\beta}_n \phi(u, x_0).
\end{aligned} \tag{4.9}$$

Consequently, the last two inequalities yield that

$$\begin{aligned}\phi(u, x_{n+1}) &\leq \phi(u, \tilde{x}_n) + \tilde{\beta}_n \phi(u, x_0) \\ &\leq \phi(u, x_n) + \alpha_n \phi(u, x_0) + \tilde{\beta}_n \phi(u, x_0) \\ &= \phi(u, x_n) + (\alpha_n + \tilde{\beta}_n) \phi(u, x_0)\end{aligned}\tag{4.10}$$

for all $n \geq 0$. So, from $\sum_{n=0}^{\infty} \alpha_n < \infty$, $\sum_{n=0}^{\infty} \tilde{\beta}_n < \infty$, and Lemma 2.12, we deduce that $\lim_{n \rightarrow \infty} \phi(u, x_n)$ exists. This implies that $\{\phi(u, x_n)\}$ is bounded. Thus, $\{x_n\}$ is bounded and so are $\{u_n\}$, $\{\tilde{u}_n\}$, $\{J_{r_n} u_n\}$, and $\{\tilde{J}_{r_n} \tilde{u}_n\}$.

Define $z_n = \Pi_{\Omega} x_n$ for all $n \geq 0$. Let us show that $\{z_n\}$ is bounded. Indeed, observe that

$$\begin{aligned}(\|z_n\| - \|x_n\|)^2 &\leq \phi(z_n, x_n) = \phi(\Pi_{\Omega} x_n, x_n) \leq \phi(p, x_n) - \phi(p, \Pi_{\Omega} x_n) \\ &= \phi(p, x_n) - \phi(p, z_n) \leq \phi(p, x_n),\end{aligned}\tag{4.11}$$

for each $p \in \Omega$. This, together with the boundedness of $\{x_n\}$, implies that $\{z_n\}$ is bounded and so is $\phi(z_n, x_0)$. Furthermore, from $z_n \in \Omega$ and (4.10) we have

$$\phi(z_n, x_{n+1}) \leq \phi(z_n, x_n) + (\alpha_n + \tilde{\beta}_n) \phi(z_n, x_0).\tag{4.12}$$

Since Π_{Ω} is the generalized projection, then, from Lemma 2.3 we obtain

$$\begin{aligned}\phi(z_{n+1}, x_{n+1}) &= \phi(\Pi_{\Omega} x_{n+1}, x_{n+1}) \leq \phi(z_n, x_{n+1}) - \phi(z_n, \Pi_{\Omega} x_{n+1}) \\ &= \phi(z_n, x_{n+1}) - \phi(z_n, z_{n+1}) \leq \phi(z_n, x_{n+1}).\end{aligned}\tag{4.13}$$

Hence, from (4.12), it follows that $\phi(z_{n+1}, x_{n+1}) \leq \phi(z_n, x_n) + (\alpha_n + \tilde{\beta}_n) \phi(z_n, x_0)$.

Note that $\sum_{n=0}^{\infty} \alpha_n < \infty$, $\sum_{n=0}^{\infty} \tilde{\beta}_n < \infty$, and $\{\phi(z_n, x_0)\}$ is bounded, so that $\sum_{n=0}^{\infty} (\alpha_n + \tilde{\beta}_n) \phi(z_n, x_0) < \infty$. Therefore, $\{\phi(z_n, x_n)\}$ is a convergent sequence. On the other hand, from (4.10) we derive, for all $m \geq 0$,

$$\phi(u, x_{n+m}) \leq \phi(u, x_n) + \sum_{j=0}^{m-1} (\alpha_{n+j} + \tilde{\beta}_{n+j}) \phi(u, x_0).\tag{4.14}$$

In particular, we have

$$\phi(z_n, x_{n+m}) \leq \phi(z_n, x_n) + \sum_{j=0}^{m-1} (\alpha_{n+j} + \tilde{\beta}_{n+j}) \phi(z_n, x_0),\tag{4.15}$$

Consequently, from $z_{n+m} = \Pi_{\Omega}x_{n+m}$ and Lemma 2.3, we have

$$\phi(z_n, z_{n+m}) + \phi(z_{n+m}, x_{n+m}) \leq \phi(z_n, x_{n+m}) \leq \phi(z_n, x_n) + \sum_{j=0}^{m-1} (\alpha_{n+j} + \tilde{\beta}_{n+j}) \phi(z_n, x_0) \quad (4.16)$$

and hence

$$\phi(z_n, z_{n+m}) \leq \phi(z_n, x_n) - \phi(z_{n+m}, x_{n+m}) + \sum_{j=0}^{m-1} (\alpha_{n+j} + \tilde{\beta}_{n+j}) \phi(z_n, x_0). \quad (4.17)$$

Let $r = \sup\{\|z_n\| : n \geq 0\}$. From Lemma 2.6, there exists a continuous, strictly increasing, and convex function g with $g(0) = 0$ such that

$$g(\|x - y\|) \leq \phi(x, y), \quad \forall x, y \in B_r. \quad (4.18)$$

So, we have

$$\begin{aligned} g(\|z_n - z_{n+m}\|) &\leq \phi(z_n, z_{n+m}) \\ &\leq \phi(z_n, x_n) - \phi(z_{n+m}, x_{n+m}) + \sum_{j=0}^{m-1} (\alpha_{n+j} + \tilde{\beta}_{n+j}) \phi(z_n, x_0). \end{aligned} \quad (4.19)$$

Since $\{\phi(z_n, x_n)\}$ is a convergent sequence, $\{\phi(z_n, x_0)\}$ is bounded and $\sum_{n=0}^{\infty} (\alpha_n + \tilde{\beta}_n)$ is convergent, from the property of g we have that $\{z_n\}$ is a Cauchy sequence. Since Ω is closed, $\{z_n\}$ converges strongly to $z \in \Omega$. This completes the proof. \square

Now, we are in a position to prove the following theorem.

Theorem 4.2. *Suppose that Assumption A is fulfilled and let $\{x_n\}$ be a sequence defined by (4.1), where $\{\alpha_n\}, \{\beta_n\}, \{\tilde{\alpha}_n\}, \{\tilde{\beta}_n\} \subset [0, 1]$ satisfy the following conditions:*

$$\sum_{n=0}^{\infty} \alpha_n < \infty, \quad \sum_{n=0}^{\infty} \tilde{\beta}_n < \infty, \quad \liminf_{n \rightarrow \infty} \beta_n(1 - \beta_n) > 0, \quad \liminf_{n \rightarrow \infty} \tilde{\alpha}_n(1 - \tilde{\alpha}_n) > 0, \quad (4.20)$$

and $\{r_n\} \subset (0, \infty)$ satisfies $\liminf_{n \rightarrow \infty} r_n > 0$. If J is weakly sequentially continuous, then $\{x_n\}$ converges weakly to $z \in T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$, where $z = \lim_{n \rightarrow \infty} \Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP} x_n$.

Proof. We consider the notations (4.3). As in the proof of Proposition 4.1, we have that $\{x_n\}$, $\{u_n\}$, $\{J_{r_n}u_n\}$, $\{\tilde{x}_n\}$, $\{\tilde{u}_n\}$, and $\{\tilde{J}_{r_n}\tilde{u}_n\}$ are bounded sequences. Let

$$r = \sup\{\|u_n\|, \|J_{r_n}u_n\|, \|\tilde{u}_n\|, \|\tilde{y}_n\| : n \geq 0\}. \quad (4.21)$$

From Lemma 2.5 and as in the proof of Theorem 3.2, there exists a continuous, strictly increasing, and convex function g with $g(0) = 0$ such that

$$\|\alpha x^* + (1 - \alpha)y^*\|^2 \leq \alpha\|x^*\|^2 + (1 - \alpha)\|y^*\|^2 - \alpha(1 - \alpha)g(\|x^* - y^*\|) \quad (4.22)$$

for $x^*, y^* \in B_r^*$ and $\alpha \in [0, 1]$. Observe that for $u \in \Omega := T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP$,

$$\begin{aligned} \phi(u, y_n) &= \phi\left(u, J^{-1}(\beta_n J u_n + (1 - \beta_n) J J_{r_n} u_n)\right) \\ &= \|u\|^2 - 2\langle u, \beta_n J u_n + (1 - \beta_n) J J_{r_n} u_n \rangle + \|\beta_n J u_n + (1 - \beta_n) J J_{r_n} u_n\|^2 \\ &\leq \|u\|^2 - 2\beta_n \langle u, J u_n \rangle - 2(1 - \beta_n) \langle u, J J_{r_n} u_n \rangle \\ &\quad + \beta_n \|u_n\|^2 + (1 - \beta_n) \|J_{r_n} u_n\|^2 - \beta_n(1 - \beta_n) g(\|J u_n - J J_{r_n} u_n\|) \\ &\leq \beta_n \phi(u, u_n) + (1 - \beta_n) \phi(u, J_{r_n} u_n) - \beta_n(1 - \beta_n) g(\|J u_n - J J_{r_n} u_n\|) \\ &\leq \beta_n \phi(u, u_n) + (1 - \beta_n) \phi(u, u_n) - \beta_n(1 - \beta_n) g(\|J u_n - J J_{r_n} u_n\|) \\ &= \phi(u, u_n) - \beta_n(1 - \beta_n) g(\|J u_n - J J_{r_n} u_n\|), \\ \phi(u, \tilde{y}_n) &= \phi\left(u, \tilde{J}_{r_n} J^{-1}(\tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J \tilde{u}_n)\right) \quad (4.23) \\ &\leq \phi\left(u, J^{-1}(\tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J \tilde{u}_n)\right) \\ &= \|u\|^2 - 2\langle u, \tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J \tilde{u}_n \rangle + \|\tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J \tilde{u}_n\|^2 \\ &\leq \|u\|^2 - 2\tilde{\beta}_n \langle u, J x_0 \rangle - 2(1 - \tilde{\beta}_n) \langle u, J \tilde{u}_n \rangle + \tilde{\beta}_n \|x_0\|^2 + (1 - \tilde{\beta}_n) \|\tilde{u}_n\|^2 \\ &= \tilde{\beta}_n \phi(u, x_0) + (1 - \tilde{\beta}_n) \phi(u, \tilde{u}_n) \\ &= \tilde{\beta}_n \phi(u, x_0) + (1 - \tilde{\beta}_n) \phi(u, K_{r_n} \tilde{x}_n) \\ &\leq \tilde{\beta}_n \phi(u, x_0) + \phi(u, \tilde{x}_n). \end{aligned}$$

Hence,

$$\begin{aligned}
\phi(u, \tilde{x}_n) &= \phi\left(u, J^{-1}(\alpha_n Jx_0 + (1 - \alpha_n)Jy_n)\right) \\
&= \|u\|^2 - 2\langle u, \alpha_n Jx_0 + (1 - \alpha_n)Jy_n \rangle + \|\alpha_n Jx_0 + (1 - \alpha_n)Jy_n\|^2 \\
&\leq \|u\|^2 - 2\alpha_n \langle u, Jx_0 \rangle - 2(1 - \alpha_n) \langle u, Jy_n \rangle + \alpha_n \|x_0\|^2 + (1 - \alpha_n) \|y_n\|^2 \\
&= \alpha_n \phi(u, x_0) + (1 - \alpha_n) \phi(u, y_n) \\
&\leq \alpha_n \phi(u, x_0) + \phi(u, y_n) \\
&\leq \alpha_n \phi(u, x_0) + \phi(u, u_n) - \beta_n(1 - \beta_n)g(\|Ju_n - JJ_{r_n}u_n\|) \\
&= \alpha_n \phi(u, x_0) + \phi(u, K_{r_n}x_n) - \beta_n(1 - \beta_n)g(\|Ju_n - JJ_{r_n}u_n\|) \\
&\leq \alpha_n \phi(u, x_0) + \phi(u, x_n) - \beta_n(1 - \beta_n)g(\|Ju_n - JJ_{r_n}u_n\|), \\
\phi(u, x_{n+1}) &= \phi\left(u, J^{-1}(\tilde{\alpha}_n J\tilde{u}_n + (1 - \tilde{\alpha}_n)J\tilde{y}_n)\right) \\
&= \|u\|^2 - 2\langle u, \tilde{\alpha}_n J\tilde{u}_n + (1 - \tilde{\alpha}_n)J\tilde{y}_n \rangle + \|\tilde{\alpha}_n J\tilde{u}_n + (1 - \tilde{\alpha}_n)J\tilde{y}_n\|^2 \\
&\leq \|u\|^2 - 2\tilde{\alpha}_n \langle u, J\tilde{u}_n \rangle - 2(1 - \tilde{\alpha}_n) \langle u, J\tilde{y}_n \rangle + \tilde{\alpha}_n \|\tilde{u}_n\|^2 + (1 - \tilde{\alpha}_n) \|\tilde{y}_n\|^2 \\
&\quad - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{u}_n - J\tilde{y}_n\|) \\
&= \tilde{\alpha}_n \phi(u, \tilde{u}_n) + (1 - \tilde{\alpha}_n) \phi(u, \tilde{y}_n) - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{u}_n - J\tilde{y}_n\|) \\
&\leq \tilde{\alpha}_n \phi(u, K_{r_n}\tilde{x}_n) + (1 - \tilde{\alpha}_n) \left[\tilde{\beta}_n \phi(u, x_0) + \phi(u, \tilde{x}_n) \right] - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{u}_n - J\tilde{y}_n\|) \\
&\leq \phi(u, \tilde{x}_n) + \tilde{\beta}_n \phi(u, x_0) - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{u}_n - J\tilde{y}_n\|).
\end{aligned} \tag{4.24}$$

Consequently, the last two inequalities yield that

$$\begin{aligned}
\phi(u, x_{n+1}) &\leq \phi(u, \tilde{x}_n) + \tilde{\beta}_n \phi(u, x_0) - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{u}_n - J\tilde{y}_n\|) \\
&\leq \alpha_n \phi(u, x_0) + \phi(u, x_n) - \beta_n(1 - \beta_n)g(\|Ju_n - JJ_{r_n}u_n\|) \\
&\quad + \tilde{\beta}_n \phi(u, x_0) - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{u}_n - J\tilde{y}_n\|) \\
&= \phi(u, x_n) + (\alpha_n + \tilde{\beta}_n) \phi(u, x_0) - \beta_n(1 - \beta_n)g(\|Ju_n - JJ_{r_n}u_n\|) \\
&\quad - \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{u}_n - J\tilde{y}_n\|).
\end{aligned} \tag{4.26}$$

Thus, we have

$$\begin{aligned}
&\beta_n(1 - \beta_n)g(\|Ju_n - JJ_{r_n}u_n\|) + \tilde{\alpha}_n(1 - \tilde{\alpha}_n)g(\|J\tilde{u}_n - J\tilde{y}_n\|) \\
&\leq \phi(u, x_n) - \phi(u, x_{n+1}) + (\alpha_n + \tilde{\beta}_n) \phi(u, x_0).
\end{aligned} \tag{4.27}$$

By the proof of Proposition 4.1, it is known that $\{\phi(u, x_n)\}$ is convergent; since $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\lim_{n \rightarrow \infty} \tilde{\beta}_n = 0$, $\liminf_{n \rightarrow \infty} \beta_n(1 - \beta_n) > 0$, and $\liminf_{n \rightarrow \infty} \tilde{\alpha}_n(1 - \tilde{\alpha}_n) > 0$, then we have

$$\lim_{n \rightarrow \infty} g(\|Ju_n - JJ_{r_n}u_n\|) = \lim_{n \rightarrow \infty} g(\|J\tilde{u}_n - J\tilde{y}_n\|) = 0. \tag{4.28}$$

Taking into account the properties of g , as in the proof of Theorem 3.2, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|Ju_n - JJ_{r_n}u_n\| &= \lim_{n \rightarrow \infty} \|u_n - J_{r_n}u_n\| = 0, \\ \lim_{n \rightarrow \infty} \|J\tilde{u}_n - J\tilde{y}_n\| &= \lim_{n \rightarrow \infty} \|\tilde{u}_n - \tilde{y}_n\| = 0, \end{aligned} \tag{4.29}$$

since J^{-1} is uniformly norm-to-norm continuous on bounded subsets of X^* . Note that $\|\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{u}_n - J\tilde{u}_n\| = \tilde{\beta}_n\|Jx_0 - J\tilde{u}_n\| \rightarrow 0$. Hence, from the uniform norm-to-norm continuity of J^{-1} on bounded subsets of X^* we obtain $\|J^{-1}(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{u}_n) - \tilde{u}_n\| \rightarrow 0$. Also, observe that

$$\begin{aligned} \|\tilde{J}_{r_n}\tilde{u}_n - \tilde{u}_n\| &\leq \|\tilde{J}_{r_n}\tilde{u}_n - \tilde{J}_{r_n}J^{-1}(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{u}_n)\| \\ &\quad + \|\tilde{J}_{r_n}J^{-1}(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{u}_n) - \tilde{u}_n\| \\ &\leq \|\tilde{u}_n - J^{-1}(\tilde{\beta}_n Jx_0 + (1 - \tilde{\beta}_n)J\tilde{u}_n)\| + \|\tilde{y}_n - \tilde{u}_n\|. \end{aligned} \tag{4.30}$$

From $\|\tilde{u}_n - \tilde{y}_n\| \rightarrow 0$ it follows that $\|\tilde{J}_{r_n}\tilde{u}_n - \tilde{u}_n\| \rightarrow 0$. Since J is uniformly norm-to-norm continuous on bounded subsets of X , we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|Ju_n - JJ_{r_n}u_n\| &= \lim_{n \rightarrow \infty} \|u_n - J_{r_n}u_n\| = 0, \\ \lim_{n \rightarrow \infty} \|J\tilde{u}_n - J\tilde{J}_{r_n}\tilde{u}_n\| &= \lim_{n \rightarrow \infty} \|\tilde{u}_n - \tilde{J}_{r_n}\tilde{u}_n\| = 0. \end{aligned} \tag{4.31}$$

Now let us show that

$$\lim_{n \rightarrow \infty} \phi(u, x_n) = \lim_{n \rightarrow \infty} \phi(u, \tilde{x}_n) = \lim_{n \rightarrow \infty} \phi(u, u_n) = \lim_{n \rightarrow \infty} \phi(u, \tilde{u}_n). \tag{4.32}$$

Indeed, from (4.10) we get

$$\phi(u, x_{n+1}) - \tilde{\beta}_n\phi(u, x_0) \leq \phi(u, \tilde{x}_n) \leq \phi(u, x_n) + \alpha_n\phi(u, x_0), \tag{4.33}$$

which, together with $\lim_{n \rightarrow \infty} \alpha_n = \lim_{n \rightarrow \infty} \tilde{\beta}_n = 0$, yields that

$$\lim_{n \rightarrow \infty} \phi(u, \tilde{x}_n) = \lim_{n \rightarrow \infty} \phi(u, x_n). \tag{4.34}$$

From (4.9) it follows that

$$\begin{aligned}\phi(u, x_{n+1}) &\leq \tilde{\alpha}_n \phi(u, \tilde{u}_n) + (1 - \tilde{\alpha}_n) \phi(u, \tilde{y}_n) \\ &= \phi(u, \tilde{y}_n) + \tilde{\alpha}_n (\phi(u, \tilde{u}_n) - \phi(u, \tilde{y}_n)) \leq \phi(u, \tilde{x}_n) + \tilde{\beta}_n \phi(u, x_0).\end{aligned}\tag{4.35}$$

Note that

$$\begin{aligned}|\phi(u, \tilde{u}_n) - \phi(u, \tilde{y}_n)| &= \left| \|\tilde{u}_n\|^2 - \|\tilde{y}_n\|^2 + 2\langle u, J\tilde{y}_n - J\tilde{u}_n \rangle \right| \\ &\leq \left| \|\tilde{u}_n\| - \|\tilde{y}_n\| \right| (\|\tilde{u}_n\| + \|\tilde{y}_n\|) + 2\|u\| \|J\tilde{y}_n - J\tilde{u}_n\| \\ &\leq \|\tilde{u}_n - \tilde{y}_n\| (\|\tilde{u}_n\| + \|\tilde{y}_n\|) + 2\|u\| \|J\tilde{y}_n - J\tilde{u}_n\|.\end{aligned}\tag{4.36}$$

Since $\|\tilde{u}_n - \tilde{y}_n\| \rightarrow 0$ and $\|J\tilde{u}_n - J\tilde{y}_n\| \rightarrow 0$, we obtain $\lim_{n \rightarrow \infty} (\phi(u, \tilde{u}_n) - \phi(u, \tilde{y}_n)) = 0$, which, together with $\lim_{n \rightarrow \infty} \phi(u, \tilde{x}_n) = \lim_{n \rightarrow \infty} \phi(u, x_n)$, yields that

$$\lim_{n \rightarrow \infty} \phi(u, \tilde{y}_n) = \lim_{n \rightarrow \infty} \phi(u, x_n).\tag{4.37}$$

We have from (4.8) that

$$\phi(u, \tilde{x}_n) - \alpha_n \phi(u, x_0) \leq \phi(u, y_n) \leq \phi(u, x_n),\tag{4.38}$$

which, together with $\lim_{n \rightarrow \infty} \phi(u, \tilde{x}_n) = \lim_{n \rightarrow \infty} \phi(u, x_n)$, yields that

$$\lim_{n \rightarrow \infty} \phi(u, y_n) = \lim_{n \rightarrow \infty} \phi(u, x_n).\tag{4.39}$$

Also from (4.7) it follows that

$$\phi(u, \tilde{y}_n) - \tilde{\beta}_n \phi(u, x_0) \leq \phi(u, \tilde{u}_n) \leq \phi(u, \tilde{x}_n),\tag{4.40}$$

which, together with $\lim_{n \rightarrow \infty} \phi(u, \tilde{x}_n) = \lim_{n \rightarrow \infty} \phi(u, \tilde{y}_n) = \lim_{n \rightarrow \infty} \phi(u, x_n)$, yields that

$$\lim_{n \rightarrow \infty} \phi(u, \tilde{u}_n) = \lim_{n \rightarrow \infty} \phi(u, x_n).\tag{4.41}$$

Similarly from (4.6) it follows that

$$\phi(u, y_n) \leq \phi(u, u_n) \leq \phi(u, x_n) \quad (4.42)$$

which, together with $\lim_{n \rightarrow \infty} \phi(u, y_n) = \lim_{n \rightarrow \infty} \phi(u, x_n)$, yields that

$$\lim_{n \rightarrow \infty} \phi(u, u_n) = \lim_{n \rightarrow \infty} \phi(u, x_n). \quad (4.43)$$

On the other hand, let us show that

$$\lim_{n \rightarrow \infty} \|x_n - \tilde{x}_n\| = 0. \quad (4.44)$$

Indeed, let $s = \sup\{\|x_n\|, \|u_n\|, \|\tilde{x}_n\|, \|\tilde{u}_n\| : n \geq 0\}$. From Lemma 2.6, there exists a continuous, strictly increasing, and convex function g_1 with $g_1(0) = 0$ such that

$$g_1(\|x - y\|) \leq \phi(x, y), \quad \forall x, y \in B_s. \quad (4.45)$$

Since $u_n = K_{r_n} x_n$ and $\tilde{u}_n = K_{r_n} \tilde{x}_n$, we deduce from Proposition 2.10 that for $u \in \Omega$,

$$\begin{aligned} g_1(\|u_n - x_n\|) &\leq \phi(u_n, x_n) \leq \phi(u, x_n) - \phi(u, u_n), \\ g_1(\|\tilde{u}_n - \tilde{x}_n\|) &\leq \phi(\tilde{u}_n, \tilde{x}_n) \leq \phi(u, \tilde{x}_n) - \phi(u, \tilde{u}_n). \end{aligned} \quad (4.46)$$

This implies that

$$\lim_{n \rightarrow \infty} g_1(\|u_n - x_n\|) = \lim_{n \rightarrow \infty} g_1(\|\tilde{u}_n - \tilde{x}_n\|) = 0. \quad (4.47)$$

Since J is uniformly norm-to-norm continuous on bounded subsets of X , from the properties of g_1 we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \|u_n - x_n\| &= \lim_{n \rightarrow \infty} \|Ju_n - Jx_n\| = 0, \\ \lim_{n \rightarrow \infty} \|\tilde{u}_n - \tilde{x}_n\| &= \lim_{n \rightarrow \infty} \|J\tilde{u}_n - J\tilde{x}_n\| = 0. \end{aligned} \quad (4.48)$$

Note that

$$\begin{aligned}
 \phi(x_n, u_n) - \phi(u_n, x_n) &= \|x_n\|^2 - 2\langle x_n, Ju_n \rangle + \|u_n\|^2 - \left[\|x_n\|^2 - 2\langle u_n, Jx_n \rangle + \|u_n\|^2 \right] \\
 &= -2\langle x_n, Ju_n \rangle + 2\langle u_n, Jx_n \rangle \\
 &= 2\langle x_n, Jx_n - Ju_n \rangle + 2\langle u_n - x_n, Jx_n \rangle \\
 &\leq 2\|x_n\| \|Jx_n - Ju_n\| + 2\|u_n - x_n\| \|x_n\|,
 \end{aligned} \tag{4.49}$$

$$\begin{aligned}
 \phi(x_n, Jr_n u_n) &= \|x_n\|^2 - 2\langle x_n, JJ_r_n u_n \rangle + \|J_r_n u_n\|^2 \\
 &= \|x_n\|^2 - \|x_n\|^2 + \|J_r_n u_n\|^2 - \|x_n\|^2 + 2\langle x_n, Jx_n - JJ_r_n u_n \rangle \\
 &= (\|J_r_n u_n\| - \|x_n\|)(\|J_r_n u_n\| + \|x_n\|) + 2\langle x_n, Jx_n - JJ_r_n u_n \rangle \\
 &\leq \|J_r_n u_n - x_n\|(\|J_r_n u_n\| + \|x_n\|) + 2\|x_n\| \|Jx_n - JJ_r_n u_n\| \\
 &= \|J_r_n u_n - u_n + u_n - x_n\|(\|J_r_n u_n\| + \|x_n\|) \\
 &\quad + 2\|x_n\| \|Jx_n - Ju_n + Ju_n - JJ_r_n u_n\| \\
 &\leq (\|J_r_n u_n - u_n\| + \|u_n - x_n\|)(\|J_r_n u_n\| + \|x_n\|) \\
 &\quad + 2\|x_n\|(\|Jx_n - Ju_n\| + \|Ju_n - JJ_r_n u_n\|).
 \end{aligned} \tag{4.50}$$

Since $\phi(u_n, x_n) \rightarrow 0$, it follows from (4.31) and (4.35) that $\phi(x_n, u_n) \rightarrow 0$ and $\phi(x_n, Jr_n u_n) \rightarrow 0$. Also, observe that

$$\begin{aligned}
 \phi(x_n, y_n) &= \phi\left(x_n, J^{-1}(\beta_n Ju_n + (1 - \beta_n) JJ_r_n u_n)\right) \\
 &= \|x_n\|^2 - 2\langle x_n, \beta_n Ju_n + (1 - \beta_n) JJ_r_n u_n \rangle + \|\beta_n Ju_n + (1 - \beta_n) JJ_r_n u_n\|^2 \\
 &\leq \|x_n\|^2 - 2\beta_n \langle x_n, Ju_n \rangle - 2(1 - \beta_n) \langle x_n, JJ_r_n u_n \rangle + \beta_n \|u_n\|^2 + (1 - \beta_n) \|J_r_n u_n\|^2 \\
 &= \beta_n \phi(x_n, u_n) + (1 - \beta_n) \phi(x_n, Jr_n u_n) \\
 &\leq \phi(x_n, u_n) + \phi(x_n, Jr_n u_n),
 \end{aligned} \tag{4.51}$$

and hence

$$\begin{aligned}
 \phi(x_n, \tilde{x}_n) &= \phi\left(x_n, J^{-1}(\alpha_n Jx_0 + (1 - \alpha_n) Jy_n)\right) \\
 &= \|x_n\|^2 - 2\langle x_n, \alpha_n Jx_0 + (1 - \alpha_n) Jy_n \rangle + \|\alpha_n Jx_0 + (1 - \alpha_n) Jy_n\|^2 \\
 &\leq \|x_n\|^2 - 2\alpha_n \langle x_n, Jx_0 \rangle - 2(1 - \alpha_n) \langle x_n, Jy_n \rangle + \alpha_n \|x_0\|^2 + (1 - \alpha_n) \|y_n\|^2 \\
 &= \alpha_n \phi(x_n, x_0) + (1 - \alpha_n) \phi(x_n, y_n) \\
 &\leq \alpha_n \phi(x_n, x_0) + \phi(x_n, y_n) \\
 &\leq \alpha_n \phi(x_n, x_0) + \phi(x_n, u_n) + \phi(x_n, Jr_n u_n).
 \end{aligned} \tag{4.52}$$

Thus, from $\alpha_n \rightarrow 0$, $\phi(x_n, u_n) \rightarrow 0$, and $\phi(x_n, J_{r_n}u_n) \rightarrow 0$, it follows that $\phi(x_n, \tilde{x}_n) \rightarrow 0$. In terms of Lemma 2.1, we derive $\|x_n - \tilde{x}_n\| \rightarrow 0$.

Next, let us show that $x_n \rightharpoonup z$, where $z = \lim_{n \rightarrow \infty} \Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP} x_n$.

Indeed, since $\{x_n\}$ is bounded, there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $x_{n_k} \rightharpoonup \hat{x} \in C$. Hence it follows from (4.31), (4.35), and $\|x_n - \tilde{x}_n\| \rightarrow 0$ that both $\{u_{n_k}\}$, $\{\tilde{u}_{n_k}\}$, $\{J_{r_{n_k}}u_{n_k}\}$ and $\tilde{J}_{r_{n_k}}\tilde{u}_{n_k}$ converge weakly to the same point \hat{x} . Furthermore, from $\liminf_{n \rightarrow \infty} r_n > 0$ and (4.31) we have that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|A_{r_n}u_n\| &= \lim_{n \rightarrow \infty} \frac{1}{r_n} \|Ju_n - JJ_{r_n}u_n\| = 0, \\ \lim_{n \rightarrow \infty} \|\tilde{A}_{r_n}\tilde{u}_n\| &= \lim_{n \rightarrow \infty} \frac{1}{r_n} \|J\tilde{u}_n - J\tilde{J}_{r_n}\tilde{u}_n\| = 0. \end{aligned} \tag{4.53}$$

If $z^* \in Tz$ and $\tilde{z}^* \in \tilde{T}\tilde{z}$, then it follows from (2.17) and the monotonicity of the operators T, \tilde{T} that for all $k \geq 1$

$$\langle z - J_{r_{n_k}}u_{n_k}, z^* - A_{r_{n_k}}u_{n_k} \rangle \geq 0, \quad \langle \tilde{z} - \tilde{J}_{r_{n_k}}\tilde{u}_{n_k}, \tilde{z}^* - \tilde{A}_{r_{n_k}}\tilde{u}_{n_k} \rangle \geq 0. \tag{4.54}$$

Letting $k \rightarrow \infty$, we obtain that

$$\langle z - \hat{x}, z^* \rangle \geq 0, \quad \langle \tilde{z} - \hat{x}, \tilde{z}^* \rangle \geq 0. \tag{4.55}$$

Then the maximality of the operators T, \tilde{T} implies that $\hat{x} \in T^{-1}0 \cap \tilde{T}^{-1}0$.

Now, by the definition of $u_n := K_{r_n}x_n$, we have

$$F(u_n, y) + \frac{1}{r_n} \langle y - u_n, Ju_n - Jx_n \rangle \geq 0, \quad \forall y \in C, \tag{4.56}$$

where $F(x, y) = f(x, y) + \langle Ax, y - x \rangle$. Replacing n by n_k , we have from (A2) that

$$\frac{1}{r_{n_k}} \langle y - u_{n_k}, Ju_{n_k} - Jx_{n_k} \rangle \geq -F(u_{n_k}, y) \geq F(y, u_{n_k}), \quad \forall y \in C. \tag{4.57}$$

Since $y \mapsto F(x, y)$ is convex and lower semicontinuous, it is also weakly lower semicontinuous. Letting $n_k \rightarrow \infty$ in the last inequality, from (4.35) and (A4) we have

$$F(y, \hat{x}) \leq 0, \quad \forall y \in C. \tag{4.58}$$

For t , with $0 < t \leq 1$, and $y \in C$, let $y_t = ty + (1 - t)\hat{x}$. Since $y \in C$ and $\hat{x} \in C$, then $y_t \in C$ and hence $F(y_t, \hat{x}) \leq 0$. So, from (A1) we have

$$0 = F(y_t, y_t) \leq tF(y_t, y) + (1 - t)F(y_t, \hat{x}) \leq tF(y_t, y). \tag{4.59}$$

Dividing by t , we get $F(y_t, y) \geq 0, \forall y \in C$. Letting $t \downarrow 0$, from (A3) it follows that $F(\hat{x}, y) \geq 0, \forall y \in C$. So, $\hat{x} \in EP$. Therefore, $\hat{x} \in \Omega$. Let $z_n = \Pi_{\Omega} x_n$. From Lemma 2.2 and $\hat{x} \in \Omega$, we get

$$\langle z_{n_k} - \hat{x}, Jx_{n_k} - Jz_{n_k} \rangle \geq 0. \quad (4.60)$$

From Proposition 4.1, we also know that $z_n \rightarrow z \in \Omega$. Note that $x_{n_k} \rightharpoonup \hat{x}$. Since J is weakly sequentially continuous, then $\langle z - \hat{x}, J\hat{x} - Jz \rangle \geq 0$ as $k \rightarrow \infty$. In addition, taking into account the monotonicity of J , we conclude that $\langle z - \hat{x}, J\hat{x} - Jz \rangle \leq 0$. Hence

$$\langle z - \hat{x}, J\hat{x} - Jz \rangle = 0. \quad (4.61)$$

From the strict convexity of X , it follows that $z = \hat{x}$. Therefore, $x_n \rightharpoonup \hat{x}$, where $\hat{x} = \lim_{n \rightarrow \infty} \Pi_{T^{-1}0 \cap \tilde{T}^{-1}0 \cap EP} x_n$. This completes the proof. \square

Remark 4.3. In Theorem 4.2, put $A \equiv 0, \tilde{T} \equiv 0$, and $\tilde{\beta}_n = 0, \forall n \geq 0$. Then, for all $\alpha, r \in (0, \infty)$ and $x, y \in C$, we have that

$$\langle Ax - Ay, x - y \rangle \geq \alpha \|Ax - Ay\|^2, \quad (4.62)$$

$$\begin{aligned} K_r(x) &= \left\{ u \in C : f(u, y) + \langle Au, y - u \rangle + \frac{1}{r} \langle y - u, Ju - Jx \rangle \geq 0, \forall y \in C \right\} \\ &= \left\{ u \in C : f(u, y) + \frac{1}{r} \langle y - u, Ju - Jx \rangle \geq 0, \forall y \in C \right\} = T_r(x). \end{aligned} \quad (4.63)$$

Moreover, the following hold:

$$\begin{aligned} x_{n+1} &= J^{-1} \left(\tilde{\alpha}_n J K_{r_n} \tilde{x}_n + (1 - \tilde{\alpha}_n) J \tilde{J}_{r_n} J^{-1} \left(\tilde{\beta}_n J x_0 + (1 - \tilde{\beta}_n) J K_{r_n} \tilde{x}_n \right) \right) \\ &= J^{-1} \left(\tilde{\alpha}_n J T_{r_n} \tilde{x}_n + (1 - \tilde{\alpha}_n) (0 J x_0 + (1 - 0) J T_{r_n} \tilde{x}_n) \right) \\ &= J^{-1} \left(\tilde{\alpha}_n J T_{r_n} \tilde{x}_n + (1 - \tilde{\alpha}_n) J T_{r_n} \tilde{x}_n \right) \\ &= J^{-1} J T_{r_n} \tilde{x}_n = T_{r_n} \tilde{x}_n. \end{aligned} \quad (4.64)$$

In this case, Algorithm (4.1) reduces to the following one:

$$x_{n+1} = T_{r_n} J^{-1} \left(\alpha_n J x_0 + (1 - \alpha_n) (\beta_n J T_{r_n} x_n + (1 - \beta_n) J J_{r_n} T_{r_n} x_n) \right). \quad (4.65)$$

Corollary 4.4. Suppose that conditions (A1)–(A5) are fulfilled and let $\{x_n\}$ be a sequence defined by (4.65), where $T_r, r > 0$ is defined in Lemma 2.8, $\{\alpha_n\}, \{\beta_n\} \subset [0, 1]$ satisfy the conditions $\sum_{n=0}^{\infty} \alpha_n < \infty$ and $\liminf_{n \rightarrow \infty} \beta_n (1 - \beta_n) > 0$, and $\{r_n\} \subset (0, \infty)$ satisfies $\liminf_{n \rightarrow \infty} r_n > 0$. If J is weakly sequentially continuous, then $\{x_n\}$ converges weakly to $z \in T^{-1}0 \cap EP(f)$, where $z = \lim_{n \rightarrow \infty} \Pi_{T^{-1}0 \cap EP(f)} x_n$.

Acknowledgments

The first author (zenglc@hotmail.com) was partially supported by the National Science Foundation of China (10771141), Ph. D. Program Foundation of Ministry of Education of China (20070270004), and Science and Technology Commission of Shanghai Municipality grant (075105118), Leading Academic Discipline Project of Shanghai Normal University (DZL707), Shanghai Leading Academic Discipline Project (S30405) and Innovation Program of Shanghai Municipal Education Commission (09ZZ133). The third author (yaojc@math.nsysu.edu.tw) was partially supported by the Grant NSF 97-2115-M-110-001.

References

- [1] L.-C. Zeng and J.-C. Yao, "Strong convergence theorem by an extragradient method for fixed point problems and variational inequality problems," *Taiwanese Journal of Mathematics*, vol. 10, no. 5, pp. 1293–1303, 2006.
- [2] S. Schaible, J.-C. Yao, and L.-C. Zeng, "A proximal method for pseudomonotone type variational-like inequalities," *Taiwanese Journal of Mathematics*, vol. 10, no. 2, pp. 497–513, 2006.
- [3] L. C. Zeng, L. J. Lin, and J. C. Yao, "Auxiliary problem method for mixed variational-like inequalities," *Taiwanese Journal of Mathematics*, vol. 10, no. 2, pp. 515–529, 2006.
- [4] J.-W. Peng and J.-C. Yao, "Ishikawa iterative algorithms for a generalized equilibrium problem and fixed point problems of a pseudo-contraction mapping," to appear in *Journal of Global Optimization*.
- [5] L.-C. Zeng, S.-Y. Wu, and J.-C. Yao, "Generalized KKM theorem with applications to generalized minimax inequalities and generalized equilibrium problems," *Taiwanese Journal of Mathematics*, vol. 10, no. 6, pp. 1497–1514, 2006.
- [6] J.-W. Peng and J.-C. Yao, "Some new extragradient-like methods for generalized equilibrium problems, fixed points problems and variational inequality problems," to appear in *Optimization Methods & Software*.
- [7] L.-C. Ceng, C. Lee, and J.-C. Yao, "Strong weak convergence theorems of implicit hybrid steepest-descent methods for variational inequalities," *Taiwanese Journal of Mathematics*, vol. 12, no. 1, pp. 227–244, 2008.
- [8] J.-W. Peng and J.-C. Yao, "A new hybrid-extragradient method for generalized mixed equilibrium problems, fixed point problems and variational inequality problems," *Taiwanese Journal of Mathematics*, vol. 12, no. 6, pp. 1401–1432, 2008.
- [9] J.-W. Peng and J.-C. Yao, "Strong convergence theorems of iterative scheme based on the extragradient method for mixed equilibrium problems and fixed point problems," *Mathematical and Computer Modelling*, vol. 49, no. 9-10, pp. 1816–1828, 2009.
- [10] L.-C. Ceng, Q. H. Ansari, and J.-C. Yao, "Viscosity approximation methods for generalized equilibrium problems and fixed point problems," *Journal of Global Optimization*, vol. 43, no. 4, pp. 487–502, 2009.
- [11] J. W. Peng and J. C. Yao, "Some new iterative algorithms for generalized mixed equilibrium problems with strict pseudo-contractions and monotone mappings," to appear in *Taiwanese Journal of Mathematics*.
- [12] S. Kamimura and W. Takahashi, "Strong convergence of a proximal-type algorithm in a Banach space," *SIAM Journal on Optimization*, vol. 13, no. 3, pp. 938–945, 2003.
- [13] F. Kohsaka and W. Takahashi, "Strong convergence of an iterative sequence for maximal monotone operators in a Banach space," *Abstract and Applied Analysis*, vol. 2004, no. 3, pp. 239–249, 2004.
- [14] B. Martinet, "Régularisation d'inéquations variationnelles par approximations successives," *Revue Française d'Informatique et de Recherche Opérationnelle*, vol. 4, pp. 154–158, 1970.
- [15] W. Takahashi and K. Zembayashi, "Strong and weak convergence theorems for equilibrium problems and relatively nonexpansive mappings in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 1, pp. 45–57, 2009.
- [16] L. C. Ceng, G. Mastroeni, and J. C. Yao, "Hybrid proximal-point methods for common solutions of equilibrium problems and zeros of maximal monotone operators," *Journal of Optimization Theory and Applications*, vol. 142, no. 3, pp. 431–449, 2009.
- [17] S. S. Zhang, "Shrinking projection method for solving generalized equilibrium problem, variational inequality and common fixed point in Banach spaces with applications," to appear in *Science in China Series A*.

- [18] R. T. Rockafellar, "Monotone operators and the proximal point algorithm," *SIAM Journal on Control and Optimization*, vol. 14, no. 5, pp. 877–898, 1976.
- [19] L.-C. Ceng and J.-C. Yao, "A hybrid iterative scheme for mixed equilibrium problems and fixed point problems," *Journal of Computational and Applied Mathematics*, vol. 214, no. 1, pp. 186–201, 2008.
- [20] L. C. Zeng and J. C. Yao, "An inexact proximal-type algorithm in Banach spaces," *Journal of Optimization Theory and Applications*, vol. 135, no. 1, pp. 145–161, 2007.
- [21] S. Kamimura, F. Kohsaka, and W. Takahashi, "Weak and strong convergence theorems for maximal monotone operators in a Banach space," *Set-Valued Analysis*, vol. 12, no. 4, pp. 417–429, 2004.
- [22] L. Li and W. Song, "Modified proximal-point algorithm for maximal monotone operators in Banach spaces," *Journal of Optimization Theory and Applications*, vol. 138, no. 1, pp. 45–64, 2008.
- [23] O. Güler, "On the convergence of the proximal point algorithm for convex minimization," *SIAM Journal on Control and Optimization*, vol. 29, no. 2, pp. 403–419, 1991.
- [24] E. Blum and W. Oettli, "From optimization and variational inequalities to equilibrium problems," *The Mathematics Student*, vol. 63, no. 1–4, pp. 123–145, 1994.
- [25] L.-C. Ceng and J.-C. Yao, "Hybrid viscosity approximation schemes for equilibrium problems and fixed point problems of infinitely many nonexpansive mappings," *Applied Mathematics and Computation*, vol. 198, no. 2, pp. 729–741, 2008.
- [26] Y. I. Alber, "Metric and generalized projection operators in Banach spaces: properties and applications," in *Theory and Applications of Nonlinear Operators of Accretive and Monotone Type*, A. G. Kartsatos, Ed., vol. 178 of *Lecture Notes in Pure and Applied Mathematics*, pp. 15–50, Dekker, New York, NY, USA, 1996.
- [27] Y. I. Alber and S. Guerre-Delabriere, "On the projection methods for fixed point problems," *Analysis*, vol. 21, no. 1, pp. 17–39, 2001.
- [28] S. Reich, "A weak convergence theorem for the alternating method with Bregman distances," in *Theory and Applications of Nonlinear Operators of Accretive and Monotone Type*, A. G. Kartsatos, Ed., vol. 178 of *Lecture Notes in Pure and Applied Mathematics*, pp. 313–318, Dekker, New York, NY, USA, 1996.
- [29] I. Cioranescu, *Geometry of Banach Spaces, Duality Mappings and Nonlinear Problems*, vol. 62 of *Mathematics and Its Applications*, Kluwer Academic, Dordrecht, The Netherlands, 1990.
- [30] H. K. Xu, "Inequalities in Banach spaces with applications," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 16, no. 12, pp. 1127–1138, 1991.
- [31] P. L. Combettes and S. A. Hirstoaga, "Equilibrium programming in Hilbert spaces," *Journal of Nonlinear and Convex Analysis*, vol. 6, no. 1, pp. 117–136, 2005.
- [32] K.-K. Tan and H. K. Xu, "Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process," *Journal of Mathematical Analysis and Applications*, vol. 178, no. 2, pp. 301–308, 1993.