

Research Article

New Results on the Nonoscillation of Solutions of Some Nonlinear Differential Equations of Third Order

Ercan Tunç

Department of Mathematics, Faculty of Arts and Sciences, Gaziosmanpaşa University, 60240 Tokat, Turkey

Correspondence should be addressed to Ercan Tunç, ercantunc72@yahoo.com

Received 27 July 2009; Accepted 6 November 2009

Recommended by Patricia J. Y. Wong

We give sufficient conditions so that all solutions of differential equations $(r(t)y''(t))' + q(t)k(y'(t)) + p(t)y^\alpha(g(t)) = f(t)$, $t \geq t_0$, and $(r(t)y''(t))' + q(t)k(y'(t)) + p(t)h(y(g(t))) = f(t)$, $t \geq t_0$, are nonoscillatory. Depending on these criteria, some results which exist in the relevant literature are generalized. Furthermore, the conditions given for the functions k and h lead to studying more general differential equations.

Copyright © 2009 Ercan Tunç. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

This paper is concerned with study of nonoscillation of solutions of third-order nonlinear differential equations of the form

$$(r(t)y''(t))' + q(t)k(y'(t)) + p(t)y^\alpha(g(t)) = f(t), \quad t \geq t_0, \quad (1.1)$$

$$(r(t)y''(t))' + q(t)k(y'(t)) + p(t)h(y(g(t))) = f(t), \quad t \geq t_0, \quad (1.2)$$

where $t_0 \geq 0$ is a fixed real number, f, p, q, r , and $g \in C([0, \infty), \mathfrak{R})$ such that $r(t) > 0$ and $f(t) \geq 0$ for all $t \in [0, \infty)$. $k, h \in C(\mathfrak{R}, \mathfrak{R})$ are nondecreasing such that $h(y)y > 0$, $k(y')y' > 0$ for all $y \neq 0$, $y' \neq 0$. Throughout the paper, it is assumed, for all $g(t)$ and α appeared in (1.1) and (1.2), that $g(t) \leq t$ for all $t \geq t_0$; $\lim_{t \rightarrow \infty} g(t) = \infty$; $\alpha > 0$ is a quotient of odd integers.

It is well known from relevant literature that there have been deep and thorough studies on the nonoscillatory behaviour of solutions of second- and third-order nonlinear differential equations in recent years. See, for instance, [1–37] as some related papers or

books on the subject. In the most of these studies the following differential equation and some special cases of

$$(r(t)y''(t))' + q(t)(y')^\beta + p(t)y^\alpha = f(t), \quad t \geq t_0, \quad (1.3)$$

have been investigated. However, much less work has been done for nonoscillation of all solutions of nonlinear functional differential equations. In this connection, Parhi [10] established some sufficient conditions for oscillation of all solutions of the second-order forced differential equation of the form

$$(r(t)y'(t))' + p(t)y^\alpha(g(t)) = f(t) \quad (1.4)$$

and nonoscillation of all bounded solutions of the equations

$$\begin{aligned} (r(t)y'(t))' + q(t)(y'(t))^\beta + p(t)y^\alpha(g(t)) &= f(t), \\ (r(t)y'(t))' + q(t)(y'(g_1(t)))^\beta + p(t)y^\alpha(g(t)) &= f(t), \end{aligned} \quad (1.5)$$

where the real-valued functions f , p , q , r , g , and g_1 are continuous on $[0, \infty)$ with $r(t) > 0$ and $f(t) \geq 0$; $g(t) \leq t$, $g_1(t) \leq t$ for $t \geq t_0$; $\lim_{t \rightarrow \infty} g(t) = \infty$, $\lim_{t \rightarrow \infty} g_1(t) = \infty$, and both $\alpha > 0$ and $\beta > 0$ are quotients of odd integers.

Later, Nayak and Choudhury [5] considered the differential equation

$$(r(t)y''(t))' - q(t)(y'(t))^\beta - p(t)y^\alpha(g(t)) = f(t), \quad (1.6)$$

and they gave certain sufficient conditions on the functions involved for all bounded solutions of the above equation to be nonoscillatory.

Recently, in 2007, Tunç [23] investigated nonoscillation of solutions of the third-order differential equations:

$$\begin{aligned} (r(t)y''(t))' + q(t)y'(t) + p(t)y^\alpha(g(t)) &= f(t), \quad t \geq t_0, \\ (r(t)y''(t))' + q(t)(y'(g_1(t)))^\beta + p(t)y^\alpha(g(t)) &= f(t), \quad t \geq t_0. \end{aligned} \quad (1.7)$$

The motivation for the present work has come from the paper of Parhi [10], Tunç [23] and the papers mentioned above. We restrict our considerations to the real solutions of (1.1) and (1.2) which exist on the half-line $[T, \infty)$, where $T (\geq 0)$ depends on the particular solution, and are nontrivial in any neighborhood of infinity. It is well known that a solution $y(t)$ of (1.1) or (1.2) is said to be nonoscillatory on $[T, \infty)$ if there exists a $t_1 \geq T$ such that $y(t) \neq 0$ for $t \geq t_1$; it is said to be oscillatory if for any $t_1 \geq T$ there exist t_2 and t_3 satisfying $t_1 < t_2 < t_3$ such that $y(t_2) > 0$ and $y(t_3) < 0$; $y(t)$ is said to be a Z-type solution if it has arbitrarily large zeros but is ultimately nonnegative or nonpositive.

2. Nonoscillation Behaviors of Solutions of (1.1)

In this section, we obtain sufficient conditions for the nonoscillation of solutions of (1.1).

Theorem 2.1. *Let $q(t) \leq 0$. If $\lim_{t \rightarrow \infty} (f(t)/|p(t)|) = \infty$, then all bounded solutions of (1.1) are nonoscillatory.*

Proof. Let $y(t)$ be a bounded solution of (1.1) on $[T_y, \infty)$, $T_y \geq 0$, such that $|y(t)| \leq M$ for $t \geq T_y$. Since $\lim_{t \rightarrow \infty} g(t) = \infty$, there exists a $t_1 > t_0$ such that $g(t) \geq T_y$ for $t \geq t_1$. In view of the assumption $\lim_{t \rightarrow \infty} (f(t)/|p(t)|) = \infty$, it follows that there exists a $t_2 \geq t_1$ such that $f(t) > M^\alpha |p(t)|$ for $t \geq t_2$. If possible, let $y(t)$ be of nonnegative Z-type solution with consecutive double zeros at a and b ($t_2 < a < b$) such that $y(t) > 0$ for $t \in (a, b)$. So, there exists $c \in (a, b)$ such that $y'(c) = 0$ and $y'(t) > 0$ for $t \in (a, c)$. Multiplying (1.1) through by $y'(t)$, we get

$$(r(t)y'(t)y''(t))' = r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)y^\alpha(g(t))y'(t) + f(t)y'(t). \quad (2.1)$$

Integrating (2.1) from a to c , we obtain

$$\begin{aligned} 0 &= \int_a^c [r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) + f(t)y'(t) - p(t)y^\alpha(g(t))y'(t)] dt \\ &\geq \int_a^c [f(t) - p(t)y^\alpha(g(t))] y'(t) dt \\ &\geq \int_a^c [f(t) - M^\alpha |p(t)|] y'(t) dt > 0, \end{aligned} \quad (2.2)$$

which is a contradiction.

Let $y(t)$ be of nonpositive Z-type solution with consecutive double zeros at a and b ($t_2 < a < b$). Then, there exists a $c \in (a, b)$ such that $y'(c) = 0$ and $y'(t) > 0$ for $t \in (c, b)$.

Integrating (2.1) from c to b yields

$$\begin{aligned} 0 &= \int_c^b [r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) + f(t)y'(t) - p(t)y^\alpha(g(t))y'(t)] dt \\ &\geq \int_c^b [f(t) - |p(t)||y^\alpha(g(t))|] y'(t) dt \\ &\geq \int_c^b [f(t) - M^\alpha |p(t)|] y'(t) dt > 0, \end{aligned} \quad (2.3)$$

which is a contradiction.

If possible, let $y(t)$ be oscillatory with consecutive zeros at a, b and a' ($t_2 < a < b < a'$) such that $y'(a) \leq 0$, $y'(b) \geq 0$, $y'(a') \leq 0$, $y(t) < 0$ for $t \in (a, b)$ and $y(t) > 0$ for $t \in (b, a')$. So

there exists points $c \in (a, b)$ and $c' \in (b, a')$ such that $y'(c) = 0$, $y'(c') = 0$, $y'(t) > 0$ for $t \in (c, b)$ and $y'(t) > 0$ for $t \in (b, c')$. Now integrating (2.1) from c to c' , we get

$$\begin{aligned}
 0 &= \int_c^{c'} \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) + f(t)y'(t) - p(t)y^\alpha(g(t))y'(t) \right] dt \\
 &\geq \int_c^b [f(t) - p(t)y^\alpha(g(t))]y'(t)dt + \int_b^{c'} [f(t) - p(t)y^\alpha(g(t))]y'(t)dt \\
 &\geq \int_c^b [f(t) - |p(t)||y^\alpha(g(t))|]y'(t)dt + \int_b^{c'} [f(t) - |p(t)||y^\alpha(g(t))|]y'(t)dt \\
 &\geq \int_c^b [f(t) - M^\alpha|p(t)|]y'(t)dt + \int_b^{c'} [f(t) - M^\alpha|p(t)|]y'(t)dt > 0,
 \end{aligned} \tag{2.4}$$

which is a contradiction. This completes the proof of Theorem 2.1. \square

Remark 2.2. For the special case $k(y'(t)) = (y'(g_1(t)))^\beta$, $h(y(g(t))) = y^\alpha(g(t))$, Theorem 2.1 has been proved by Tunç [23]. Our results include the results established in Tunç [23].

Theorem 2.3. *Let $0 \leq p(t) < f(t)$ and $q(t) \leq 0$, then all solutions $y(t)$ of (1.1) which satisfy the inequality*

$$1 - z^\alpha(g(t)) \geq 0 \tag{2.5}$$

on any interval where $y'(t) > 0$ are nonoscillatory.

Proof. Let $y(t)$ be a solution of (1.1) on $[T_y, \infty)$, $T_y > 0$. Due to $\lim_{t \rightarrow \infty} g(t) = \infty$, there exists a $t_1 > t_0$ such that $g(t) \geq T_y$ for $t \geq t_1$. If possible, let $y(t)$ be of nonnegative Z-type solution with consecutive double zeros at a and b ($T_y \leq a < b$) such that $y(t) > 0$ for $t \in (a, b)$. So, there exists a $c \in (a, b)$ such that $y'(c) = 0$ and $y'(t) > 0$ for $t \in (a, c)$. Integrating (2.1) from a to c , we get

$$\begin{aligned}
 0 &= \int_a^c \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) + f(t)y'(t) - p(t)y^\alpha(g(t))y'(t) \right] dt \\
 &\geq \int_a^c [f(t) - p(t)y^\alpha(g(t))]y'(t)dt \\
 &\geq \int_a^c p(t)[1 - y^\alpha(g(t))]y'(t)dt > 0,
 \end{aligned} \tag{2.6}$$

which is a contradiction.

Next, let $y(t)$ be of nonpositive Z-type solution with consecutive double zeros at a and b ($T_y \leq a < b$). Then, there exists $c \in (a, b)$ such that $y'(c) = 0$ and $y'(t) > 0$ for $t \in (c, b)$.

Integrating (2.1) from c to b , we have

$$0 = \int_c^b \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) + f(t)y'(t) - p(t)y^\alpha(g(t))y'(t) \right] dt > 0, \quad (2.7)$$

which is a contradiction.

Now, if possible let $y(t)$ be oscillatory with consecutive zeros at a, b and a' ($T_y < a < b < a'$) such that $y'(a) \leq 0$, $y'(b) \geq 0$, $y'(a') \leq 0$, $y(t) < 0$ for $t \in (a, b)$ and $y(t) > 0$ for $t \in (b, a')$. Hence, there exist $c \in (a, b)$ and $c' \in (b, a')$ such that $y'(c) = y'(c') = 0$ and $y'(t) > 0$ for $t \in (c, b)$ and $t \in (b, c')$. Integrating (2.1) from c to c' , we obtain

$$\begin{aligned} 0 &= \int_c^{c'} \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) + f(t)y'(t) - p(t)y^\alpha(g(t))y'(t) \right] dt \\ &\geq \int_c^b [f(t) - p(t)y^\alpha(g(t))]y'(t)dt + \int_b^{c'} [f(t) - p(t)y^\alpha(g(t))]y'(t)dt \\ &\geq \int_b^{c'} [f(t) - p(t)y^\alpha(g(t))]y'(t)dt \\ &\geq \int_c^b p(t)[1 - y^\alpha(g(t))]y'(t)dt > 0, \end{aligned} \quad (2.8)$$

which is a contradiction. This completes the proof of Theorem 2.3. \square

Remark 2.4. For the special case $k(y') = (y')^\beta$, $y^\alpha(g(t)) = y^\alpha$, Theorem 2.3 has been proved by Tunç [25]. Our results include the results established in Tunç [25].

3. Nonoscillation Behaviors of Solutions (1.2)

In this section, we give sufficient conditions so that all solutions of (1.2) are nonoscillatory.

Theorem 3.1. *Suppose that $q(t) \leq 0$ and $0 \leq p(t) < f(t)$. If $y(t)$ is a solution (1.2) such that it satisfies the inequality*

$$1 - h(z(t)) > 0 \quad (3.1)$$

on any interval where $y'(t) > 0$, then $y(t)$ is nonoscillatory.

Proof. Let $y(t)$ be a solution of (1.2) on $[T_y, \infty)$, $T_y > 0$. Due to $\lim_{t \rightarrow \infty} g(t) = \infty$, there exists a $t_1 > t_0$ such that $g(t) \geq T_y$ for $t \geq t_1$. If possible, let $y(t)$ be of nonnegative Z-type solution with consecutive double zeros at a and b ($T_y \leq a < b$) such that $y(t) > 0$ for $t \in (a, b)$. So, there exists a $c \in (a, b)$ such that $y'(c) = 0$ and $y'(t) > 0$ for $t \in (a, c)$. Multiplying (1.2) through by $y'(t)$, we get

$$(r(t)y'(t)y''(t))' = r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t). \quad (3.2)$$

Integrating (3.2) from a to c , we get

$$\begin{aligned} 0 &= \int_a^c \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt \\ &\geq \int_a^c [f(t) - p(t)h(y(g(t)))]y'(t)dt \\ &\geq \int_a^c f(t)[1 - h(y(t))]y'(t)dt > 0, \end{aligned} \quad (3.3)$$

which is a contradiction.

Next, let $y(t)$ be of nonpositive Z -type solution with consecutive double zeros at a and b ($T_y \leq a < b$). Then, there exists $c \in (a, b)$ such that $y'(c) = 0$ and $y'(t) > 0$ for $t \in (c, b)$.

Integrating (3.2) from c to b , we have

$$0 = \int_c^b \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt > 0, \quad (3.4)$$

which is a contradiction.

Now, if possible let $y(t)$ be oscillatory with consecutive zeros at a, b and a' ($T_y < a < b < a'$) such that $y'(a) \leq 0, y'(b) \geq 0, y'(a') \leq 0, y(t) < 0$ for $t \in (a, b)$ and $y(t) > 0$ for $t \in (b, a')$. Hence, there exist $c \in (a, b)$ and $c' \in (b, a')$ such that $y'(c) = y'(c') = 0$ and $y'(t) > 0$ for $t \in (c, b)$ and $t \in (b, c')$. Integrating (3.2) from c to c' , we obtain

$$\begin{aligned} 0 &= \int_c^{c'} \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt \\ &\geq \int_c^b [f(t) - p(t)h(y(g(t)))]y'(t)dt + \int_b^{c'} [f(t) - p(t)h(y(g(t)))]y'(t)dt \\ &\geq \int_c^b [f(t) - p(t)h(y(t))]y'(t)dt + \int_b^{c'} [f(t) - p(t)h(y(t))]y'(t)dt \\ &\geq \int_b^{c'} [f(t) - p(t)h(y(t))]y'(t)dt \\ &\geq \int_b^{c'} f(t)[1 - h(y(t))]y'(t)dt > 0, \end{aligned} \quad (3.5)$$

which is a contradiction. This completes the proof of Theorem 3.1. \square

Theorem 3.2. Suppose that $0 \leq q \leq p \leq f$ and $q \neq 0$ on any subinterval of $[T_y, \infty)$, $T_y \geq 0$. If $y(t)$ is a solution of (1.2) such that it satisfies the inequality

$$1 - k(z') - h(z) > 0 \quad (3.6)$$

on any subinterval of $[T_y, \infty)$, $T_y \geq 0$, where $y'(t) > 0$, then $y(t)$ is nonoscillatory.

Proof. Let $y(t)$ be a solution of (1.2) on $[T_y, \infty)$, $T_y > 0$. Since $\lim_{t \rightarrow \infty} g(t) = \infty$, there exists a $t_1 > t_0$ such that $g(t) \geq T_y$ for $t \geq t_1$. If possible, let $y(t)$ be of nonnegative Z-type solution with consecutive double zeros at a and b ($T_y \leq a < b$) such that $y(t) > 0$ for $t \in (a, b)$. So, there exists a $c \in (a, b)$ such that $y'(c) = 0$ and $y'(t) > 0$ for $t \in (a, c)$. Integrating (3.2) from a to c , we get

$$\begin{aligned} 0 &= \int_a^c \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt \\ &\geq \int_a^c \left[-q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt \\ &\geq \int_a^c \left[-q(t)k(y'(t))y'(t) - p(t)h(y(t))y'(t) + f(t)y'(t) \right] dt \\ &\geq \int_a^c f(t) [1 - k(y'(t)) - p(t)h(y(t))] y'(t) dt > 0, \end{aligned} \tag{3.7}$$

which is a contradiction.

Next, let $y(t)$ be of nonpositive Z-type solution with consecutive double zeros at a and b ($T_y \leq a < b$). Then, there exists $c \in (a, b)$ such that $y'(c) = 0$ and $y'(t) > 0$ for $t \in (c, b)$.

Integrating (3.2) from c to b , we have

$$\begin{aligned} 0 &= \int_c^b \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt \\ &\geq \int_c^b \left[-q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt \\ &\geq \int_c^b q(t) [1 - k(y'(t)) - p(t)h(y(t))] y'(t) dt > 0, \end{aligned} \tag{3.8}$$

which is a contradiction.

Now, if possible let $y(t)$ be oscillatory with consecutive zeros at a, b and a' ($T_y < a < b < a'$) such that $y'(a) \leq 0$, $y'(b) \geq 0$, $y'(a') \leq 0$, $y(t) < 0$ for $t \in (a, b)$ and $y(t) > 0$ for $t \in (b, a')$. Hence, there exist $c \in (a, b)$ and $c' \in (b, a')$ such that $y'(c) = y'(c') = 0$ and $y'(t) > 0$ for $t \in (c, b)$ and $t \in (b, c')$. Integrating (3.2) from c to c' , we obtain

$$\begin{aligned} 0 &= \int_c^{c'} \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt \\ &\geq \int_c^b \left[-q(t)k(y'(t)) - p(t)h(y(g(t))) + f(t) \right] y'(t) dt \\ &\quad + \int_b^{c'} \left[-q(t)k(y'(t)) - p(t)h(y(g(t))) + f(t) \right] y'(t) dt \end{aligned}$$

$$\begin{aligned}
&\geq \int_c^b [-q(t)k(y'(t)) - p(t)h(y(t)) + f(t)]y'(t)dt \\
&\quad + \int_b^{c'} [-q(t)k(y'(t)) - p(t)h(y(t)) + f(t)]y'(t)dt \\
&\geq \int_c^b q(t)[1 - k(y'(t)) - h(y(t))]y'(t)dt + \int_b^{c'} f(t)[1 - k(y'(t)) - h(y(t))]y'(t)dt > 0,
\end{aligned} \tag{3.9}$$

which is a contradiction. This completes the proof of Theorem 3.2. \square

Remark 3.3. It is clear that Theorem 3.2 is not applicable to homogeneous equations:

$$(r(t)y''(t))' + q(t)k(y'(t)) + p(t)h(y(g(t))) = 0, \tag{3.10}$$

where $p(t) \geq 0$ and $q(t) \geq 0$.

Remark 3.4. For the special case $k(y') = (y')^\gamma$, $h(y(g(t))) = y^\beta$, Theorem 3.2 has been proved by N. parhi and S. parhi [19, Theorem 2.7].

Theorem 3.5. Let $p(t) \geq 0$, $q(t) \leq 0$, and $h(y) \leq y$ for all $y > 0$. If $p(t)$ and $f(t)$ are once continuously differentiable functions such that $p'(t) \geq 0$, $f'(t) \leq 0$, and $2f(t) - p(t) \geq 0$, then all solutions $y(t)$ of (1.2) for which $|y(t)| \leq 1$ ultimately are nonoscillatory.

Proof. Let $y(t)$ be a solution of (1.2) on $[T_y, \infty)$, $T_y > 0$, such that $|y(t)| \leq 1$ for $t \geq T_1 > T_y$. Since $\lim_{t \rightarrow \infty} g(t) = \infty$, there exists a $t_1 > t_0$ such that $g(t) \geq T_y$ for $t \geq t_1$. If possible, let $y(t)$ be of nonnegative Z-type solution with consecutive double zeros at a and b ($T_1 \leq a < b$) such that $y(t) > 0$ for $t \in (a, b)$. So, there exists a $c \in (a, b)$ such that $y'(c) = 0$ and $y'(t) > 0$ for $t \in (a, c)$. Integrating (3.2) from a to c , we get

$$0 = \int_a^c [r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t)]dt. \tag{3.11}$$

But

$$\begin{aligned}
\int_a^c f(t)y'(t)dt &= f(t)y(t)|_a^c - \int_a^c f'(t)y(t)dt \geq f(c)y(c), \\
\int_a^c p(t)h(y(g(t)))y'(t)dt &\leq \frac{1}{2}p(c)y^2(c).
\end{aligned} \tag{3.12}$$

Therefore

$$\begin{aligned}
&\int_a^c [-p(t)h(y(g(t)))y'(t) + f(t)y'(t)]dt \\
&\geq f(c)y(c) - \frac{1}{2}p(c)y^2(c) \geq \frac{p(c)}{2}y(c) - \frac{1}{2}p(c)y^2(c) = \frac{1}{2}p(c)[y(c) - y^2(c)] > 0,
\end{aligned} \tag{3.13}$$

since $|y(t)| \leq 1$ for $t \geq T_1$. So (3.11) yields

$$0 = \int_a^c \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt > 0, \quad (3.14)$$

which is a contradiction.

Next, let $y(t)$ be of nonpositive Z-type solution with consecutive double zeros at a and b ($T_1 \leq a < b$). Then, there exists $c \in (a, b)$ such that $y'(c) = 0$ and $y'(t) > 0$ for $t \in (c, b)$.

Integrating (3.2) from c to b , we have

$$0 = \int_c^b \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt > 0, \quad (3.15)$$

which is a contradiction.

Now, if possible let $y(t)$ be oscillatory with consecutive zeros at a, b and a' ($T_y < a < b < a'$) such that $y'(a) \leq 0$, $y'(b) \geq 0$, $y'(a') \leq 0$, $y(t) < 0$ for $t \in (a, b)$ and $y(t) > 0$ for $t \in (b, a')$. So there exist $c \in (a, b)$ and $c' \in (b, a')$ such that $y'(c) = 0$, $y'(c') = 0$ and $y'(t) > 0$ for $t \in (c, c')$. We consider two cases, namely, $y''(b) \leq 0$ and $y''(b) > 0$. Suppose that $y''(b) \leq 0$. Integrating (3.2) from c to b , we get

$$\begin{aligned} 0 &\geq r(b)y'(b)y''(b) \\ &= \int_c^b \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt \\ &> 0, \end{aligned} \quad (3.16)$$

which is a contradiction. Let $y''(b) > 0$. Integrating (3.2) from b to c' , we get

$$-r(b)y'(b)y''(b) = \int_b^{c'} \left[r(t)(y''(t))^2 - q(t)k(y'(t))y'(t) - p(t)h(y(g(t)))y'(t) + f(t)y'(t) \right] dt. \quad (3.17)$$

We proceed as in nonnegative Z-type to conclude that $0 \geq -r(b)y'(b)y''(b) > 0$. This is a contradiction. So $y(t)$ is nonoscillatory. This completes the proof of Theorem 3.5. \square

Remark 3.6. If $f \equiv 0$ in Theorem 3.5, then $p \equiv 0$ and hence the theorem is not applicable to homogeneous equation:

$$(r(t)y''(t))' + q(t)k(y'(t)) + p(t)h(y(g(t))) = 0. \quad (3.18)$$

Acknowledgment

The author would like to express sincere thanks to the anonymous referees for their invaluable corrections, comments, and suggestions.

References

- [1] S. R. Grace and B. S. Lalli, "On oscillation and nonoscillation of general functional-differential equations," *Journal of Mathematical Analysis and Applications*, vol. 109, no. 2, pp. 522–533, 1985.
- [2] J. R. Graef and M. Greguš, "Oscillatory properties of solutions of certain nonlinear third order differential equations," *Nonlinear Studies*, vol. 7, no. 1, pp. 43–50, 2000.
- [3] P. Hartman, *Ordinary Differential Equations*, Classics in Applied Mathematics, SIAM, Philadelphia, Pa, USA, 2002.
- [4] A. G. Kartsatos and M. N. Manougian, "Perturbations causing oscillations of functional-differential equations," *Proceedings of the American Mathematical Society*, vol. 43, pp. 111–117, 1974.
- [5] P. C. Nayak and R. Choudhury, "Oscillation and nonoscillation theorems for third order functional-differential equation," *The Journal of the Indian Mathematical Society. (New Series)*, vol. 62, no. 1–4, pp. 89–96, 1996.
- [6] S. Padhi, "On oscillatory solutions of third order differential equations," *Memoirs on Differential Equations and Mathematical Physics*, vol. 31, pp. 109–111, 2004.
- [7] S. Padhi, "On oscillatory linear third order forced differential equations," *Differential Equations and Dynamical Systems*, vol. 13, no. 3-4, pp. 343–358, 2005.
- [8] N. Parhi, "Nonoscillatory behaviour of solutions of nonhomogeneous third order differential equations," *Applicable Analysis*, vol. 12, no. 4, pp. 273–285, 1981.
- [9] N. Parhi, "Nonoscillation of solutions of a class of third order differential equations," *Acta Mathematica Hungarica*, vol. 54, no. 1-2, pp. 79–88, 1989.
- [10] N. Parhi, "Sufficient conditions for oscillation and nonoscillation of solutions of a class of second order functional-differential equations," *Analysis*, vol. 13, no. 1-2, pp. 19–28, 1993.
- [11] N. Parhi, "On non-homogeneous canonical third-order linear differential equations," *Australian Mathematical Society Journal*, vol. 57, no. 2, pp. 138–148, 1994.
- [12] N. Parhi and P. Das, "Oscillation criteria for a class of nonlinear differential equations of third order," *Annales Polonici Mathematici*, vol. 57, no. 3, pp. 219–229, 1992.
- [13] N. Parhi and P. Das, "On asymptotic property of solutions of linear homogeneous third order differential equations," *Unione Matematica Italiana. Bollettino B. Series VII*, vol. 7, no. 4, pp. 775–786, 1993.
- [14] N. Parhi and P. Das, "Oscillatory and asymptotic behaviour of a class of nonlinear functional-differential equations of third order," *Bulletin of the Calcutta Mathematical Society*, vol. 86, no. 3, pp. 253–266, 1994.
- [15] N. Parhi and P. Das, "On nonoscillation of third order differential equations," *Bulletin of the Institute of Mathematics Academia Sinica*, vol. 22, no. 3, pp. 267–274, 1994.
- [16] N. Parhi and S. Padhi, "On oscillatory linear differential equations of third order," *Archivum Mathematicum, Universitatis Masarykianae Brunensis*, vol. 37, no. 1, pp. 33–38, 2001.
- [17] N. Parhi and S. Padhi, "On oscillatory linear third order differential equations," *The Journal of the Indian Mathematical Society. (New Series)*, vol. 69, no. 1–4, pp. 113–128, 2002.
- [18] N. Parhi and S. Parhi, "Oscillation and nonoscillation theorems for nonhomogeneous third order differential equations," *Bulletin of the Institute of Mathematics Academia Sinica*, vol. 11, no. 2, pp. 125–139, 1983.
- [19] N. Parhi and S. Parhi, "Nonoscillation and asymptotic behaviour for forced nonlinear third order differential equations," *Bulletin of the Institute of Mathematics. Academia Sinica*, vol. 13, no. 4, pp. 367–384, 1985.
- [20] N. Parhi and S. Parhi, "On the behaviour of solutions of the differential equations $(r(t)y')' + q(t)(y'(t))^\beta + p(t)y^\alpha = f(t)$," *Polska Akademia Nauk. Annales Polonici Mathematici*, vol. 47, no. 2, pp. 137–148, 1986.
- [21] N. Parhi and S. Parhi, "Qualitative behaviour of solutions of forced nonlinear third order differential equations," *Rivista di Matematica della Università di Parma. Serie IV*, vol. 13, pp. 201–210, 1987.
- [22] C. A. Swanson, *Comparison and Oscillation Theory of Linear Differential Equations*, vol. 48 of *Mathematics in Science and Engineering*, Academic Press, New York, NY, USA, 1968.
- [23] C. Tunç, "On the non-oscillation of solutions of some nonlinear differential equations of third order," *Nonlinear Dynamics and Systems Theory*, vol. 7, no. 4, pp. 419–430, 2007.
- [24] C. Tunç, "On the nonoscillation of solutions of nonhomogeneous third order differential equations," *Soochow Journal of Mathematics*, vol. 23, no. 1, pp. 1–7, 1997.

- [25] C. Tunç, "Non-oscillation criteria for a class of nonlinear differential equations of third order," *Bulletin of the Greek Mathematical Society*, vol. 39, pp. 131–137, 1997.
- [26] C. Tunç and E. Tunç, "On the asymptotic behavior of solutions of certain second-order differential equations," *Journal of the Franklin Institute, Engineering and Applied Mathematics*, vol. 344, no. 5, pp. 391–398, 2007.
- [27] C. Tunç, "Uniform ultimate boundedness of the solutions of third-order nonlinear differential equations," *Kuwait Journal of Science & Engineering*, vol. 32, no. 1, pp. 39–48, 2005.
- [28] E. Tunç, "On the convergence of solutions of certain third-order differential equations," *Discrete Dynamics in Nature and Society*, vol. 2009, Article ID 863178, 12 pages, 2009.
- [29] E. Tunç, "Periodic solutions of a certain vector differential equation of sixth order," *The Arabian Journal for Science and Engineering A*, vol. 33, no. 1, pp. 107–112, 2008.
- [30] C. Tunç, "A new boundedness theorem for a class of second order differential equations," *The Arabian Journal for Science and Engineering A*, vol. 33, no. 1, pp. 1–10, 2008.
- [31] X.-Z. Zhong, H.-L. Xing, Y. Shi, J.-C. Liang, and D.-H. Wang, "Existence of nonoscillatory solution of third order linear neutral delay difference equation with positive and negative coefficients," *Nonlinear Dynamics and Systems Theory*, vol. 5, no. 2, pp. 201–214, 2005.
- [32] X. Zhong, J. Liang, Y. Shi, D. Wang, and L. Ge, "Existence of nonoscillatory solution of high-order nonlinear difference equation," *Nonlinear Dynamics and Systems Theory*, vol. 6, no. 2, pp. 205–210, 2006.
- [33] E. M. E. Zayed and M. A. El-Moneam, "Some oscillation criteria for second order nonlinear functional ordinary differential equations," *Acta Mathematica Scientia B*, vol. 27, no. 3, pp. 602–610, 2007.
- [34] E. M. E. Zayed, S. R. Grace, H. El-Metwally, and M. A. El-Moneam, "The oscillatory behavior of second order nonlinear functional differential equations," *The Arabian Journal for Science and Engineering A*, vol. 31, no. 1, pp. 23–30, 2006.
- [35] S. R. Grace, B. S. Lalli, and C. C. Yeh, "Oscillation theorems for nonlinear second order differential equations with a nonlinear damping term," *SIAM Journal on Mathematical Analysis*, vol. 15, no. 6, pp. 1082–1093, 1984.
- [36] S. R. Grace, "Oscillation criteria for forced functional-differential equations with deviating arguments," *Journal of Mathematical Analysis and Applications*, vol. 145, no. 1, pp. 63–88, 1990.
- [37] S. R. Grace and G. G. Hamedani, "On the oscillation of functional-differential equations," *Mathematische Nachrichten*, vol. 203, pp. 111–123, 1999.