

## Research Article

# Stability of Approximate Quadratic Mappings

**Hark-Mahn Kim, Minyoung Kim, and Juri Lee**

*Department of Mathematics, Chungnam National University, 79 Daehangno, Yuseong-gu, Daejeon 305-764, South Korea*

Correspondence should be addressed to Juri Lee, [annans@nate.com](mailto:annans@nate.com)

Received 23 October 2009; Revised 21 January 2010; Accepted 1 February 2010

Academic Editor: Yeol Je Cho

Copyright © 2010 Hark-Mahn Kim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the general solution of the quadratic functional equation  $f(2x + y) + 3f(2x - y) = 4f(x - y) + 12f(x)$ , in the class of all functions between quasi- $\beta$ -normed spaces, and then we prove the generalized Hyers-Ulam stability of the equation by using direct method and fixed point method.

## 1. Introduction

In 1940, Ulam [1] gave a talk before the Mathematics Club of the University of Wisconsin in which he discussed a number of unsolved problems. Among these was the following question concerning the stability of homomorphisms.

*Let  $G_1$  be a group and let  $G_2$  be a metric group with metric  $\rho(\cdot, \cdot)$ . Given  $\epsilon > 0$ , does there exist a  $\delta > 0$  such that if  $f : G_1 \rightarrow G_2$  satisfies  $\rho(f(xy), f(x)f(y)) < \delta$  for all  $x, y \in G_1$ , then a homomorphism  $h : G_1 \rightarrow G_2$  exists with  $\rho(f(x), h(x)) < \epsilon$  for all  $x \in G_1$ ?*

In 1941, the first result concerning the stability of functional equations was presented by Hyers [2]. And then Aoki [3] and Bourgin [4] have investigated the stability theorems of functional equations with unbounded Cauchy differences. In 1978, Th. M. Rassias [5] provided a generalization of Hyers' Theorem which allows the Cauchy difference to be unbounded. It was shown by Gajda [6] as well as by Th. M. Rassias and Šemrl [7] that one cannot prove the Rassias' type theorem when  $p = 1$ . Găvruta [8] obtained generalized result of Th. M. Rassias' Theorem which allow the Cauchy difference to be controlled by a general unbounded function. J. M. Rassias [9, 10] established a similar stability theorem linear and nonlinear mappings with the unbounded Cauchy difference.

Let  $E_1$  and  $E_2$  be real vector spaces. A function  $f : E_1 \rightarrow E_2$  is called a quadratic function if and only if  $f$  is a solution function of the quadratic functional equation:

$$f(x + y) + f(x - y) = 2f(x) + 2f(y). \quad (1.1)$$

It is well known that a function  $f$  between real vector spaces is quadratic if and only if there exists a unique symmetric biadditive function  $B$  such that  $f(x) = B(x, x)$  for all  $x$ , where the mapping  $B$  is given by  $B(x, y) = (1/4)(f(x + y) - f(x - y))$ . See [11, 12] for the details. The Hyers-Ulam stability of the quadratic functional (1.1) was first proved by Skof [13] for functions  $f : E_1 \rightarrow E_2$ , where  $E_1$  is a normed space and  $E_2$  is a Banach space. Cholewa [14] demonstrated that Skof's theorem is also valid if  $E_1$  is replaced by an abelian group. Czerwik [15] proved the Hyers-Ulam stability of quadratic functional (1.1) by the similar way to Th. M. Rassias control function [5]. According to the theorem of Borelli and Forti [16], we obtain the following generalization of stability theorem for the quadratic functional (1.1): let  $G$  be an abelian group and  $E$  a Banach space; let  $f : G \rightarrow E$  be a mapping with  $f(0) = 0$  satisfying the inequality

$$\|f(x + y) + f(x - y) - 2f(x) - 2f(y)\| \leq \varphi(x, y) \quad (1.2)$$

for all  $x, y \in G$ . Assume that one of the following conditions

$$\Phi(x, y) := \begin{cases} \sum_{k=0}^{\infty} \frac{1}{2^{2(k+1)}} \varphi(2^k x, 2^k y) < \infty, \\ \sum_{k=0}^{\infty} 2^{2k} \varphi\left(\frac{x}{2^{(k+1)}}, \frac{y}{2^{(k+1)}}\right) < \infty \end{cases} \quad (1.3)$$

holds for all  $x, y \in G$ , then there exists a unique quadratic function  $Q : G \rightarrow E$  such that

$$\|f(x) - Q(x)\| \leq \Phi(x, x) \quad (1.4)$$

for all  $x \in G$ . The stability problems of several functional equations have been extensively investigated by a number of authors and there are many interesting results concerning this problem [17–23].

In this paper, we consider a new quadratic functional equation

$$f(2x + y) + 3f(2x - y) = 4f(x - y) + 12f(x), \quad (1.5)$$

for all vectors in quasi- $\beta$ -normed spaces. First, we note that a function  $f$  is a solution of the functional (1.5) in the class of all functions between vector spaces if and only if the function  $f$  is quadratic. Further, we investigate the generalized Hyers-Ulam stability of (1.5) by using direct method and fixed point method. As a result of the paper, we have a much better possible estimation of approximate quadratic mappings by quadratic mappings than that of Czerwik [15] and Skof [13].

## 2. Stability of (1.5)

Now, we consider some basic concepts concerning quasi- $\beta$ -normed spaces and some preliminary results. We fix a realnumber  $\beta$  with  $0 < \beta \leq 1$  and let  $\mathbb{K}$  denote either  $\mathbb{R}$  or  $\mathbb{C}$ .

Let  $X$  be a linear space over  $\mathbb{K}$ . A *quasi- $\beta$ -norm*  $\|\cdot\|$  is a real-valued function on  $X$  satisfying the following.

- (1)  $\|x\| \geq 0$  for all  $x \in X$  and  $\|x\| = 0$  if and only if  $x = 0$ .
- (2)  $\|\lambda x\| = |\lambda|^\beta \cdot \|x\|$  for all  $\lambda \in \mathbb{K}$  and all  $x \in X$ .
- (3) There is a constant  $K \geq 1$  such that  $\|x + y\| \leq K(\|x\| + \|y\|)$  for all  $x, y \in X$ .

The pair  $(X, \|\cdot\|)$  is called a *quasi- $\beta$ -normed space* if  $\|\cdot\|$  is a quasi- $\beta$ -norm on  $X$ . The smallest possible  $K$  is called the *modulus of concavity* of  $\|\cdot\|$ . A *quasi- $\beta$ -Banach space* is a complete quasi- $\beta$ -normed space. A quasi- $\beta$ -norm  $\|\cdot\|$  is called a *( $\beta, p$ )-norm* ( $0 < p \leq 1$ ) if

$$\|x + y\|^p \leq \|x\|^p + \|y\|^p \quad (2.1)$$

for all  $x, y \in X$ . In this case, a quasi- $\beta$ -Banach space is called a *( $\beta, p$ )-Banach space*. We can refer to [24, 25] for the concept of quasinormed spaces and  $p$ -Banach spaces. Given a  $p$ -norm, the formula  $d(x, y) := \|x - y\|^p$  gives us a translation invariant metric on  $X$ . By the Aoki-Rolewicz theorem [25] (see also [24]), each quasinorm is equivalent to some  $p$ -norm. In [26], Tabor has investigated a version of the D. H. Hyers, Th. M. Rassias, and Z. Gajda theorem (see [5, 6]) in quasibanach spaces. Recently, J. M. Rassias and Kim [27] have obtained stability results of general additive equations in quasi- $\beta$ -normed spaces.

From now on, let  $X$  be a quasi- $\alpha$ -normed space with norm  $\|\cdot\|_\alpha$  and let  $Y$  be a *( $\beta, p$ )-Banach space* with norm  $\|\cdot\|_\beta$  unless we give any specific reference. Now, we are ready to investigate the generalized Hyers-Ulam stability problem for the functional (1.5) using direct method.

**Theorem 2.1.** *Assume that a function  $f : X \rightarrow Y$  satisfies*

$$\|Df(x, y) := f(2x + y) + 3f(2x - y) - 4f(x - y) - 12f(x)\|_\beta \leq \varphi(x, y) \quad (2.2)$$

for all  $x, y \in X$  and that  $\varphi$  satisfies the following control conditions

$$\Phi_1(x) := \sum_{i=0}^{\infty} \frac{\varphi(3^i x, 3^i x)^p}{9^{ip\beta}} < \infty, \quad \lim_{n \rightarrow \infty} \frac{\varphi(3^n x, 3^n y)^p}{9^{np\beta}} = 0 \quad (2.3)$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $Q : X \rightarrow Y$  satisfying

$$\left\| f(x) + \frac{f(0)}{2} - Q(x) \right\|_\beta \leq \frac{1}{9^\beta} \sqrt[p]{\Phi_1(x)} \quad (2.4)$$

for all  $x \in X$ , where  $\|f(0)\|_\beta \leq \varphi(0, 0)/12^\beta$ . The function  $Q$  is defined as

$$Q(x) = \lim_{k \rightarrow \infty} \frac{f(3^k x)}{3^{2k}} \quad (2.5)$$

for all  $x \in X$ .

*Proof.* Putting  $x, y := 0$  in (2.2), we get  $\|f(0)\|_\beta \leq \varphi(0, 0)/12^\beta$ . Replacing  $y$  by  $x$  in (2.2), we obtain

$$\|f(3x) - 9f(x) - 4f(0)\|_\beta \leq \varphi(x, x) \quad (2.6)$$

for all  $x \in X$ . Dividing (2.6) by  $9^\beta$ , we get

$$\left\| \frac{1}{9} \bar{f}(3x) - \bar{f}(x) \right\|_\beta \leq \frac{1}{9^\beta} \varphi(x, x) \quad (2.7)$$

for all  $x \in X$  where  $\bar{f}(x) = f(x) + f(0)/2$ ,  $x \in X$ . Now letting  $x := 3^i x$  and dividing  $3^{2ip\beta}$  in (2.7), we have

$$\left\| \frac{1}{3^{2(i+1)}} \bar{f}(3^{i+1}x) - \frac{1}{3^{2i}} \bar{f}(3^i x) \right\|_\beta^p \leq \frac{1}{9^{(i+1)p\beta}} \varphi(3^i x, 3^i x)^p \quad (2.8)$$

for all  $x \in X$ . Therefore we prove from the inequality (2.8) that for any integers  $m, n$  with  $m > n \geq 0$

$$\begin{aligned} \left\| \frac{1}{3^{2m}} \bar{f}(3^m x) - \frac{1}{3^{2n}} \bar{f}(3^n x) \right\|_\beta^p &\leq \sum_{i=n}^{m-1} \left\| \frac{\bar{f}(3^{i+1}x)}{3^{2(i+1)}} - \frac{\bar{f}(3^i x)}{3^{2i}} \right\|_\beta^p \\ &\leq \sum_{i=n}^{m-1} \frac{1}{9^{(i+1)p\beta}} \varphi(3^i x, 3^i x)^p. \end{aligned} \quad (2.9)$$

Since the right-hand side of (2.9) tends to zero as  $n \rightarrow \infty$ , the sequence  $\{(1/3^{2n})\bar{f}(3^n x)\}$  is Cauchy for all  $x \in X$  and thus converges by the completeness of  $Y$ . Define  $Q : X \rightarrow Y$  by

$$Q(x) = \lim_{n \rightarrow \infty} \frac{1}{3^{2n}} \left( f(3^n x) + \frac{f(0)}{2} \right) = \lim_{n \rightarrow \infty} \frac{f(3^n x)}{3^{2n}}, \quad x \in X. \quad (2.10)$$

Letting  $x := 3^n x$ ,  $y := 3^n y$  in (2.2), respectively, and dividing both sides by  $3^{2np\beta}$  and after then taking the limit in the resulting inequality, we have

$$\begin{aligned} &\|Q(2x + y) + 3Q(2x - y) - 4Q(x - y) - 12Q(x)\|_\beta^p \\ &= \lim_{n \rightarrow \infty} \frac{\|f(3^n(2x + y)) + 3f(3^n(2x - y)) - 4f(3^n(x - y)) - 12f(3^n x)\|_\beta^p}{9^{np\beta}} \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{9^{np\beta}} \varphi y(3^n x, 3^n x)^p = 0, \end{aligned} \quad (2.11)$$

and so the function  $Q$  is quadratic.

Taking the limit in (2.9) with  $n = 0$  as  $m \rightarrow \infty$ , we obtain that

$$\left\| f(x) + \frac{f(0)}{2} - Q(x) \right\|_{\beta}^p \leq \frac{1}{9^{p\beta}} \sum_{i=0}^{\infty} \frac{\varphi(3^i x, 3^i x)^p}{9^{ip\beta}}, \tag{2.12}$$

which yields the estimation (2.4).

To prove the uniqueness of the quadratic function  $Q$  subject to (2.4), let us assume that there exists a quadratic function  $Q' : X \rightarrow Y$  which satisfies (1.5) and the inequality (2.4). Obviously, we obtain that

$$Q(x) = 3^{-2n} Q(3^n x), \quad Q'(x) = 3^{-2n} Q'(3^n x) \tag{2.13}$$

for all  $x \in X$ . Hence it follows from (2.4) that

$$\begin{aligned} \|Q(x) - Q'(x)\|_{\beta}^p &\leq \frac{1}{3^{2np\beta}} \left( \left\| Q(3^n x) - f(3^n x) - \frac{f(0)}{2} \right\|_{\beta}^p + \left\| f(3^n x) + \frac{f(0)}{2} - Q'(3^n x) \right\|_{\beta}^p \right) \\ &\leq \frac{2}{9^{p\beta}} \sum_{i=0}^{\infty} \frac{1}{3^{2(n+i)p\beta}} \varphi(3^{n+i} x, 3^{n+i} x)^p = \frac{2}{9^{p\beta}} \sum_{j=n}^{\infty} \frac{1}{3^{2jp\beta}} \varphi(3^j x, 3^j x)^p \end{aligned} \tag{2.14}$$

for all  $n \in \mathbb{N}$ . Therefore letting  $n \rightarrow \infty$ , one has  $Q(x) - Q'(x) = 0$  for all  $x \in X$ , completing the proof of uniqueness.  $\square$

**Theorem 2.2.** Assume that a function  $f : X \rightarrow Y$  satisfies

$$\|Df(x, y)\|_{\beta} \leq \varphi(x, y) \tag{2.15}$$

for all  $x, y \in X$  and that  $\varphi$  satisfies conditions

$$\Phi_2(x) := \sum_{i=1}^{\infty} 9^{ip\beta} \varphi\left(\frac{x}{3^i}, \frac{x}{3^i}\right)^p < \infty, \quad \lim_{n \rightarrow \infty} 9^{np\beta} \varphi\left(\frac{x}{3^n}, \frac{y}{3^n}\right)^p = 0 \tag{2.16}$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $Q : X \rightarrow Y$  satisfying

$$\|f(x) - Q(x)\|_{\beta}^p \leq \frac{1}{9^{\beta}} \sqrt[p]{\Phi_2(x)} \tag{2.17}$$

for all  $x \in X$ . The function  $Q$  is given by

$$Q(x) = \lim_{n \rightarrow \infty} 3^{2n} f\left(\frac{x}{3^n}\right) \tag{2.18}$$

for all  $x \in X$ .

*Proof.* In this case,  $f(0) = 0$  since  $\sum_{i=1}^{\infty} (1/9^i)\varphi(0,0) < \infty$  and so  $\varphi(0,0) = 0$  by assumption.

Replacing  $x$  by  $x/3$  in (2.6), we obtain

$$\left\| f(x) - 9f\left(\frac{x}{3}\right) \right\|_{\beta} \leq \varphi\left(\frac{x}{3}, \frac{x}{3}\right) \quad (2.19)$$

for  $x \in X$ . Therefore we prove from inequality (2.19) that for any integers  $m, n$  with  $m > n \geq 0$

$$\begin{aligned} \left\| 9^m f\left(\frac{x}{3^m}\right) - 9^n f\left(\frac{x}{3^n}\right) \right\|_{\beta}^p &\leq \sum_{i=n}^{m-1} \left\| 9^i f\left(\frac{x}{3^i}\right) - 9^{i+1} f\left(\frac{x}{3^{i+1}}\right) \right\|_{\beta}^p \\ &\leq \sum_{i=n}^{m-1} 9^{ip\beta} \varphi\left(\frac{x}{3^{i+1}}, \frac{x}{3^{i+1}}\right)^p \\ &= \frac{1}{9^{p\beta}} \sum_{i=n}^{m-1} 9^{(i+1)p\beta} \varphi\left(\frac{x}{3^{i+1}}, \frac{x}{3^{i+1}}\right)^p \end{aligned} \quad (2.20)$$

for all  $x \in X$ . Since the right-hand side of (2.20) tends to zero as  $n \rightarrow \infty$ , the sequence  $\{3^{2n}f(x/3^n)\}$  is Cauchy for all  $x \in X$  and thus converges by the completeness of  $Y$ . Define  $Q : X \rightarrow Y$  by

$$Q(x) = \lim_{n \rightarrow \infty} 3^{2n} f\left(\frac{x}{3^n}\right) \quad (2.21)$$

for all  $x \in X$ .

Thereafter, applying the same argument as in the proof of Theorem 2.1, we obtain the desired result.  $\square$

We now introduce a fundamental result of fixed point theory. We refer to [28] for the proof of it, and the reader is referred to papers [29–31].

**Theorem 2.3.** *Let  $(\Omega, d)$  be a generalized complete metric space (i.e.,  $d$  may assume infinite values). Assume that  $\Lambda : \Omega \rightarrow \Omega$  is a strictly contractive operator with the Lipschitz constant  $0 < L < 1$ . Then for a given element  $x \in \Omega$  one of the following assertions is true:*

(A<sub>1</sub>)  $d(\Lambda^{k+1}x, \Lambda^kx) = \infty$  for all  $k \geq 0$ ;

(A<sub>2</sub>) there exists a nonnegative integer  $n_0$  such that

(A<sub>2.1</sub>)  $d(\Lambda^{n+1}x, \Lambda^n x) < \infty$  for all  $n \geq n_0$ ;

(A<sub>2.2</sub>) the sequence  $\{\Lambda^n x\}$  converges to a fixed point  $x^*$  of  $\Lambda$ ;

(A<sub>2.3</sub>)  $x^*$  is the unique fixed point of  $\Lambda$  in the set  $\Delta = \{y \in \Omega : d(\Lambda^{n_0}x, y) < \infty\}$ ;

(A<sub>2.4</sub>)  $d(y, x^*) \leq (1/1 - L)d(y, \Lambda y)$  for all  $y \in \Delta$ .

For an extensive theory of fixed point theorems and other nonlinear methods, the reader is referred to the book of Hyers et al. [32]. In 1996, Isac and Th. M. Rassias [33] applied the stability theory of functional equations to prove fixed point theorems and study some new applications in nonlinear analysis. Cădariu and Radu [29, 31] and Radu [34] applied the fixed

point theorem of alternative to the investigation of Cauchy and Jensen functional equations. Recently, Jung et al. [35–40] and Jung and Rassias [41] have obtained the generalized Hyers-Ulam stability of functional equations via the fixed point method.

Now we are ready to investigate the generalized Hyers-Ulam stability problem for the functional (1.5) using the fixed point method.

**Theorem 2.4.** *Let  $f : X \rightarrow Y$  be a function with  $f(0) = 0$  for which there exists a function  $\varphi : X^2 \rightarrow [0, \infty)$  such that there exists a constant  $L$ ,  $0 < L < 1$ , satisfying the inequalities*

$$\|Df(x, y)\|_{\beta} \leq \varphi(x, y), \quad (2.22)$$

$$\varphi(3x, 3y) \leq 9^{\beta} L \varphi(x, y) \quad (2.23)$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $Q : X \rightarrow Y$  defined by  $\lim_{k \rightarrow \infty} (f(3^k x) / 3^{2k}) = Q(x)$  such that

$$\|f(x) - Q(x)\|_{\beta} \leq \frac{1}{9^{\beta}(1-L)} \varphi(x, x) \quad (2.24)$$

for all  $x \in X$ .

*Proof.* Let us define  $\Omega$  to be the set of all functions  $g : X \rightarrow Y$  and introduce a generalized metric  $d$  on  $\Omega$  as follows:

$$d(g, h) = \inf \left\{ C \in [0, \infty) : \|g(x) - h(x)\|_{\beta} \leq C \varphi(x, x), \forall x \in X \right\}. \quad (2.25)$$

Then it is easy to show that  $(\Omega, d)$  is complete (see [37, Proof of Theorem 3.1]). Now we define an operator  $\Lambda : \Omega \rightarrow \Omega$  by

$$\Lambda g(x) = \frac{g(3x)}{9}, \quad g \in \Omega \quad (2.26)$$

for all  $x \in X$ . First, we assert that  $\Lambda$  is strictly contractive with constant  $L$  on  $\Omega$ . Given  $g, h \in \Omega$ , let  $C \in [0, \infty)$  be an arbitrary constant with  $d(g, h) \leq C$ , that is,  $\|g(x) - h(x)\|_{\beta} \leq C \varphi(x, x)$ . Then it follows from (2.23) that

$$\begin{aligned} \|\Lambda g(x) - \Lambda h(x)\|_{\beta} &= \frac{1}{9^{\beta}} \|g(3x) - h(3x)\|_{\beta} \leq \frac{1}{9^{\beta}} C \varphi(3x, 3x) \\ &\leq LC \varphi(x, x) \end{aligned} \quad (2.27)$$

for all  $x \in X$ , that is,  $d(\Lambda g, \Lambda h) \leq LC$  for any  $C \in [0, \infty)$  with  $d(g, h) \leq C$ . Thus we see that  $d(\Lambda g, \Lambda h) \leq Ld(g, h)$  for any  $g, h \in \Omega$  and so  $\Lambda$  is strictly contractive with constant  $L$  on  $\Omega$ .

Next, if we put  $(x, y) := (x, x)$  in (2.22) and we divide both sides by 9, then we get

$$\begin{aligned} \left\| \frac{f(3x)}{9} - f(x) \right\|_{\beta} &= \frac{1}{9^{\beta}} \|f(3x) - 9f(x)\|_{\beta} \\ &\leq \frac{1}{9^{\beta}} \varphi(x, x) \end{aligned} \quad (2.28)$$

for all  $x \in X$ , which implies  $d(\Lambda f, f) \leq 1/9^{\beta} < \infty$ .

Thus applying Theorem 2.3 to the complete generalized metric space  $(\Omega, d)$  with contractive constant  $L$ , we see from  $(A_{2.2})$  of Theorem 2.3 that there exists a function  $Q : X \rightarrow Y$  which is a fixed point of  $\Lambda$ , that is,  $Q(x) = \Lambda Q(x) = Q(3x)/9$ , such that  $d(\Lambda^k f, Q) \rightarrow 0$  as  $k \rightarrow \infty$ . By mathematical induction we know that

$$\Lambda^k Q(x) = \frac{Q(3^k x)}{3^{2k}} = Q(x) \quad (2.29)$$

for all  $k \in \mathbb{N}$ .

Since  $d(\Lambda^k f, Q) \rightarrow 0$  as  $k \rightarrow \infty$  by  $(A_{2.3})$  of Theorem 2.3, there exists a sequence  $\{C_k\}$  such that  $C_k \rightarrow 0$  as  $k \rightarrow \infty$ , and  $d(\Lambda^k f, Q) \leq C_k$  for every  $k \in \mathbb{N}$ . Hence, it follows from the definition of  $d$  that

$$\left\| \Lambda^k f(x) - Q(x) \right\|_{\beta} \leq C_k \varphi(x, x) \quad (2.30)$$

for all  $x \in X$ . This implies

$$\lim_{k \rightarrow \infty} \left\| \Lambda^k f(x) - Q(x) \right\|_{\beta} = 0, \quad \text{i.e.,} \quad \lim_{k \rightarrow \infty} \frac{f(3^k x)}{3^{2k}} = Q(x) \quad (2.31)$$

for all  $x \in X$ .

In turn, it follows from (2.22) and (2.23) that

$$\begin{aligned} \|DQ(x, y)\|_{\beta} &= \lim_{k \rightarrow \infty} \frac{1}{3^{2k\beta}} \left\| Df(3^k x, 3^k y) \right\|_{\beta} \\ &\leq \lim_{k \rightarrow \infty} \frac{1}{3^{2k\beta}} \varphi(3^k x, 3^k y) \leq \lim_{k \rightarrow \infty} L^k \varphi(x, y) \\ &= 0 \end{aligned} \quad (2.32)$$

for all  $x, y \in X$ , which implies that  $Q$  is a solution of (1.5) and so the mapping  $Q$  is quadratic. By  $(A_{2.4})$  of Theorem 2.3, we obtain

$$d(f, Q) \leq \frac{1}{1-L} d(\Lambda f, f) \leq \frac{1}{9^{\beta}(1-L)}, \quad (2.33)$$

which yields the inequality (2.24).

To prove the uniqueness of  $Q$ , assume now that  $Q_1 : X \rightarrow Y$  is another quadratic mapping satisfying the inequality (2.24). Then  $Q_1$  is a fixed point of  $\Lambda$  with  $d(f, Q_1) < \infty$  in view of the inequality (2.24). This implies that  $Q_1 \in \Delta = \{g \in \Omega : d(f, g) < \infty\}$  and so  $Q = Q_1$  by  $(A_{2.3})$  of Theorem 2.3. The proof is complete.  $\square$

By a similar way, one can prove the following theorem using the fixed point method.

**Theorem 2.5.** *Let  $f : X \rightarrow Y$  be a function with  $f(0) = 0$  for which there exists a function  $\varphi : X^2 \rightarrow [0, \infty)$  such that there exists a constant  $L$ ,  $0 < L < 1$ , satisfying the inequalities*

$$\|Df(x, y)\|_\beta \leq \varphi(x, y), \quad (2.34)$$

$$\varphi(x, y) \leq \frac{L}{9^\beta} \varphi(3x, 3y) \quad (2.35)$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $Q : X \rightarrow Y$  defined by  $\lim_{k \rightarrow \infty} 3^{2k} f(x/3^k) = Q(x)$  such that

$$\|f(x) - Q(x)\|_\beta \leq \frac{L}{9^\beta(1-L)} \varphi(x, x) \quad (2.36)$$

for all  $x \in X$ .

*Proof.* We use the same notations for  $\Omega$  and  $d$  as in the proof of Theorem 2.4. Thus  $(\Omega, d)$  is a complete generalized metric space. Let us define an operator  $\Lambda : \Omega \rightarrow \Omega$  by

$$\Lambda g(x) = 9g\left(\frac{x}{3}\right), \quad g \in \Omega \quad (2.37)$$

for all  $x \in X$ . Then it follows from (2.35) that

$$\|\Lambda g(x) - \Lambda h(x)\|_\beta = 9^\beta \left\| g\left(\frac{x}{3}\right) - h\left(\frac{x}{3}\right) \right\|_\beta \leq 9^\beta C \varphi\left(\frac{x}{3}, \frac{x}{3}\right) \leq LC \varphi(x, x) \quad (2.38)$$

for all  $x \in X$ , that is,  $d(\Lambda g, \Lambda h) \leq LC$ . Thus we see that  $d(\Lambda g, \Lambda h) \leq Ld(g, h)$  for any  $g, h \in \Omega$  and so  $\Lambda$  is strictly contractive with constant  $L$  on  $\Omega$ .

Next, if we put  $(x, y) := (x/3, x/3)$  in (2.34) and we divide both sides by  $1/9$ , then we get by virtue of (2.35)

$$\left\| f(x) - 9f\left(\frac{x}{3}\right) \right\|_\beta = \varphi\left(\frac{x}{3}, \frac{x}{3}\right) \leq \frac{L}{9^\beta} \varphi(x, x) \quad (2.39)$$

for all  $x \in X$ , which implies  $d(f, \Lambda f) \leq L/9^\beta < \infty$ . Thereafter, applying the same argument as in the proof of Theorem 2.4, we obtain the desired results.  $\square$

### 3. Applications of Main Results

In the following corollary, we have a stability result of (1.5) in the sense of Th. M. Rassias.

**Corollary 3.1.** *Let  $r_i$  and  $\varepsilon_i$  be real numbers such that  $\alpha(\max\{r_i : i = 1, 2\}) < 2\beta$  and  $\varepsilon_i \geq 0$  for  $i = 1, 2$ . Assume that a function  $f : X \rightarrow Y$  satisfies the inequality*

$$\|Df(x, y)\|_\beta \leq \varepsilon_1 \|x\|_\alpha^{r_1} + \varepsilon_2 \|y\|_\alpha^{r_2} \quad (3.1)$$

for all  $x, y \in X$ , and for all  $x, y \in X \setminus \{0\}$  if  $r_1, r_2 < 0$ . Then there exists a unique quadratic function  $Q : X \rightarrow Y$  which satisfies the inequality

$$\left\| f(x) + \frac{f(0)}{2} - Q(x) \right\|_\beta \leq \left[ \frac{\varepsilon_1^p \|x\|_\alpha^{pr_1}}{3^{p2\beta} - 3^{p\alpha r_1}} + \frac{\varepsilon_2^p \|x\|_\alpha^{pr_2}}{3^{p2\beta} - 3^{p\alpha r_2}} \right]^{1/p} \quad (3.2)$$

for all  $x \in X$ , and for all  $x \in X \setminus \{0\}$  if  $r_1, r_2 < 0$ . The function  $Q$  is given by

$$Q(x) = \lim_{n \rightarrow \infty} \frac{f(3^n x)}{3^{2n}}, \quad (3.3)$$

for all  $x \in X$ , where  $f(0) = 0$  if  $r_1, r_2 > 0$ .

*Proof.* If  $r_1, r_2 > 0$ , then we get  $f(0) = 0$  by putting  $x, y := 0$  in (3.1). Letting  $\varphi(x, y) := \varepsilon_1 \|x\|_\alpha^{r_1} + \varepsilon_2 \|y\|_\alpha^{r_2}$  for all  $x, y \in X$  and then applying Theorem 2.1 we obtain easily the desired results.  $\square$

**Corollary 3.2.** *Let  $r_i$  and  $\varepsilon_i$  be real numbers such that  $\alpha(\min\{r_i : i = 1, 2\}) > 2\beta$  and  $\varepsilon_i \geq 0$  for  $i = 1, 2$ . Assume that a function  $f : X \rightarrow Y$  satisfies the inequality*

$$\|Df(x, y)\|_\beta \leq \varepsilon_1 \|x\|_\alpha^{r_1} + \varepsilon_2 \|y\|_\alpha^{r_2} \quad (3.4)$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $Q : X \rightarrow Y$  which satisfies the inequality

$$\|f(x) - Q(x)\|_\beta \leq \left[ \frac{\varepsilon_1^p \|x\|_\alpha^{pr_1}}{3^{p\alpha r_1} - 3^{p2\beta}} + \frac{\varepsilon_2^p \|x\|_\alpha^{pr_2}}{3^{p\alpha r_2} - 3^{p2\beta}} \right]^{1/p} \quad (3.5)$$

for all  $x \in X$ . The function  $Q$  is given by

$$Q(x) = \lim_{n \rightarrow \infty} 3^{2n} f\left(\frac{x}{3^n}\right) \quad (3.6)$$

for all  $x \in X$ .

In the following corollary, we have a stability result of (1.5) in the sense of Hyers.

**Corollary 3.3.** *Let  $\delta$  be a nonnegative real number. Assume that a function  $f : X \rightarrow Y$  satisfies the inequality*

$$\|Df(x, y)\|_{\beta} \leq \delta \quad (3.7)$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $Q : X \rightarrow Y$ , defined by  $Q(x) = \lim_{n \rightarrow \infty} (f(3^n x)/3^{2n})$ , which satisfies the inequality

$$\left\| f(x) + \frac{f(0)}{2} - Q(x) \right\|_{\beta} \leq \frac{\delta}{9^{p\beta} - 1} \quad (3.8)$$

for all  $x \in X$ .

In the next corollary, we get a stability result of (1.5) in the sense of J. M. Rassias.

**Corollary 3.4.** *Let  $\varepsilon, r_1, r_2$  be real numbers such that  $\varepsilon \geq 0$  and  $\alpha r \neq 2\beta$ , where  $r := r_1 + r_2$ . Suppose that a function  $f : X \rightarrow Y$  satisfies*

$$\|Df(x, y)\|_{\beta} \leq \varepsilon \|x\|_{\alpha}^{r_1} \|y\|_{\alpha}^{r_2} \quad (3.9)$$

for all  $x, y \in X$ , and for all  $x, y \in X \setminus \{0\}$  if  $r_1, r_2 < 0$ . Then there exists a unique quadratic function  $Q : X \rightarrow Y$  which satisfies the inequality

$$\left\| f(x) + \frac{f(0)}{2} - Q(x) \right\|_{\beta} \leq \frac{\varepsilon \|x\|_{\alpha}^r}{\sqrt{|3^{p\alpha r} - 3^{p2\beta}|}} \quad (3.10)$$

for all  $x \in X$  and all  $x, y \in X \setminus \{0\}$  if  $r_1, r_2 < 0$ , where  $f(0) = 0$  if  $r_1, r_2 > 0$ .

*Proof.* Letting  $\varphi(x, y) := \varepsilon \|x\|_{\alpha}^{r_1} \|y\|_{\alpha}^{r_2}$  and applying Theorems 2.1 and 2.2, we get the results.  $\square$

## Acknowledgment

This work was supported by the National Research Foundation of Korea Grant funded by the Korean Government (no. 2009-0070940).

## References

- [1] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Tracts in Pure and Applied Mathematics, no. 8, Interscience, New York, NY, USA, 1960.
- [2] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [3] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.
- [4] D. G. Bourgin, "Classes of transformations and bordering transformations," *Bulletin of the American Mathematical Society*, vol. 57, pp. 223–237, 1951.
- [5] Th. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [6] Z. Gajda, "On stability of additive mappings," *International Journal of Mathematics and Mathematical Sciences*, vol. 14, no. 3, pp. 431–434, 1991.
- [7] Th. M. Rassias and P. Šemrl, "On the behavior of mappings which do not satisfy Hyers-Ulam stability," *Proceedings of the American Mathematical Society*, vol. 114, no. 4, pp. 989–993, 1992.
- [8] P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431–436, 1994.
- [9] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Bulletin des Sciences Mathématiques*, vol. 108, no. 4, pp. 445–446, 1984.
- [10] J. M. Rassias, "On the stability of the non-linear Euler-Lagrange functional equation in real normed linear spaces," *Journal of Mathematical and Physical Sciences*, vol. 28, no. 5, pp. 231–235, 1994.
- [11] J. Aczél and J. Dhombres, *Functional Equations in Several Variables*, vol. 31 of *Encyclopedia of Mathematics and Its Applications*, Cambridge University Press, Cambridge, UK, 1989.
- [12] D. H. Hyers and Th. M. Rassias, "Approximate homomorphisms," *Aequationes Mathematicae*, vol. 44, no. 2-3, pp. 125–153, 1992.
- [13] F. Skof, "Local properties and approximation of operators," *Rendiconti del Seminario Matematico e Fisico di Milano*, vol. 53, pp. 113–129, 1983.
- [14] P. W. Cholewa, "Remarks on the stability of functional equations," *Aequationes Mathematicae*, vol. 27, no. 1-2, pp. 76–86, 1984.
- [15] St. Czerwik, "On the stability of the quadratic mapping in normed spaces," *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, vol. 62, pp. 59–64, 1992.
- [16] C. Borelli and G. L. Forti, "On a general Hyers-Ulam stability result," *International Journal of Mathematics and Mathematical Sciences*, vol. 18, no. 2, pp. 229–236, 1995.
- [17] I.-S. Chang and H.-M. Kim, "On the Hyers-Ulam stability of quadratic functional equations," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 3, no. 3, article 33, 12 pages, 2002.
- [18] G.-L. Forti, "Comments on the core of the direct method for proving Hyers-Ulam stability of functional equations," *Journal of Mathematical Analysis and Applications*, vol. 295, no. 1, pp. 127–133, 2004.
- [19] M. Eshaghi Gordji, T. Karimi, and S. Kaboli Gharetapeh, "Approximately  $n$ -Jordan homomorphisms on Banach algebras," *Journal of Inequalities and Applications*, vol. 2009, Article ID 870843, 8 pages, 2009.
- [20] M. Eshaghi Gordji, S. Zolfaghari, J. M. Rassias, and M. B. Savadkouhi, "Solution and stability of a mixed type cubic and quartic functional equation in quasi-Banach spaces," *Abstract and Applied Analysis*, vol. 2009, Article ID 417473, 14 pages, 2009.
- [21] K.-W. Jun and H.-M. Kim, "Ulam stability problem for generalized  $A$ -quadratic mappings," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 2, pp. 466–476, 2005.
- [22] J. M. Rassias, "On the stability of the Euler-Lagrange functional equation," *Chinese Journal of Mathematics*, vol. 20, no. 2, pp. 185–190, 1992.
- [23] J. M. Rassias, "On the stability of the general Euler-Lagrange functional equation," *Demonstratio Mathematica*, vol. 29, no. 4, pp. 755–766, 1996.
- [24] Y. Benyamini and J. Lindenstrauss, *Geometric Nonlinear Functional Analysis. Vol. 1*, vol. 48 of *American Mathematical Society Colloquium Publications*, American Mathematical Society, Providence, RI, USA, 2000.
- [25] S. Rolewicz, *Metric Linear Spaces*, PWN-Polish Scientific, Warsaw, Poland, 2nd edition, 1984.
- [26] J. Tabor, "Stability of the Cauchy functional equation in quasi-Banach spaces," *Annales Polonici Mathematici*, vol. 83, no. 3, pp. 243–255, 2004.

- [27] J. M. Rassias and H.-M. Kim, "Generalized Hyers-Ulam stability for general additive functional equations in quasi- $\beta$ -normed spaces," *Journal of Mathematical Analysis and Applications*, vol. 356, no. 1, pp. 302–309, 2009.
- [28] J. B. Diaz and B. Margolis, "A fixed point theorem of the alternative, for contractions on a generalized complete metric space," *Bulletin of the American Mathematical Society*, vol. 74, no. 2, pp. 305–309, 1968.
- [29] L. Cădariu and V. Radu, "Fixed points and the stability of Jensen's functional equation," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 1, article 4, 7 pages, 2003.
- [30] L. Cădariu and V. Radu, "Fixed points and the stability of quadratic functional equations," *Analele Universității din Timișoara. Seria Matematică-Informatică*, vol. 41, no. 1, pp. 25–48, 2003.
- [31] L. Cădariu and V. Radu, "On the stability of the Cauchy functional equation: a fixed point approach," in *Iteration Theory (ECIT '02)*, vol. 346 of *Grazer Mathematische Berichte*, pp. 43–52, Karl-Franzens-Universität Graz, Graz, Austria, 2004.
- [32] D. H. Hyers, G. Isac, and Th. M. Rassias, *Topics in Nonlinear Analysis and Applications*, World Scientific, River Edge, NJ, USA, 1997.
- [33] G. Isac and Th. M. Rassias, "Stability of  $\psi$ -additive mappings: applications to nonlinear analysis," *International Journal of Mathematics and Mathematical Sciences*, vol. 19, no. 2, pp. 219–228, 1996.
- [34] V. Radu, "The fixed point alternative and the stability of functional equations," *Fixed Point Theory*, vol. 4, no. 1, pp. 91–96, 2003.
- [35] S.-M. Jung, "Stability of the quadratic equation of Pexider type," *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, vol. 70, pp. 175–190, 2000.
- [36] S.-M. Jung and T.-S. Kim, "A fixed point approach to the stability of the cubic functional equation," *Boletín de la Sociedad Matemática Mexicana*, vol. 12, no. 1, pp. 51–57, 2006.
- [37] S.-M. Jung and Z.-H. Lee, "A fixed point approach to the stability of quadratic functional equation with involution," *Fixed Point Theory and Applications*, vol. 2008, Article ID 732086, 11 pages, 2008.
- [38] S.-M. Jung, "A fixed point approach to the stability of isometries," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 2, pp. 879–890, 2007.
- [39] S.-M. Jung, "A fixed point approach to the stability of a Volterra integral equation," *Fixed Point Theory and Applications*, vol. 2007, Article ID 57064, 9 pages, 2007.
- [40] S.-M. Jung, T.-S. Kim, and K.-S. Lee, "A fixed point approach to the stability of quadratic functional equation," *Bulletin of the Korean Mathematical Society*, vol. 43, no. 3, pp. 531–541, 2006.
- [41] S.-M. Jung and J. M. Rassias, "A fixed point approach to the stability of a functional equation of the spiral of Theodorus," *Fixed Point Theory and Applications*, vol. 2008, Article ID 945010, 7 pages, 2008.