

Research Article

On Some Integral Inequalities on Time Scales and Their Applications

Run Xu,¹ Fanwei Meng,¹ and Cuihua Song²

¹ School of Mathematical Sciences, Qufu Normal University, Qufu, 273165 Shandong, China

² School of Chemistry and Chemical Engineering, Qufu Normal University, Qufu, 273165 Shandong, China

Correspondence should be addressed to Run Xu, xurun.2005@163.com

Received 15 January 2010; Revised 3 March 2010; Accepted 18 March 2010

Academic Editor: Martin Bohner

Copyright © 2010 Run Xu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The purpose of this paper is to investigate some new dynamic inequalities on time scales. We establish some new dynamic inequalities; the results unify and extend some continuous inequalities and their corresponding discrete analogues. The inequalities given here can be used as tools in the qualitative theory of certain dynamic equations. Some examples are given in the end of this paper.

1. Introduction

The theory of time scales was introduced by Hilger [1] in 1988 in order to contain both difference and differential calculus in a consistent way. Recently, many authors have extended some fundamental integral inequalities used in the theory of differential and integral equations on time scales. For example, we refer the reader to the papers [2–12] and the references cited there in.

In this paper, we investigate some nonlinear integral inequalities on time scales, which extend some inequalities established by Li and Sheng [8] and Li [9]. The obtained inequalities can be used as important tools in the study of dynamic equations on time scales.

Throughout this paper, let us assume that we have already acquired the knowledge of time scales and time scales notation; for an excellent introduction to the calculus on time scales, we refer the reader to Bohner and Peterson [4] for general overview.

2. Some Preliminaries on Time Scales

In what follows, \mathbb{R} denotes the set of real numbers, \mathbb{Z} denotes the set of integers, \mathbb{N}_0 denotes the set of nonnegative integers, \mathbb{C} denotes the set of complex numbers, and $C(M, S)$ denotes

the class of all continuous functions defined on set M with range in the set S . \mathbb{T} is an arbitrary time scale. If \mathbb{T} has a right-scattered maximum m , then the set $\mathbb{T}^k = \mathbb{T} - \{m\}$; otherwise, $\mathbb{T}^k = \mathbb{T}$. C_{rd} denotes the set of rd-continuous functions; \mathcal{R} denotes the set of all regressive and rd-continuous functions. We define the set of all positively regressive functions by $\mathcal{R}^+ = \{p \in \mathcal{R} : 1 + \mu(t)p(t) > 0, t \in \mathbb{T}\}$. Obviously, if $p \in C_{rd}$ and $p(t) \geq 0$ for $t \in \mathbb{T}$, then $p \in \mathcal{R}^+$.

For $f : \mathbb{T} \rightarrow \mathbb{R}$ and $t \in \mathbb{T}^k$, we define $f^\Delta(t)$ as follows (provided it exists):

$$f^\Delta(t) := \lim_{s \rightarrow t} \frac{f^\sigma(t) - f(s)}{\sigma(t) - s}; \quad (2.1)$$

we call $f^\Delta(t)$ the *delta derivative* of f at t .

The following lemmas are very useful in our main results.

Lemma 2.1 (see [4]). *If $p \in \mathcal{R}$ and fix $t_0 \in \mathbb{T}$, then the exponential function $e_p(\cdot, t_0)$ is for the unique solution of the initial value problem*

$$x^\Delta = p(t)x, \quad x(t_0) = 1 \quad \text{on } \mathbb{T}. \quad (2.2)$$

Lemma 2.2 (see [4]). *Let $t_0 \in \mathbb{T}^k$ and $w : \mathbb{T} \times \mathbb{T}^k \rightarrow \mathbb{R}$ be continuous at (t, t) , where $t \geq t_0$. Assume that $w^\Delta(t, \cdot)$ is rd-continuous on $[t_0, \sigma(t)]$. If for any $\varepsilon > 0$, there exists a neighborhood \mathcal{U} of t , independent of $\tau \in [t_0, \sigma(t)]$, such that*

$$\left| w(\sigma(t), \tau) - w(s, \tau) - w^\Delta(t, \tau)(\sigma(t) - s) \right| \leq \varepsilon |\sigma(t) - s|, \quad s \in \mathcal{U}, \quad (2.3)$$

where w^Δ denotes the derivative of w with respect to the first variable, then

$$g(t) := \int_{t_0}^t w(t, \tau) \Delta \tau \quad (2.4)$$

implies

$$g^\Delta(t) := \int_{t_0}^t w^\Delta(t, \tau) \Delta \tau + w(\sigma(t), t). \quad (2.5)$$

The following theorem is a foundational result in dynamic inequalities.

Lemma 2.3 (Comparison Theorem [4]). *Suppose $u, b \in C_{rd}$, $a \in \mathcal{R}^+$; then*

$$u^\Delta(t) \leq a(t)u(t) + b(t), \quad t \geq t_0, \quad t \in \mathbb{T}^k, \quad (2.6)$$

implies

$$u(t) \leq u(t_0)e_a(t, t_0) + \int_{t_0}^t b(\tau)e_a(t, \sigma(\tau)) \Delta \tau, \quad t \geq t_0, \quad t \in \mathbb{T}^k. \quad (2.7)$$

The following lemma is useful in our main results.

Lemma 2.4 (see [7]). *Let $a \geq 0$, $p \geq q > 0$, then*

$$a^{q/p} \leq \frac{q}{p} K^{(q-p)/p} a + \frac{p-q}{p} K^{q/p}, \quad K > 0. \quad (2.8)$$

3. Main Results

In this section, we study some integral inequalities on time scales. We always assume that p, q, r, m are constants, $p \geq q > 0$, $p \geq m > 0$, $p \geq r > 0$, and $t \geq t_0, t \in \mathbb{T}^k$.

Theorem 3.1. *Assume that $u, a, b, f, g, h \in C_{\text{rd}}$; $u(t), a(t), b(t), f(t), g(t)$, and $h(t)$ are nonnegative; then*

$$u^p(t) \leq a(t) + b(t) \int_{t_0}^t \left[f(s)u^q(s) + g(s)u^r(s) + \int_{t_0}^s h(\tau)u^m(\tau)\Delta\tau \right] \Delta s, \quad t \in \mathbb{T}^k, \quad (3.1)$$

implies

$$u(t) \leq \left\{ a(t) + b(t) \int_{t_0}^t B(\tau)e_{A(\tau)}(t, \sigma(\tau))\Delta\tau \right\}^{1/p}, \quad K > 0, \quad t \in \mathbb{T}^k, \quad (3.2)$$

where

$$\begin{aligned} A(t) &= \left[\frac{q}{p} K^{(q-p)/p} f(t) + \frac{r}{p} K^{(r-p)/p} g(t) \right] b(t) + \frac{m}{p} K^{(m-p)/p} \int_{t_0}^t b(\tau)h(\tau)\Delta\tau, \\ B(t) &= f(t) \left[\frac{q}{p} K^{(q-p)/p} a(t) + \frac{p-q}{p} K^{q/p} \right] + g(t) \left[\frac{r}{p} K^{(r-p)/p} a(t) + \frac{p-r}{p} K^{p/r} \right] \\ &\quad + \int_{t_0}^t \left[\frac{m}{p} K^{(m-p)/p} a(\tau) + \frac{p-m}{p} K^{m/p} \right] h(\tau)\Delta\tau, \quad t \in \mathbb{T}^k. \end{aligned} \quad (3.3)$$

Proof. Define $z(t)$ by

$$z(t) = \int_{t_0}^t \left[f(s)u^q(s) + g(s)u^r(s) + \int_{t_0}^s h(\tau)u^m(\tau)\Delta\tau \right] \Delta s, \quad (3.4)$$

then $z(t_0) = 0$, and (3.1) can be restated as

$$u^p(t) \leq a(t) + b(t)z(t). \quad (3.5)$$

Using Lemma 2.1, for any $k > 0$, we obtain

$$\begin{aligned} u^q(t) &\leq [a(t) + b(t)z(t)]^{q/p} \leq \frac{q}{p}K^{(q-p)/p}[a(t) + b(t)z(t)] + \frac{p-q}{p}K^{q/p}, \\ u^r(t) &\leq [a(t) + b(t)z(t)]^{r/p} \leq \frac{r}{p}K^{(r-p)/p}[a(t) + b(t)z(t)] + \frac{p-r}{p}K^{r/p}, \\ u^m(t) &\leq [a(t) + b(t)z(t)]^{m/p} \leq \frac{m}{p}K^{(m-p)/p}[a(t) + b(t)z(t)] + \frac{p-m}{p}K^{m/p}. \end{aligned} \quad (3.6)$$

It follows from (3.4) and (3.6) that

$$\begin{aligned} z^\Delta(t) &= f(t)u^q(t) + g(t)u^r(t) + \int_{t_0}^t h(\tau)u^m(\tau)\Delta\tau \\ &\leq f(t)\left[\frac{q}{p}K^{(q-p)/p}(a(t) + b(t)z(t)) + \frac{p-q}{p}K^{q/p}\right] \\ &\quad + g(t)\left[\frac{r}{p}K^{(r-p)/p}(a(t) + b(t)z(t)) + \frac{p-r}{p}K^{r/p}\right] \\ &\quad + \int_{t_0}^t h(\tau)\left[\frac{m}{p}K^{(m-p)/p}(a(\tau) + b(\tau)z(\tau)) + \frac{p-m}{p}K^{m/p}\right]\Delta\tau \\ &\leq B(t) + A(t)z(t), \quad t \in \mathbb{T}^k, \end{aligned} \quad (3.7)$$

where $A(t)$, and $B(t)$ are defined as in (3.3) and $A(t)$ is regressive obviously.

From Lemma 2.3 and (3.7), noting $z(t_0) = 0$, we obtain

$$z(t) \leq \int_{t_0}^t B(\tau)e_{A(\tau)}(t, \sigma(\tau))\Delta\tau. \quad (3.8)$$

Therefore, the desired inequality (3.2) follows from (3.5) and (3.8). \square

Remark 3.2. Theorem 3.1 extends some known inequalities on time scales. If $q = 1, r = 0, h(t) = 0$, then Theorem 3.1 reduces to [7, Theorem 3.1]. If $q = p, h(t) = 0$, then Theorem 3.1 reduces to [8, Theorem 3.2].

Remark 3.3. The result of Theorem 3.1 holds for an arbitrary time scale. If $\mathbb{T} = \mathbb{R}$, then Theorem 3.1 becomes the Theorem 1 established by Yuan et al. [13]. If $\mathbb{T} = \mathbb{Z}$, we can have the following Corollary.

Corollary 3.4. *Let $\mathbb{T} = \mathbb{Z}$ and assume that $u(t), a(t), b(t), f(t), g(t)$, and $h(t)$ are nonnegative functions defined for $t \in \mathbb{N}_0$. Then the inequality*

$$u^p(t) \leq a(t) + b(t) \sum_{s=0}^{t-1} \left[f(s)u^q(s) + g(s)u^r(s) + \sum_{\tau=0}^{s-1} h(\tau)u^m(\tau) \right], \quad t \in \mathbb{N}_0, \quad (3.9)$$

implies

$$u(t) \leq \left\{ a(t) + b(t) \sum_{s=0}^{t-1} \bar{B}(s) \prod_{\tau=s+1}^{t-1} (1 + \bar{A}(\tau)) \right\}^{1/p}, \quad K > 0, \quad t \in \mathbb{N}_0, \quad (3.10)$$

where

$$\begin{aligned} \bar{A}(t) &= \left[\frac{q}{p} K^{(q-p)/p} f(t) + \frac{r}{p} K^{(r-p)/p} g(t) \right] b(t) + \frac{m}{p} K^{(m-p)/p} \sum_{\tau=0}^{t-1} b(\tau) h(\tau), \\ \bar{B}(t) &= f(t) \left[\frac{q}{p} K^{(q-p)/p} a(t) + \frac{p-q}{p} K^{q/p} \right] + g(t) \left[\frac{r}{p} K^{(r-p)/p} a(t) + \frac{p-r}{p} K^{p/r} \right] \\ &\quad + \sum_{\tau=0}^{t-1} \left[\frac{m}{p} K^{(m-p)/p} a(\tau) + \frac{p-m}{p} K^{m/p} \right] h(\tau), \quad t \in \mathbb{N}_0. \end{aligned} \quad (3.11)$$

Corollary 3.5. Let $\mathbb{T} = l\mathbb{Z} \cap [0, \infty)$, where $l\mathbb{Z} = \{lk : k \in \mathbb{Z}, l > 0\}$. We assume that $u(t), a(t), b(t), f(t), g(t)$, and $h(t)$ are nonnegative functions defined for $t \in \mathbb{T}$. Then the inequality

$$u^p(t) \leq a(t) + b(t) \sum_{s=0}^{t/l-1} \left[f(ls) u^q(ls) + g(ls) u^r(ls) + \sum_{\tau=0}^{s/l-1} h(l\tau) u^m(l\tau) \right], \quad t \in \mathbb{T} \quad (3.12)$$

implies

$$u(t) \leq \left\{ a(t) + b(t) \sum_{s=0}^{t/l-1} \bar{B}(ls) \prod_{\tau=s/l+1}^{t/l-1} (1 + \bar{A}(\tau)) \right\}^{1/p}, \quad K > 0, \quad t \in \mathbb{T}, \quad (3.13)$$

where

$$\begin{aligned} \bar{A}(t) &= \left[\frac{q}{p} K^{(q-p)/p} f(t) + \frac{r}{p} K^{(r-p)/p} g(t) \right] b(t) + \frac{m}{p} K^{(m-p)/p} \sum_{\tau=0}^{t/l-1} b(l\tau) h(l\tau), \\ \bar{B}(t) &= f(t) \left[\frac{q}{p} K^{(q-p)/p} a(t) + \frac{p-q}{p} K^{q/p} \right] + g(t) \left[\frac{r}{p} K^{(r-p)/p} a(t) + \frac{p-r}{p} K^{p/r} \right] \\ &\quad + \sum_{\tau=0}^{t/l-1} \left[\frac{m}{p} K^{(m-p)/p} a(l\tau) + \frac{p-m}{p} K^{m/p} \right] h(l\tau), \quad t \in \mathbb{T}. \end{aligned} \quad (3.14)$$

Theorem 3.6. Assume that u, a, b, f, h are defined as in Theorem 3.1, $L(t, y), M(t, y) : \mathbb{T}^k \times \mathbb{R} \rightarrow \mathbb{R}_+$ are continuous functions, and $L(t, y)$ is nondecreasing about the second variable and satisfies

$$0 \leq L(t, x) - L(t, y) \leq M(t, y)(x - y) \quad (3.15)$$

for $t \in \mathbb{T}^k$ and $x \geq y \geq 0$; then

$$u^p(t) \leq a(t) + b(t) \int_{t_0}^t \left[f(s)u^q(s) + L(s, u^r(s)) + \int_{t_0}^s h(\tau)u^m(\tau)\Delta\tau \right] \Delta s, \quad t \in \mathbb{T}^k \quad (3.16)$$

implies

$$u(t) \leq \left\{ a(t) + b(t) \int_{t_0}^t B_1(\tau)e_{A_1}(t, \sigma(\tau))\Delta\tau \right\}^{1/p}, \quad K > 0, \quad t \in \mathbb{T}^k, \quad (3.17)$$

where

$$\begin{aligned} A_1(t) &= \frac{q}{p}K^{(q-p)/p}f(t)b(t) + \frac{m}{p}K^{(m-p)/p} \int_{t_0}^t h(\tau)b(\tau)\Delta\tau \\ &\quad + \frac{r}{p}K^{(r-p)/p}M\left(t, \frac{r}{p}K^{(r-p)/p}a(t) + \frac{p-r}{p}K^{r/p}\right)b(t), \\ B_1(t) &= f(t)\left(\frac{q}{p}K^{(q-p)/p}a(t) + \frac{p-q}{p}K^{q/p}\right) \\ &\quad + \int_{t_0}^t h(\tau)\left(\frac{m}{p}K^{(m-p)/p}a(\tau) + \frac{p-m}{p}K^{m/p}\right)\Delta\tau \\ &\quad + L\left(t, \frac{r}{p}K^{(r-p)/p}a(t) + \frac{p-r}{p}K^{r/p}\right), \quad t \in \mathbb{T}^k. \end{aligned} \quad (3.18)$$

Proof. Define $z(t)$ by

$$z(t) = \int_{t_0}^t \left[f(s)u^q(s) + L(s, u^r(s)) + \int_{t_0}^s h(\tau)u^m(\tau)\Delta\tau \right] \Delta s, \quad (3.19)$$

then $z(t_0) = 0$, and (3.16) can be written as (3.5).

Therefore, from (3.6) and (3.19), we have

$$\begin{aligned}
z^\Delta(t) &= f(t)u^q(t) + L(t, u^r(t)) + \int_{t_0}^t h(\tau)u^m(\tau)\Delta\tau \\
&\leq f(t) \left[\frac{q}{p}K^{(q-p)/p}(a(t) + b(t)z(t)) + \frac{p-q}{p}K^{q/p} \right] \\
&\quad + L \left(t, \frac{r}{p}K^{(r-p)/p}(a(t) + b(t)z(t)) + \frac{p-r}{p}K^{r/p} \right) \\
&\quad - L \left(t, \frac{r}{p}K^{(r-p)/p}a(t) + \frac{p-r}{p}K^{r/p} \right) \\
&\quad + \int_{t_0}^t h(\tau) \left[\frac{m}{p}K^{(m-p)/p}(a(\tau) + b(\tau)z(\tau)) + \frac{p-m}{p}K^{m/p} \right] \Delta\tau \\
&\quad + L \left(t, \frac{r}{p}K^{(r-p)/p}a(t) + \frac{p-r}{p}K^{r/p} \right) \\
&\leq f(t) \left[\frac{q}{p}K^{(q-p)/p}a(t) + \frac{p-q}{p}K^{q/p} \right] + \int_{t_0}^t h(\tau) \left(\frac{m}{p}K^{(m-p)/p}a(\tau) + \frac{p-m}{p}K^{m/p} \right) \Delta\tau \\
&\quad + \left[\frac{q}{p}K^{(q-p)/p}f(t)b(t) + \frac{m}{p}K^{(m-p)/p} \int_{t_0}^t h(\tau)b(\tau)\Delta\tau \right] z(t) \\
&\quad + M \left(t, \frac{r}{p}K^{(r-p)/p}a(t) + \frac{p-r}{p}K^{r/p} \right) \frac{r}{p}K^{(r-p)/p}b(t)z(t) \\
&\quad + L \left(t, \frac{r}{p}K^{(r-p)/p}a(t) + \frac{p-r}{p}K^{r/p} \right) = A_1(t)z(t) + B_1(t), \quad t \in \mathbb{T}^k,
\end{aligned} \tag{3.20}$$

where $A_1(t)$, and $B_1(t)$ are defined as in (3.18) and $A_1(t)$ is regressive obviously.

From Lemma 2.3 and (3.20), noting $z(t_0) = 0$, we obtain

$$z(t) \leq \int_{t_0}^t B_1(\tau)e_{A_1(\tau)}(t, \sigma(\tau))\Delta\tau. \tag{3.21}$$

Therefore, the desired inequality (3.17) follows from (3.5) and (3.21). \square

Remark 3.7. If $\mathbb{T} = \mathbb{R}$, then Theorem 3.6 becomes [13, Theorem 3]. If $\mathbb{T} = \mathbb{Z}$, we can have the following Corollary.

Corollary 3.8. Let $\mathbb{T} = \mathbb{Z}$ and assume that $u(t), a(t), b(t), f(t), g(t)$, and $h(t)$ are nonnegative functions defined for $t \in \mathbb{N}_0$. $L, M \in C(\mathbb{R}_+^2, \mathbb{R}_+)$ satisfy

$$0 \leq L(t, x) - L(t, y) \leq M(t, y)(x - y) \quad (3.22)$$

for $x \geq y \geq 0$ and $L(t, y)$ is nondecreasing about the second variable. Then the inequality

$$u^p(t) \leq a(t) + b(t) \sum_{s=0}^{t-1} \left[f(s)u^q(s) + L(s, u^r(s)) + \sum_{\tau=0}^{s-1} h(\tau)u^m(\tau) \right], \quad t \in \mathbb{N}_0 \quad (3.23)$$

implies

$$u(t) \leq \left\{ a(t) + b(t) \sum_{s=0}^{t-1} \bar{B}_1(\tau) \prod_{\tau=s+1}^{t-1} (1 + \bar{A}_1(\tau)) \right\}^{1/p}, \quad K > 0, \quad t \in \mathbb{N}_0, \quad (3.24)$$

where

$$\begin{aligned} \bar{A}_1(t) &= \frac{q}{p} K^{(q-p)/p} f(t)b(t) + \frac{m}{p} K^{(m-p)/p} \sum_{\tau=0}^{t-1} h(\tau)b(\tau) \\ &\quad + \frac{r}{p} K^{(r-p)/p} M \left(t, \frac{r}{p} K^{(r-p)/p} a(t) + \frac{p-r}{p} K^{r/p} \right) b(t), \\ \bar{B}_1(t) &= f(t) \left(\frac{q}{p} K^{(q-p)/p} a(t) + \frac{q-p}{p} K^{q/p} \right) + \sum_{\tau=0}^{t-1} h(\tau) \left(\frac{m}{p} K^{(m-p)/p} a(\tau) + \frac{p-m}{p} K^{m/p} \right) \\ &\quad + L \left(t, \frac{r}{p} K^{(r-p)/p} a(t) + \frac{p-r}{p} K^{r/p} \right), \quad t \in \mathbb{N}_0. \end{aligned} \quad (3.25)$$

Theorem 3.9. Assume that $u(t), a(t), b(t), f(t), g(t)$, and $h(t)$ are defined as in Theorem 3.1, $w(t, s)$ is defined as in Lemma 2.2 such that $w(\sigma(t), t) \geq 0, w^\Delta(t, s) \geq 0$ for $t, s \in \mathbb{T}$ with $s \leq t$; then

$$u^p(t) \leq a(t) + b(t) \int_{t_0}^t w(t, s) \left[f(s)u^q(s) + g(s)u^r(s) + \int_{t_0}^s h(\tau)u^m(\tau)\Delta\tau \right] \Delta s, \quad t \in \mathbb{T}^k, \quad (3.26)$$

implies

$$u(t) \leq \left\{ a(t) + b(t) \int_{t_0}^t B_2(\tau) e_{A_2}(t, \sigma(\tau)) \Delta\tau \right\}^{1/p}, \quad K > 0, \quad t \in \mathbb{T}^k, \quad (3.27)$$

where

$$\begin{aligned} A_2(t) &= w(\sigma(t), t) \left[\left(\frac{q}{p} K^{(q-p)/p} f(t) + \frac{r}{p} K^{(r-p)/p} g(t) \right) b(t) + \frac{m}{p} K^{(m-p)/p} \int_{t_0}^t b(\tau) h(\tau) \Delta\tau \right] \\ &\quad + \int_{t_0}^t w^\Delta(t, s) \left[\left(\frac{q}{p} K^{(q-p)/p} f(s) + \frac{r}{p} K^{(r-p)/p} g(s) \right) b(s) \right. \\ &\quad \left. + \frac{m}{p} K^{(m-p)/p} \int_{t_0}^s b(\tau) h(\tau) \Delta\tau \right] \Delta s, \\ B_2(t) &= w(\sigma(t), t) \left[f(t) \left(\frac{q}{p} K^{(q-p)/p} a(t) + \frac{p-q}{p} K^{q/p} \right) + g(t) \left(\frac{r}{p} K^{(r-p)/p} a(t) + \frac{p-r}{p} K^{p/r} \right) \right. \\ &\quad \left. + \int_{t_0}^t h(\tau) \left(\frac{m}{p} K^{(m-p)/p} a(\tau) + \frac{p-m}{p} K^{m/p} \right) \Delta\tau \right] \\ &\quad + \int_{t_0}^t w^\Delta(t, s) \left[f(s) \left(\frac{q}{p} K^{(q-p)/p} a(s) + \frac{p-q}{p} K^{q/p} \right) \right. \\ &\quad \left. + g(s) \left(\frac{r}{p} K^{(r-p)/p} a(s) + \frac{p-r}{p} K^{p/r} \right) \right. \\ &\quad \left. + \int_{t_0}^s h(\tau) \left(\frac{m}{p} K^{(m-p)/p} a(\tau) + \frac{p-m}{p} K^{m/p} \right) \Delta\tau \right] \Delta s, \quad t \in \mathbb{T}^k. \end{aligned} \quad (3.28)$$

Proof. Define $z(t)$ by

$$z(t) = \int_{t_0}^t w(t, s) \left[f(s) u^q(s) + g(s) u^r(s) + \int_{t_0}^s h(\tau) u^m(\tau) \Delta\tau \right] \Delta s, \quad (3.29)$$

then $z(t_0) = 0$, and (3.26) can be written as (3.5).

Therefore, from (3.6) and (3.29) we have

$$\begin{aligned}
z^\Delta(t) &= w(\sigma(t), t) \left[f(t)u^q(t) + g(t)u^r(t) + \int_{t_0}^t h(\tau)u^m(\tau)\Delta\tau \right] \\
&\quad + \int_{t_0}^t w^\Delta(t, s) \left[f(s)u^q(s) + g(s)u^r(s) + \int_{t_0}^s h(\tau)u^m(\tau)\Delta\tau \right] \Delta s \\
&\leq w(\sigma(t), t) \left[f(t) \left(\frac{q}{p} K^{(q-p)/p} (a(t) + b(t)z(t)) + \frac{p-q}{p} K^{q/p} \right) \right. \\
&\quad \left. + g(t) \left(\frac{r}{p} K^{(r-p)/p} (a(t) + b(t)z(t)) + \frac{p-r}{p} K^{r/p} \right) \right. \\
&\quad \left. + \int_{t_0}^t h(\tau) \left(\frac{m}{p} K^{(m-p)/p} (a(\tau) + b(\tau)z(\tau)) + \frac{p-m}{p} K^{m/p} \right) \right] \Delta\tau \quad (3.30) \\
&\quad + \int_{t_0}^t w^\Delta(t, s) \left[f(s) \left(\frac{q}{p} K^{(q-p)/p} (a(s) + b(s)z(s)) + \frac{p-q}{p} K^{q/p} \right) \right. \\
&\quad \left. + g(s) \left(\frac{r}{p} K^{(r-p)/p} (a(s) + b(s)z(s)) + \frac{p-r}{p} K^{r/p} \right) \right. \\
&\quad \left. + \int_{t_0}^s h(\tau) \left(\frac{m}{p} K^{(m-p)/p} (a(\tau) + b(\tau)z(\tau)) + \frac{p-m}{p} K^{m/p} \right) \right] \Delta s \\
&\leq B_2(t) + A_2(t)z(t), \quad t \in \mathbb{T}^k,
\end{aligned}$$

where $A_2(t)$, and $B_2(t)$ are defined by (3.28) and $A_2(t)$ is regressive obviously.

From Lemma 2.3 and (3.30), noting $z(t_0) = 0$, we obtain

$$z(t) \leq \int_{t_0}^t B_2(\tau) e_{A_2(\tau)}(t, \sigma(\tau)) \Delta\tau. \quad (3.31)$$

Therefore, the desired inequality (3.27) follows from (3.5) and (3.31). \square

Remark 3.10. If $q = p$, $h(t) = 0$, then Theorem 3.9 reduces to [8, Theorem 3.8].

Using our results, we can also obtain many dynamic inequalities for some peculiar time scales; here, we omit them.

4. Some Applications

In this section, we present some applications of Theorem 3.9 to investigate certain properties of solution $u(t)$ of the following dynamic equation:

$$[u^p(t)]^\Delta = F\left(t, U(t, u(t)), \int_{t_0}^t H(s, u(s)) \Delta s\right), \quad u^p(t_0) = C, \quad t \in \mathbb{T}^k, \quad (4.1)$$

where C is a constant, $F : \mathbb{T}^k \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function, and $U : \mathbb{T}^k \times \mathbb{R} \rightarrow \mathbb{R}$, $H : \mathbb{T}^k \times \mathbb{R} \rightarrow \mathbb{R}$ are also continuous functions.

Example 4.1. Assume that

$$\begin{aligned} |F(t, U, V)| &\leq |U| + |V|, \\ |U(t, u)| &\leq f(t)|u|^q + g(t)|u|^r, \\ |H(t, u)| &\leq h(t)|u|^m, \quad t \in \mathbb{T}^k, \end{aligned} \quad (4.2)$$

where p, q, r , and m are constants, $p \geq q > 0$, and $p \geq m > 0, p \geq r > 0$. $f, g, h \in C_{rd}$, $f(t), g(t)$ and $h(t)$ are nonnegative. Then every solution $u(t)$ of (4.1) satisfies

$$|u(t)| \leq \left\{ |C| + \int_{t_0}^t B(\tau) e_{A(\tau)}(t, \sigma(\tau)) \Delta\tau \right\}^{1/p}, \quad K > 0, \quad t \in \mathbb{T}^k, \quad (4.3)$$

where A, B are defined as in (3.3) with $a(t) = |C|, b(t) = 1$.

Indeed, the solution $u(t)$ of (4.1) satisfies the following equivalent equation

$$u^p(t) = C + \int_{t_0}^t F\left(\tau, U(\tau, u(\tau)), \int_{t_0}^{\tau} H(s, u(s)) \Delta s\right) \Delta\tau, \quad t \in \mathbb{T}^k. \quad (4.4)$$

It follows from (4.2) and (4.4) that

$$\begin{aligned} |u^p(t)| &\leq |C| + \int_{t_0}^t \left| F\left(\tau, U(\tau, u(\tau)), \int_{t_0}^{\tau} H(s, u(s)) \Delta s\right) \right| \Delta\tau \\ &\leq |C| + \int_{t_0}^t \left[f(\tau)|u(\tau)|^q + g(\tau)|u(\tau)|^r + \int_{t_0}^{\tau} h(s)|u(s)|^m \Delta s \right] \Delta\tau. \end{aligned} \quad (4.5)$$

Using Theorem 3.1, the inequality (4.3) is obtained from (4.5).

Example 4.2. Assume that

$$\begin{aligned} |F(t, U_1, V_1) - F(t, U_2, V_2)| &\leq |U_1 - U_2| + |V_1 - V_2|, \\ |U(t, u_1) - U(t, u_2)| &\leq f(t) \left| u_1^p - u_2^p \right|, \\ |H(t, u_1) - H(t, u_2)| &\leq h(t) \left| u_1^p - u_2^p \right|, \quad t \in \mathbb{T}^k, \end{aligned} \quad (4.6)$$

p, f , and h are defined as in Example 4.1. If $p = (m/n)$ ($m, n \in \mathbb{N}$) and m is odd, then (4.1) has at most one solution; otherwise, the two solutions $u_1(t)$, and $u_2(t)$ of (4.1) have the relation $u_1^p(t) = u_2^p(t)$.

Proof. Let $u_1(t)$, and $u_2(t)$ be two solutions of (4.1). Then we have

$$u_1^p(t) - u_2^p(t) = \int_{t_0}^t \left[F \left(\tau, U(\tau, u_1(\tau)), \int_{t_0}^{\tau} H(s, u_1(s)) \Delta s \right) - F \left(\tau, U(\tau, u_2(\tau)), \int_{t_0}^{\tau} H(s, u_2(s)) \Delta s \right) \right] \Delta \tau, \quad t \in \mathbb{T}^k. \quad (4.7)$$

It follows from (4.6) and (4.7) that

$$|u_1^p(t) - u_2^p(t)| \leq \int_{t_0}^t \left[f(\tau) |u_1^p(\tau) - u_2^p(\tau)| + \int_{t_0}^{\tau} h(s) |u_1^p(s) - u_2^p(s)| \Delta s \right] \Delta \tau, \quad t \in \mathbb{T}^k. \quad (4.8)$$

By Theorem 3.1, we have $u_1^p(t) - u_2^p(t) \equiv 0$, $t \in \mathbb{T}^k$. The results are obtained. \square

Example 4.3. Consider the equation

$$u^p = a(t) + b(t) \int_{t_0}^t F \left(t, s, U(s, u), \int_{t_0}^s H(\tau, u) \Delta \tau \right) \Delta s, \quad t \in \mathbb{T}^k. \quad (4.9)$$

If

$$\begin{aligned} |F(t, s, U, V)| &\leq w(t, s)(|U| + |V|), \\ |U(t, u)| &\leq f(t)|u|^q + g(t)|u|^r, \\ |H(t, u)| &\leq h(t)|u|^m, \quad t \in \mathbb{T}^k, \end{aligned} \quad (4.10)$$

where p, q, r, m are constants, $p \geq q > 0$, $p \geq m > 0$, $p \geq r > 0$. $a, b, f, g, h \in C_{rd}$, $a(t), b(t), f(t), g(t)$ and $h(t)$ are nonnegative, $w(t, s)$ is defined as in Lemma 2.2 such that $w(\sigma(t), t) \geq 0$, $w^\Delta(t, s) \geq 0$ for $t, s \in \mathbb{T}$ with $s \leq t$.

Then we have the estimate of the solution $u(t)$ of (4.9) that

$$|u(t)| \leq \left\{ a(t) + b(t) \int_{t_0}^t B_2(\tau) e_{A_2}(t, \sigma(\tau)) \Delta \tau \right\}^{1/p}, \quad K > 0, \quad t \in \mathbb{T}^k, \quad (4.11)$$

where A_2, B_2 are defined as in (3.28).

Proof. From (4.10) and (4.9), we have

$$|u(t)|^p \leq a(t) + b(t) \int_{t_0}^t w(t, s) \left[f(s)|u(s)|^q + g(s)|u(s)|^r + \int_{t_0}^s h(\tau)|u(\tau)|^m \Delta \tau \right] \Delta s, \quad t \in \mathbb{T}^k. \quad (4.12)$$

By Theorem 3.9 and (4.12), we have that (4.11) holds. \square

Acknowledgments

The authors thank the referees very much for their careful comments and valuable suggestions on this paper. This research is supported by the National Natural Science Foundation of China (10771118), the Natural Science Foundation of Shandong Province (ZR2009AM011), and the Science Foundation of the Education Department of Shandong Province, China (J07yh05).

References

- [1] S. Hilger, "Analysis on measure chains—a unified approach to continuous and discrete calculus," *Results in Mathematics*, vol. 18, no. 1-2, pp. 18–56, 1990.
- [2] R. Agarwal, M. Bohner, and A. Peterson, "Inequalities on time scales: a survey," *Mathematical Inequalities & Applications*, vol. 4, no. 4, pp. 535–557, 2001.
- [3] E. Akin-Bohner, M. Bohner, and F. Akin, "Pachpatte inequalities on time scales," *Journal of Inequalities in Pure Applied Mathematics*, vol. 6, no. 1, article 6, 23 pages, 2005.
- [4] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Applications*, An introduction with application, Birkhäuser, Boston, Mass, USA, 2001.
- [5] M. Bohner and A. Peterson, *Advances in Dynamic Equations on Time Scales*, Birkhäuser, Boston, Mass, USA, 2003.
- [6] D. B. Pachpatte, "Explicit estimates on integral inequalities with time scale," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 7, no. 4, article 143, p. 8, 2006.
- [7] W. N. Li, "Some new dynamic inequalities on time scales," *Journal of Mathematical Analysis and Applications*, vol. 319, no. 2, pp. 802–814, 2006.
- [8] W. N. Li and W. Sheng, "Some nonlinear integral inequalities on time scales," *Journal of Inequalities and Applications*, Article ID 70465, 15 pages, 2007.
- [9] W. N. Li, "Some Pachpatte type inequalities on time scales," *Computers & Mathematics with Applications*, vol. 57, no. 2, pp. 275–282, 2009.
- [10] W. N. Li, "Some delay integral inequalities on time scales," *Computers and Mathematics with Applications*, vol. 59, no. 6, pp. 1929–1936, 2010.
- [11] W. N. Li, "Bounds for certain new integral inequalities on time scales," *Advances in Difference Equations*, vol. 2009, Article ID 484185, 16 pages, 2009.
- [12] D. R. Anderson, "Nonlinear dynamic integral inequalities in two independent variables on time scale Pairs," *Advances in Dynamical Systems and Applications*, vol. 3, no. 1, pp. 1–13, 2008.
- [13] Z. Yuan, X. Yuan, F. Meng, and H. Zhang, "Some new delay integral inequalities and their applications," *Applied Mathematics and Computation*, vol. 208, no. 1, pp. 231–237, 2009.