Research Article

A Fixed Point Approach to the Stability of Pexider Quadratic Functional Equation with Involution

**M. M. Pourpasha,¹ J. M. Rassias,² R. Saadati,³
and S. M. Vaezpour³**

¹ Department of Mathematics, Science and Research Branch, Islamic Azad University (IAU), Tehran, Iran

² Section of Mathematics and Informatics, Pedagogical Department, National and Kapodistrian University of Athens, 4, Agamemnonos St., Aghia Paraskevi, Athens 15342, Greece

³ Department of Mathematics, Amirkabir University of Technology, Hafez Avenue, P. O. Box 15914, Tehran, Iran

Correspondence should be addressed to

R. Saadati, rsaadati@eml.cc and S. M. Vaezpour, vaezpour@aut.ac.ir

Received 10 May 2010; Revised 13 July 2010; Accepted 31 July 2010

Academic Editor: S. Reich

Copyright © 2010 M. M. Pourpasha et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We apply the fixed point method to investigate the Hyers-Ulam stability of the Pexider functional equation $f(x + y) + g(x + \sigma(y)) = h(x) + k(y)$, for all $x, y \in E$, where E is a normed space and $\sigma : E \rightarrow E$ is an involution.

1. Introduction and Preliminary

A basic question in the theory of functional equations is as follows. “When is it true that a function, which approximately satisfies a functional equation must be close to an exact solution of the equation?” The first stability problem concerning group homomorphisms was raised by Ulam [1] in 1940 and affirmatively answered by Hyers in [2]. Subsequently, the result of Hyers was generalized by Aoki [3] for additive mappings and by Rassias [4] for linear mappings by considering an unbounded Cauchy difference. The paper of Rassias has provided a lot of influence in the development of what we now call Hyers-Ulam-Rassias stability of functional equations. For more information, see [5–7]. Specially, Maligranda [8] and Moszner [9] provided a very interesting discussion on the definition of functional equations’ stability.

Recently, the stability of functional equations has been investigated by many mathematicians. They have many applications in the Information Theory, Physics, Economic Theory and Social and Behavior Sciences. See [10–14].

A Hyers-Ulam stability theorem for the quadratic functional equation

$$f(x+y) + f(x-y) = 2f(x) + 2f(y) \quad (1.1)$$

was proved by Skof [15] for the function $f : E_1 \rightarrow E_2$, where E_1 is a normed space and E_2 is a Banach space. Cholewa [16] noticed that the theorem of Skof is still true if the relevant domain E_1 is replaced by an abelian group. Czerwik [17] proved the generalized Hyers-Ulam stability of the quadratic functional equation (1.1). Recently, Brzdęk [18], Jung [19], and Jung and Sahoo [20] investigated the Hyers-Ulam-Rassias stability of (1.1). Furthermore they proved the Hyers-Ulam-Rassias stability of the functional equation of Pexider type

$$f_1(x+y) + f_2(x-y) = f_3(x) + f_4(y). \quad (1.2)$$

The stability problem of several functional equations has been extensively investigated by a number of authors, and there are many interesting results concerning this problem (see [4, 21–37]).

Let E_1 and E_2 be real vector spaces. If an additive function $\sigma : E_1 \rightarrow E_1$ satisfies $\sigma(x+y) = \sigma(x) + \sigma(y)$ and $\sigma(\sigma(x)) = x$ for all $x, y \in E_1$, then σ is called an involution of E_1 , see [21, 37]. For a given involution $\sigma : E_1 \rightarrow E_1$, the functional equation

$$f(x+y) + f(x+\sigma(y)) = 2f(x) + 2f(y), \quad \forall x, y \in E \quad (1.3)$$

is called the quadratic functional equation with involution. According to [37, Corollary 8], a function $f : E_1 \rightarrow E_2$ is a solution of (1.3) if and only if there exists an additive function $A : E_1 \rightarrow E_2$, and a biadditive symmetric function $B : E_1 \times E_1 \rightarrow E_2$ such that $A(\sigma(x)) = A(x)$, $B(\sigma(x), y) = -B(\sigma(x), y)$ and $f(x) = B(x, y) + A(x)$ for all $x \in E_1$.

Indeed, if we set $\sigma(x) = I$ in (1.3), where $I : E_1 \rightarrow E_1$ denotes the identity function, then (1.3) reduces to the additive functional equation

$$f(x+y) = f(x) + f(y), \quad \forall x, y \in E. \quad (1.4)$$

On the other hand, if $\sigma(x) = -I$ in (1.3), then (1.3) is transformed into the quadratic functional equation

$$f(x+y) + f(x-y) = 2f(x) + 2f(y), \quad \forall x, y \in E. \quad (1.5)$$

Recently, Bouikhalene et al. have proved the Hyers-Ulam-Rassias stability of the quadratic functional equation with involution (1.2), see [21].

In this paper, we will apply the fixed point method to prove the Hyers-Ulam-Rassias stability of the functional equation (1.3) in the Pexider type

$$f(x+y) + g(x+\sigma(y)) = 2h(x) + 2k(y). \quad (1.6)$$

To see the different approaches to the problem of the Pexiderized Cauchy equations' stability and further references concerning that subject we refer to [38–43].

Let X be a set. A function $d : X \times X \rightarrow [0, \infty]$ is called a generalized metric on X if and only if d satisfies the following

- (1) $d(x, y) = 0$, if and only if $x = y$
- (2) $d(x, y) = d(y, x)$, for all $x, y \in X$;
- (3) $d(x, z) \leq d(x, y) + d(y, z)$, for all $x, y, z \in X$.

For an extensive theory of fixed point and other nonlinear methods, the reader is referred to the book of Hyers et al. [44] and Cădariu and Radu [45].

Theorem 1.1. *Let (X, d) be a generalized complete metric space. Assume that $J : X \rightarrow X$ is a strictly contractive operator with the Lipschitz constant $0 < L < 1$. If there exists a nonnegative integer k such that $d(J^{k+1}x, J^kx) < \infty$ for some $x \in X$, then the following are true:*

- (a) *the sequence $\{J^n x\}$ converges to a fixed point x^* of J*
- (b) *x^* is the unique fixed point of J in*

$$X^* = \{y \in X : d(J^k x, x) < \infty\}; \quad (1.7)$$

- (c) *if $y \in X^*$, then*

$$d(y, x^*) \leq \frac{1}{1-L} d(Jy, y). \quad (1.8)$$

2. Main Results

In this section, we prove the Hyers-Ulam-Rassias stability of the quadratic functional equation with involution (1.6) by applying the fixed point method.

Theorem 2.1. *Let E_1 be a commutative semigroup (with the divisibility by 2), and let E_2 be a real Banach space. Suppose that a function $\varphi : E_1 \times E_1 \rightarrow [0, \infty)$ is given and there exists a constant L , $0 < L < 1$, such that*

$$\begin{aligned} \varphi(2x, 2y) &\leq 2L\varphi(x, y), \\ \varphi(x + \sigma(x), y + \sigma(y)) &\leq 2L\varphi(x, y), \end{aligned} \quad (2.1)$$

for all $x, y \in E_1$. Furthermore, let $f, g, h, k : E_1 \rightarrow E_2$ be even functions satisfying the inequality

$$\|f(x+y) + g(x+\sigma(y)) - 2h(x) - 2k(y)\| \leq \varphi(x, y) \quad (2.2)$$

for all $x, y \in E_1$, where $\sigma : E_1 \rightarrow E_1$ is an involution of E_1 and $f(0) = g(0) = h(0) = k(0) = 0$. Then there exists a unique solution $T : E_1 \rightarrow E_2$ of (2.2) such that

$$\begin{aligned} \|2f(x) - T(x)\| &\leq \frac{1}{4(1-L)} M'(x, x) + M(x, 0), \\ \|2g(x) - T(x)\| &\leq \frac{1}{4(1-L)} M'(x, x) + M(x, 0) + \frac{1}{2}(\varphi(x, x) + \varphi(x, -x)), \\ \|h(x) - T(x)\| &\leq \frac{1}{4(1-L)} M'(x, x), \\ \|k(x) - T(x)\| &\leq \frac{1}{4(1-L)} M'(x, x) + \frac{1}{4}(\varphi(0, x) + \varphi(x, 0)), \end{aligned} \quad (2.3)$$

for all $x \in E_1$, where

$$\begin{aligned} M(x, y) &= \varphi(x, y) + \varphi(0, y) + \varphi(y, 0) + \varphi\left(\frac{x}{2}, \frac{x}{2}\right) + \varphi\left(\frac{x}{2}, -\left(\frac{x}{2}\right)\right), \\ M'(x, y) &= M(x, y) + M(x + y, 0) + M(x + \sigma(y), 0), \\ T(x) &= \lim_{n \rightarrow \infty} \frac{1}{2^{2n}} \left[h(2^n x) + (2^n - 1)h\left(2^{n-1}x + 2^{n-1}\sigma(x)\right) \right]. \end{aligned} \quad (2.4)$$

Proof. Letting $y = 0$ in (2.2), we obtain

$$\|f(x) + g(x) - 2h(x)\| \leq \varphi(x, 0). \quad (2.5)$$

Similarly, for every $y \in E_1$, we can put $x = 0$ in (2.2) to obtain

$$\|f(y) + g(\sigma(y)) - 2k(y)\| \leq \varphi(0, y). \quad (2.6)$$

Since $\sigma : E_1 \rightarrow E_1$ is an involution, we replace y by $x + \sigma(x) := d$ in (2.6), then we have

$$\|f(d) + g(d) - 2k(d)\| \leq \varphi(0, d). \quad (2.7)$$

Also, we can replace y by x and replace y by $-x$ in (2.2) to get

$$\|f(2x) + g(x + \sigma(x)) - 2h(x) - 2k(x)\| \leq \varphi(x, x), \quad (2.8)$$

$$\|g(x - \sigma(x)) - 2h(x) - 2k(-x)\| \leq \varphi(x, -x). \quad (2.9)$$

Since $\sigma : E_1 \rightarrow E_1$ is an involution, by replacing x by $x + \sigma(x) := d$ in (2.8) and using (2.1), we have

$$\|f(2d) - 2h(d) - 2k(d)\| \leq \varphi(d, d). \quad (2.10)$$

Also, by replacing x by $x - \sigma(x) := d$ in (2.9), we have

$$\|g(2d) - 2h(d) - 2k(d)\| \leq \varphi(d, -d). \quad (2.11)$$

In view of (2.5) and (2.7), we see that

$$\|h(d) - k(d)\| \leq \frac{1}{2}(\varphi(0, d) + \varphi(d, 0)), \quad (2.12)$$

and it follows from (2.10) and (2.11) that

$$\|f(2d) - g(2d)\| \leq \varphi(d, d) + \varphi(d, -d). \quad (2.13)$$

By using (2.2), (2.12) and (2.13), we have

$$\begin{aligned} & \|f(x+y) + f(x+\sigma(y)) - 2h(x) - 2h(y)\| \\ & \leq \|f(x+y) + g(x+\sigma(y)) - 2h(x) - 2k(y)\| + 2\|k(y) - h(y)\| \\ & \quad + \|f(x+\sigma(y)) - g(x+\sigma(y))\| \\ & \leq \varphi(x, y) + \varphi(0, y) + \varphi(y, 0) + \varphi\left(\frac{x+\sigma(y)}{2}, \frac{x+\sigma(y)}{2}\right) \\ & \quad + \varphi\left(\frac{x+\sigma(y)}{2}, -\left(\frac{x+\sigma(y)}{2}\right)\right) \\ & \leq \varphi(x, y) + \varphi(0, y) + \varphi(y, 0) + \varphi\left(\frac{x}{2}, \frac{x}{2}\right) + \varphi\left(\frac{x}{2}, -\left(\frac{x}{2}\right)\right) = M(x, y). \end{aligned} \quad (2.14)$$

Therefore

$$\|f(x+y) + f(x+\sigma(y)) - 2h(x) - 2h(y)\| \leq M(x, y). \quad (2.15)$$

By putting $y = 0$ in (2.15), we get

$$\|f(x) - h(x)\| \leq M(x, 0). \quad (2.16)$$

Hence, (2.15) and (2.16) imply that

$$\begin{aligned} & \|h(x+y) + h(x+\sigma(y)) - 2h(x) - 2h(y)\| \\ & \leq \|f(x+y) + f(x+\sigma(y)) - 2h(x) - 2h(y)\| + \|f(x+y) - h(x+y)\| \\ & \quad + \|f(x+\sigma(y)) - h(x+\sigma(y))\| \\ & \leq M(x, y) + M(x+y, 0) + M(x+\sigma(y), 0) = M'(x, y). \end{aligned} \quad (2.17)$$

Therefore

$$\|h(x+y) + h(x+\sigma(y)) - 2h(x) - 2h(y)\| \leq M'(x, y). \quad (2.18)$$

Now, we define X to be the set of all functions $f : E_1 \rightarrow E_2$ and introduce a generalized metric on X as follows:

$$d(g, h) = \inf\{C \in [0, \infty) : \|g(x) - h(x)\| \leq CM'(x, x), \forall x \in E_1\}. \quad (2.19)$$

Let $\{f_n\}$ be a Cauchy sequence in (X, d) . According to the definition of Cauchy sequences, for any given $\epsilon > 0$, there exists a positive integer N_ϵ such that

$$d(f_m, f_n) \leq \epsilon \quad (2.20)$$

for all $m, n \geq N_\epsilon$.

By considering the definition of the generalized metric d , we see that

$$\forall \epsilon > 0, \quad \exists N_\epsilon \in \mathbb{N}, \quad \forall m, n \geq N_\epsilon, \quad \forall x \in E_1 : \|f_m(x) - f_n(x)\| \leq \epsilon M'(x, x). \quad (2.21)$$

If x is any given point in E_1 , (2.21) implies that $\{f_n(x)\}$ is a Cauchy sequence in E_2 . Since E_2 is complete, $\{f_n(x)\}$ converges in E_2 for each $x \in E_1$. Hence, we can define a function $f : E_1 \rightarrow E_2$ by

$$f(x) = \lim_{n \rightarrow \infty} f_n(x). \quad (2.22)$$

We define an operator $J : X \rightarrow X$ by

$$JL(x) = \frac{1}{4}[L(2x) + L(x + \sigma(x))] \quad (2.23)$$

for all $x \in E_1$.

First, we assert that J is strictly contractive on X . Given $g, h \in X$, let $C \in [0, \infty)$ be an arbitrary constant with

$$d(g, h) \leq C, \quad (2.24)$$

that is,

$$\|g(x) - h(x)\| \leq CM'(x, x) \quad (2.25)$$

for all $x \in E_1$.

If we replace y by x in (2.18), then we obtain

$$\|h(2x) + h(x + \sigma(x)) - 4h(x)\| \leq M'(x, x) \quad (2.26)$$

for every $x \in E_1$.

It follows from (2.23) and (2.25) that

$$\begin{aligned} \|Jg(x) - Jh(x)\| &= \frac{1}{4} \|g(2x) + g(x + \sigma(x)) - h(2x) - h(x + \sigma(x))\| \\ &\leq \frac{1}{4} \|g(2x) - h(2x)\| + \frac{1}{4} \|g(x + \sigma(x)) - h(x + \sigma(x))\| \\ &\leq \frac{1}{4} CM'(2x, 2x) + \frac{1}{4} CM'(x + \sigma(x), x + \sigma(x)) \\ &\leq LCM'(x, x) \end{aligned} \quad (2.27)$$

for all $x \in E_1$, that is, $d(Jg, Jh) \leq LC$. Hence, we conclude that $d(Jg, Jh) \leq Ld(g, h)$ for any $g, h \in X$. Therefore, J is strictly contractive because L is a constant with $0 < L < 1$.

Now, we claim that $d(Jh, h) < \infty$. If we put $y = x$ in (2.26) and divide both sides by $1/4$, then we get

$$\|Jh(x) - h(x)\| = \left\| \frac{1}{4} [h(2x) + h(x + \sigma(x)) - h(x)] \right\| \leq \frac{1}{4} M'(x, x) \quad (2.28)$$

for all $x \in E_1$, that is,

$$d(Jh, h) \leq \frac{1}{4} < \infty. \quad (2.29)$$

Now, by Theorem 1.1 there exists a function $T : E_1 \rightarrow E_2$ which is a fixed point of J , such that $d(J^n h, T) \rightarrow 0$ as $n \rightarrow \infty$. By induction, we can easily show that

$$(J^n h)(x) = \frac{1}{2^{2n}} \left[h(2^n x) + (2^n - 1)h(2^{n-1}x + 2^{n-1}\sigma(x)) \right] \quad (2.30)$$

for each $n \in \mathbb{N}$. Since $d(J^n h, T) \rightarrow 0$ as $n \rightarrow \infty$, there exists a sequence $\{C_n\}$ such that $C_n \rightarrow 0$ as $n \rightarrow \infty$ and $d(J^n h, T) \leq C_n$ for every $n \in \mathbb{N}$. Hence, by the definition of d , we have

$$\|J^n h(x) - T(x)\| \leq C_n M'(x, x) \quad (2.31)$$

for all $x \in E_1$. Thus, for each fixed $x \in E_1$, we have

$$\lim_{n \rightarrow \infty} \|J^n h(x) - T(x)\| = 0. \quad (2.32)$$

Therefore

$$T(x) = \lim_{n \rightarrow \infty} \frac{1}{2^{2n}} \left[h(2^n x) + (2^n - 1)h\left(2^{n-1}x + 2^{n-1}\sigma(x)\right) \right] \quad (2.33)$$

for all $x \in E_1$. It follows from (2.1), (2.2), and (2.33) that

$$\begin{aligned} & \|T(x+y) + T(x+\sigma(y)) - 2T(x) - 2T(y)\| \\ &= \lim_{n \rightarrow \infty} \frac{1}{2^{2n}} \left\| h(2^n x + 2^n y) + (2^n - 1)h\left(2^{n-1}(x+y) + 2^{n-1}(\sigma(x) + \sigma(y))\right) \right. \\ &\quad + h(2^n x + 2^n \sigma(y)) + (2^n - 1)h\left(2^{n-1}(x + \sigma(y)) + 2^{n-1}(\sigma(x) + y)\right) \\ &\quad - 2h(2^n x) + 2(2^n - 1)h\left(2^{n-1}x + 2^{n-1}\sigma(x)\right) \\ &\quad \left. - 2h(2^n y) + 2(2^n - 1)h\left(2^{n-1}y + 2^{n-1}\sigma(y)\right) \right\| \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{2^{2n}} \|h(2^n x + 2^n y) + h(2^n x + 2^n \sigma(y)) - 2h(2^n x) - 2h(2^n y)\| \\ &\quad + \lim_{n \rightarrow \infty} \frac{(2^n - 1)}{2^{2n}} \left\| h\left(2^{n-1}(x + \sigma(x)) + 2^{n-1}(y + \sigma(y))\right) \right. \\ &\quad + h\left(2^{n-1}(x + \sigma(x)) + 2^{n-1}(y + \sigma(y))\right) \\ &\quad \left. - 2h\left(2^{n-1}x + 2^{n-1}\sigma(x)\right) - 2h\left(2^{n-1}y + 2^{n-1}\sigma(y)\right) \right\| \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{2^{2n}} M'(2^n x, 2^n y) + \lim_{n \rightarrow \infty} \frac{(2^n - 1)}{2^{2n}} M'(2^n(x + \sigma(x)), 2^n(y + \sigma(y))) = 0 \end{aligned} \quad (2.34)$$

for all $x, y \in E_1$, which implies that T is a solution of (1.6).

By Theorem 1.1 (c) and (2.29), we obtain

$$d(h, T) \leq \frac{1}{1-L} d(h, Jh) < \frac{1}{4(1-L)}, \quad (2.35)$$

that is, (2.3) is true for all $x \in E_1$. Assume that $T_1 : E_1 \rightarrow E_2$ is another solution of (2.2) satisfying (2.3). We know that T_1 is a fixed point of J . In view of (2.3) and the definition of d , we can conclude that (2.35) is true with T_1 in place of T . Due to Theorem 1.1 (b), we get $T = T_1$. This proves the uniqueness of T .

By (2.16), (2.28), we obtain

$$\begin{aligned}\|f(x) - T(x)\| &\leq \|f(x) - h(x)\| + \|h(x) - T(x)\| \\ &\leq M(x, 0) + \frac{1}{4(1-L)} M'(x, x).\end{aligned}\quad (2.36)$$

Also by (2.12), (2.28), we obtain

$$\begin{aligned}\|k(x) - T(x)\| &\leq \|k(x) - h(x)\| + \|h(x) - T(x)\| \\ &\leq \frac{1}{4}(\varphi(0, x) + \varphi(x, 0)) + \frac{1}{4(1-L)} M'(x, x),\end{aligned}\quad (2.37)$$

and by (2.13), (2.28), we obtain

$$\begin{aligned}\|g(x) - T(x)\| &\leq \|f(x) - g(x)\| + \|f(x) - T(x)\| \\ &\leq \frac{1}{2}(\varphi(x, x) + \varphi(x, -x)) + M(x, 0) + \frac{1}{4(1-L)} M'(x, x).\end{aligned}\quad (2.38)$$

□

In the following, we will investigate some special cases of Theorem 2.1.

Remark 2.2. Let E_1 be a commutative semigroup (with the divisibility by 2), and let E_2 be a real Banach space. Suppose that a function $\varphi : E_1 \times E_1 \rightarrow [0, \infty)$ is given and there exists a constant L , $0 < L < 1$, such that

$$\begin{aligned}\varphi(x, y) &\leq \frac{L}{8} \varphi(2x, 2y), \\ \varphi(x + \sigma(x), y + \sigma(y)) &\leq \frac{L}{4} \varphi(2x, 2y),\end{aligned}\quad (2.39)$$

for all $x, y \in E_1$. Furthermore, let $f, g, h, k : E_1 \rightarrow E_2$ be even functions satisfying the inequality

$$\|f(x + y) + g(x + \sigma(y)) - 2h(x) - 2k(y)\| \leq \varphi(x, y) \quad (2.40)$$

for all $x, y \in E_1$, where $\sigma : E_1 \rightarrow E_1$ is an involution of E_1 and $f(0) = g(0) = h(0) = k(0) = 0$. Then there exists a unique solution $T : E_1 \rightarrow E_2$ of (2.40) such that

$$\begin{aligned}\|2f(x) - T(x)\| &\leq \frac{L}{4(1-L)}M'(x, x) + M(x, 0), \\ \|2g(x) - T(x)\| &\leq \frac{L}{4(1-L)}M'(x, x) + M(x, 0) + \frac{1}{2}(\varphi(x, x) + \varphi(x, -x)), \\ \|h(x) - T(x)\| &\leq \frac{L}{4(1-L)}M'(x, x), \\ \|k(x) - T(x)\| &\leq \frac{L}{4(1-L)}M'(x, x) + \frac{1}{4}(\varphi(0, x) + \varphi(x, 0))\end{aligned}\tag{2.41}$$

for all $x \in E_1$, where

$$\begin{aligned}M(x, y) &= \varphi(x, y) + \varphi(0, y) + \varphi(y, 0) + \varphi\left(\frac{x}{2}, \frac{x}{2}\right) + \varphi\left(\frac{x}{2}, -\left(\frac{x}{2}\right)\right), \\ M'(x, y) &= M(x, y) + M(x + y, 0) + M(x + \sigma(y), 0), \\ T(x) &= \lim_{n \rightarrow \infty} \left(2^{2n} \left[h\left(\frac{x}{2^n}\right) + \left(\frac{1}{2^n} - 1\right) h\left(\frac{x}{2^{n+1}} + \frac{\sigma(x)}{2^{n+1}}\right) \right] \right).\end{aligned}\tag{2.42}$$

Remark 2.3. Let E_1 and E_2 be real Banach spaces. Let the hypotheses of Theorem 2.1 hold. If we put

$$\phi(x, y) = \delta, \quad \delta > 0\tag{2.43}$$

for all $x, y \in E_1$, then, there exists a unique solution $T : E_1 \rightarrow E_2$ such that

$$\begin{aligned}\|2f(x) - T(x)\| &\leq \frac{25}{2}\delta, \\ \|2g(x) - T(x)\| &\leq \frac{27}{2}\delta, \\ \|h(x) - T(x)\| &\leq \frac{15}{2}\delta, \\ \|K(x) - T(x)\| &\leq 8\delta,\end{aligned}\tag{2.44}$$

for all $x \in E_1$, where

$$T(x) = \lim_{n \rightarrow \infty} \frac{1}{2^{2n}} \left[h(2^n x) + (2^n - 1) h\left(2^{n-1}x + 2^{n-1}\sigma(x)\right) \right].\tag{2.45}$$

Also, if we put $\phi(x, y) = \epsilon(\|x\|^p + \|y\|^p)$ for $0 \leq p < 1$ and $\epsilon > 0$, then there exists a unique solution $T : E_1 \rightarrow E_2$ such that

$$\begin{aligned}
 \|2f(x) - T(x)\| &\leq \frac{1}{2(2-2^p)} \epsilon \left(10 + \frac{3+(-1)^p}{2^{p-1}} + 2(-2)^p \right) \epsilon \|x\|^p \\
 &\quad + \left(1 + \frac{3+(-1)^p}{2^p} \right) \epsilon \|x\|^p, \\
 \|2g(x) - T(x)\| &\leq \frac{1}{2(2-2^p)} \epsilon \left(10 + \frac{3+(-1)^p}{2^{p-1}} + 2(-2)^p \right) \epsilon \|x\|^p \\
 &\quad + \left(\frac{5}{2} + \frac{3+(-1)^p}{2^p} + \frac{(-1)^p}{2} \right) \epsilon \|x\|^p, \\
 \|h(x) - T(x)\| &\leq \frac{1}{2(2-2^p)} \epsilon \left(10 + \frac{3+(-1)^p}{2^{p-1}} + 2(-2)^p \right) \epsilon \|x\|^p, \\
 \|k(x) - T(x)\| &\leq \frac{1}{2(2-2^p)} \epsilon \left(10 + \frac{3+(-1)^p}{2^{p-1}} + 2(-2)^p \right) \epsilon \|x\|^p + \frac{1}{2} \epsilon \|x\|^p.
 \end{aligned} \tag{2.46}$$

Similarly, let $\epsilon, p, q \geq 0$ be real numbers such that $p + q < 1$. If we put $\phi(x, y) = \epsilon(\|x\|^p)(\|y\|^q)$ (see [22, 28]), then there exists a unique solution $T : E_1 \rightarrow E_2$ such that

$$\begin{aligned}
 \|2f(x) - T(x)\| &\leq \frac{1}{4-2^{p+q+1}} \left(5 + \frac{1}{2^{p+q-1}} \right) \epsilon (\|x\|^{p+q}) + \frac{\epsilon}{2^{p+q}} (\|x\|^{p+q}), \\
 \|2g(x) - T(x)\| &\leq \frac{1}{4-2^{p+q+1}} \left(5 + \frac{1}{2^{p+q-1}} \right) \epsilon (\|x\|^{p+q}) + \left(\frac{1}{2^{p+q}} + 1 \right) \epsilon (\|x\|^{p+q}), \\
 \|h(x) - T(x)\| &\leq \frac{1}{4-2^{p+q+1}} \left(5 + \frac{1}{2^{p+q-1}} \right) \epsilon (\|x\|^{p+q}), \\
 \|k(x) - T(x)\| &\leq \frac{1}{4-2^{p+q+1}} \left(5 + \frac{1}{2^{p+q-1}} \right) \epsilon (\|x\|^{p+q}).
 \end{aligned} \tag{2.47}$$

Let $\epsilon, p, q \geq 0$ be real numbers such that $p + q < 1$. Another control function is $\epsilon(\|x\|^p \|y\|^q + \|x\|^{p+q} + \|y\|^{p+q})$ (see [35]). Then, there exists a unique solution $T : E_1 \rightarrow E_2$ such that

$$\begin{aligned}
 \|2f(x) - T(x)\| &\leq \frac{1}{4-2^{p+q+1}} \left(15 + \frac{9}{2^{p+q-1}} \right) \|x\|^{p+q} + \left(1 + \frac{3}{2^{p+q-1}} \right) \epsilon \|x\|^{p+q}, \\
 \|2g(x) - T(x)\| &\leq \frac{1}{4-2^{p+q+1}} \left(15 + \frac{9}{2^{p+q-1}} \right) \|x\|^{p+q} + \left(4 + \frac{3}{2^{p+q-1}} \right) \epsilon \|x\|^{p+q}, \\
 \|h(x) - T(x)\| &\leq \frac{1}{4-2^{p+q+1}} \left(15 + \frac{9}{2^{p+q-1}} \right) \|x\|^{p+q}, \\
 \|k(x) - T(x)\| &\leq \frac{1}{4-2^{p+q+1}} \left(15 + \frac{9}{2^{p+q-1}} \right) \|x\|^{p+q} + \frac{1}{2} \epsilon \|x\|^{p+q}
 \end{aligned} \tag{2.48}$$

for all $x \in E_1$.

Theorem 2.4. Let E_1 be a commutative semigroup (with the divisibility by 2) and let E_2 be a real Banach space. Suppose that a function $\varphi : E_1 \times E_1 \rightarrow [0, \infty)$ is given and there exists a constant L , $0 < L < 1$, such that

$$\begin{aligned}\varphi(2x, 2y) &\leq L\varphi(x, y), \\ \varphi(x + \sigma(x), y + \sigma(y)) &\leq L\varphi(x, y),\end{aligned}\tag{2.49}$$

for all $x, y \in E_1$. Furthermore, let $f, g, h, k : E_1 \rightarrow E_2$ be odd functions satisfying the inequality

$$\|f(x + y) + g(x + \sigma(y)) - h(x) - k(y)\| \leq \varphi(x, y),\tag{2.50}$$

for all $x, y \in E_1$, where $\sigma : E_1 \rightarrow E_1$ is an involution of E_1 . Then, there exists a unique solution $T : E_1 \rightarrow E_2$ of (2.50) such that

$$\begin{aligned}\|f(x) - T(x)\| &\leq \frac{1}{4(1-L)}M'(x, x) + \frac{3}{2}(\varphi(x, 0) + \varphi(0, x)), \\ \|g(x) - T(x)\| &\leq \frac{1}{2(1-L)}M'(x, x) + 2\varphi(x, 0) + \varphi(0, x), \\ \|h(x) - T(x)\| &\leq \frac{1}{2(1-L)}M'(x, x), \\ \|k(x) - T(x)\| &\leq \frac{1}{2(1-L)}M'(x, x) + \frac{1}{2}(\varphi(x, 0) + \varphi(0, x)),\end{aligned}\tag{2.51}$$

for all $x \in E_1$, where

$$\begin{aligned}M(x, y) &= \varphi(x, y) + \varphi(x + y, 0) + \varphi(0, x + y) + \frac{3}{2}(\varphi(x, 0) + \varphi(0, x)) \\ &\quad + \varphi(x, -x) + \frac{1}{2}(\varphi(y, 0) + \varphi(0, y)) + \varphi(y, -y), \\ M'(x, y) &= M(x, y) + M(x, \sigma(y)), \\ T(x) &= \lim_{n \rightarrow \infty} \frac{1}{2^n} \left[h(2^n x) + (2^n - 1)h(2^{n-1}x + 2^{n-1}\sigma(x)) \right].\end{aligned}\tag{2.52}$$

Proof. As in the Theorem 2.1, if we put $y = 0$, $x = 0$ (and replace y by x), $y = x$, and $y = -x$ in (2.50) separately, then we obtain

$$\|f(x) + g(x) - 2h(x)\| \leq \varphi(x, 0),\tag{2.53}$$

$$\|f(y) + g(\sigma(y)) - 2k(y)\| \leq \varphi(0, y),\tag{2.54}$$

$$\|f(2x) + g(x + \sigma(x)) - 2h(x) - 2k(x)\| \leq \varphi(x, x),\tag{2.55}$$

$$\|g(x - \sigma(x)) - 2h(x) + 2k(x)\| \leq \varphi(x, -x),\tag{2.56}$$

for all $x \in E_1$, respectively.

We replace y by $x - \sigma(x) := d$ in (2.54). Since $\sigma : E_1 \rightarrow E_1$ is an involution, then

$$\|f(d) - g(d) - 2k(d)\| \leq \varphi(0, d). \quad (2.57)$$

Also, replace y by $x + \sigma(x) := d$ in (2.54), then

$$\|f(d) + g(d) - 2k(d)\| \leq \varphi(0, d). \quad (2.58)$$

Also, we replace x by $x - \sigma(x) := d$ in (2.55), then

$$\|f(2d) - 2h(d) - 2k(d)\| \leq \varphi(d, d). \quad (2.59)$$

We replace x by $x - \sigma(x) := d$ in (2.56), then

$$\|g(2d) - 2h(d) + 2k(d)\| \leq \varphi(d, -d). \quad (2.60)$$

Due to (2.53) and (2.57), we have

$$\|2f(d) - 2h(d) - 2k(d)\| \leq \varphi(d, 0) + \varphi(0, d), \quad (2.61)$$

$$\|2g(d) - 2h(d) + 2k(d)\| \leq \varphi(d, 0) + \varphi(0, d). \quad (2.62)$$

Combining (2.54) with (2.61) yields

$$\|h(2d) + k(2d) - 2h(d) - 2k(d)\| \leq \frac{1}{2}(\varphi(d, 0) + \varphi(0, d)) + \varphi(d, d). \quad (2.63)$$

Due to (2.60) and (2.62), we have

$$\|h(2d) - k(2d) - 2h(d) + 2k(d)\| \leq \frac{1}{2}(\varphi(d, 0) + \varphi(0, d)) + \varphi(d, -d). \quad (2.64)$$

Now, it follows from (2.63) and (2.64) that

$$\begin{aligned} \|h(2d) - 2h(d)\| &\leq \frac{1}{2}(\varphi(d, 0) + \varphi(0, d)) + \varphi(d, -d), \\ \|k(2d) - 2k(d)\| &\leq \frac{1}{2}(\varphi(d, 0) + \varphi(0, d)) + \varphi(d, -d). \end{aligned} \quad (2.65)$$

By (2.50), (2.61), (2.62), and (2.65), we have

$$\begin{aligned}
 & \|h(x+y) + k(x+y) + h(x+\sigma(y)) - k(x+\sigma(y)) - h(2x) - k(2y)\| \\
 & \leq \|f(x+y) + g(x+\sigma(y)) - 2h(x) - 2k(y)\| \\
 & \quad + \|h(x+y) + k(x+y) - f(x+y)\| \\
 & \quad + \|h(x+\sigma(y)) - k(x+\sigma(y)) - g(x+\sigma(y))\| \\
 & \quad + \|h(2x) - 2h(x)\| + \|k(2y) - 2k(y)\| \\
 & \leq \varphi(x, y) + \varphi(x+y, 0) + \varphi(0, x+y) + \frac{3}{2}(\varphi(x, 0) + \varphi(0, x)) \\
 & \quad + \varphi(x, -x) + \frac{1}{2}(\varphi(y, 0) + \varphi(0, y)) + \varphi(y, -y) = M(x, y)
 \end{aligned} \tag{2.66}$$

for all $x, y \in E_1$. If we replace y in (2.66) by $\sigma(y)$, we get

$$\|h(x+\sigma(y)) + k(x+\sigma(y)) + h(x+y) - k(x+y) - h(2x) - k(2\sigma(y))\| \leq M(x, \sigma(y)). \tag{2.67}$$

By (2.66) and (2.67), we get

$$\|2h(x+y) + 2h(x+\sigma(y)) - h(2x) - h(2\sigma(y))\| \leq M(x, y) + M(x, \sigma(y)). \tag{2.68}$$

If we replace y in (2.66) by $\sigma(y)$ and combine (2.53) with (2.58), we get

$$\|2h(x+y) + 2h(x+\sigma(y)) - h(2x) - h(2y)\| \leq M(x, y) + M(x, \sigma(y)). \tag{2.69}$$

By (2.65) and (2.69), we get

$$\begin{aligned}
 \|2h(x+y) + 2h(x+\sigma(y)) - 2h(x) - 2h(y)\| & \leq M(x, y) + M(x, \sigma(y)) \\
 & = M'(x, y).
 \end{aligned} \tag{2.70}$$

Therefore

$$\|h(x+y) + h(x+\sigma(y)) - h(x) - h(y)\| \leq M'(x, y). \tag{2.71}$$

We define X to be the set of all functions $f : E_1 \rightarrow E_2$ and introduce a generalized metric on X as follows:

$$d(g, h) = \inf\{C \in [0, \infty) : \|g(x) - h(x)\| \leq CM'(x, x), \forall x \in E_1\}. \tag{2.72}$$

Also, we define an operator $J : X \rightarrow X$ by

$$JL(x) = \frac{1}{2}[L(2x) + L(x + \sigma(x))] \quad (2.73)$$

for all $x \in E_1$.

Then, there exists a function $T : E_1 \rightarrow E_2$ which is a fixed point of J , such that $d(J^n h, T) \rightarrow 0$ as $n \rightarrow \infty$. By induction, we can show that

$$(J^n h)(x) = \frac{1}{2^n} \left[h(2^n x) + (2^n - 1)h(2^{n-1}x + 2^{n-1}\sigma(x)) \right] \quad (2.74)$$

for each $n \in \mathbb{N}$. Since $d(J^n h, T) \rightarrow 0$ as $n \rightarrow \infty$, there exists a sequence $\{C_n\}$ such that $C_n \rightarrow 0$ as $n \rightarrow \infty$ and $d(J^n h, T) \leq C_n$ for every $n \in \mathbb{N}$. Hence,

$$\|J^n h(x) - T(x)\| \leq C_n M'(x, x) \quad (2.75)$$

for all $x \in E_1$. Thus, for each fixed $x \in E_1$, we have

$$\lim_{n \rightarrow \infty} \|J^n h(x) - T(x)\| = 0. \quad (2.76)$$

Then

$$T(x) = \lim_{n \rightarrow \infty} \frac{1}{2^n} \left[h(2^n x) + (2^n - 1)h(2^{n-1}x + 2^{n-1}\sigma(x)) \right] \quad (2.77)$$

for all $x \in E_1$. Hence, T is a solution of (2.51). $\square$

Remark 2.5. Let E_1 and E_2 be real Banach spaces. Let the hypotheses of Theorem 2.4 hold. If we replace the control function $\phi(x, y)$ by $\delta > 0$, then, there exists a unique solution $T : E_1 \rightarrow E_2$ such that

$$\begin{aligned} \|f(x) - T(x)\| &\leq 12\delta, \\ \|g(x) - T(x)\| &\leq 21\delta, \\ \|h(x) - T(x)\| &\leq 18\delta, \\ \|k(x) - T(x)\| &\leq 19\delta, \end{aligned} \quad (2.78)$$

for all $x \in E_1$, where

$$T(x) = \lim_{n \rightarrow \infty} \frac{1}{2^n} \left[h(2^n x) + (2^n - 1)h(2^{n-1}x + 2^{n-1}\sigma(x)) \right]. \quad (2.79)$$

If we put $\phi(x, y) = \epsilon(\|x\|^p + \|y\|^p)$ in which $\epsilon > 0$ and p is a nonnegative number less than 1, then, there exists a unique solution $T : E_1 \rightarrow E_2$ such that

$$\begin{aligned}\|f(x) - T(x)\| &\leq \frac{2^{p-1}}{(2^p - 1)} (12 + 2(-1)^p) \epsilon \|x\|^p + 3(\epsilon \|x\|^p), \\ \|g(x) - T(x)\| &\leq \frac{2^p}{(2^p - 1)} (12 + 2(-1)^p) \epsilon \|x\|^p + 3(\epsilon \|x\|^p), \\ \|h(x) - T(x)\| &\leq \frac{2^p}{(2^p - 1)} (12 + 2(-1)^p) \epsilon \|x\|^p, \\ \|k(x) - T(x)\| &\leq \frac{2^p}{(2^p - 1)} (12 + 2(-1)^p) \epsilon \|x\|^p + (\epsilon \|x\|^p).\end{aligned}\tag{2.80}$$

Let $\epsilon > 0$ and $p, q > 2$. If we put $\phi(x, y) = \epsilon(\|x\|^p)(\|y\|^q)$, then, there exists a unique solution $T : E_1 \rightarrow E_2$ such that

$$\begin{aligned}\|f(x) - T(x)\| &\leq \frac{3}{2(2^{(p+q+2)} - 1)} \epsilon \|x\|^{p+q}, \\ \|g(x) - T(x)\| &\leq \frac{3}{(2^{(p+q+2)} - 1)} \epsilon \|x\|^{p+q}, \\ \|h(x) - T(x)\| &\leq \frac{3}{(2^{(p+q+2)} - 1)} \epsilon \|x\|^{p+q}, \\ \|k(x) - T(x)\| &\leq \frac{3}{(2^{(p+q+2)} - 1)} \epsilon \|x\|^{p+q}.\end{aligned}\tag{2.81}$$

Finally, if we put $\phi(x, y) = \epsilon(\|x\|^p \|y\|^q + \|x\|^{p+q} + \|y\|^{p+q})$, then, there exists a unique solution $T : E_1 \rightarrow E_2$ such that

$$\begin{aligned}\|f(x) - T(x)\| &\leq \frac{9}{2(2^{(p+q+2)} - 1)} \epsilon \|x\|^{p+q}, \\ \|g(x) - T(x)\| &\leq \frac{9}{(2^{(p+q+2)} - 1)} \epsilon \|x\|^{p+q}, \\ \|h(x) - T(x)\| &\leq \frac{9}{(2^{(p+q+2)} - 1)} \epsilon \|x\|^{p+q}, \\ \|k(x) - T(x)\| &\leq \frac{9}{(2^{(p+q+2)} - 1)} \epsilon \|x\|^{p+q}\end{aligned}\tag{2.82}$$

for all $x \in E_1$.

Remark 2.6. The methods of proofs, used in this paper, can be also applied to the problem of stability of (1.6) on the restricted domain, analogously as in the papers [18, 46]. Also, this corresponds to the results in [38, 41–43], where the Pexiderized equations' stability on restricted domains has been investigated.

Acknowledgments

The authors would like to thank the referee and the associate editor Professor S. Reich for giving useful suggestions and comments for the improvement of this paper.

References

- [1] S. M. Ulam, *Problems in Modern Mathematics*, chapter 6, John Wiley & Sons, New York, NY, USA, 1964.
- [2] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [3] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.
- [4] Th. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [5] S. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific, River Edge, NJ, USA, 2002.
- [6] K.-W. Jun, D.-S. Shin, and B.-D. Kim, "On Hyers-Ulam-Rassias stability of the Pexider equation," *Journal of Mathematical Analysis and Applications*, vol. 239, no. 1, pp. 20–29, 1999.
- [7] Th. M. Rassias, Ed., *Functional Equations, Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2003.
- [8] L. Maligranda, "A result of Tosio Aoki about a generalization of Hyers-Ulam stability of additive functions—a question of priority," *Aequationes Mathematicae*, vol. 75, no. 3, pp. 289–296, 2008.
- [9] Z. Moszner, "On the stability of functional equations," *Aequationes Mathematicae*, vol. 77, no. 1-2, pp. 33–88, 2009.
- [10] J. Aczél, *A Short Course on Functional Equations*, Theory and Decision Library. Series B: Mathematical and Statistical Methods, D. Reidel, Dordrecht, The Netherlands, 1987.
- [11] G. L. Forti, "An existence and stability theorem for a class of functional equations," *Stochastica*, vol. 4, no. 1, pp. 23–30, 1980.
- [12] P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431–436, 1994.
- [13] M. Mirzavaziri and M. S. Moslehian, "A fixed point approach to stability of a quadratic equation," *Bulletin of the Brazilian Mathematical Society*, vol. 37, no. 3, pp. 361–376, 2006.
- [14] J. Rätz, "On orthogonally additive mappings," *Aequationes Mathematicae*, vol. 28, no. 1-2, pp. 35–49, 1985.
- [15] F. Skof, "Local properties and approximation of operators," *Rendiconti del Seminario Matematico e Fisico di Milano*, vol. 53, pp. 113–129, 1983.
- [16] P. W. Cholewa, "Remarks on the stability of functional equations," *Aequationes Mathematicae*, vol. 27, no. 1-2, pp. 76–86, 1984.
- [17] S. Czerwik, "On the stability of the quadratic mapping in normed spaces," *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, vol. 62, pp. 59–64, 1992.
- [18] J. Brzdęk, "On a method of proving the Hyers-Ulam stability of functional equations on restricted domains," *The Australian Journal of Mathematical Analysis and Applications*, vol. 6, no. 1, article 4, pp. 1–10, 2009.
- [19] S.-M. Jung, "Stability of the quadratic equation of Pexider type," *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, vol. 70, pp. 175–190, 2000.
- [20] S.-M. Jung and P. K. Sahoo, "Hyers-Ulam stability of the quadratic equation of Pexider type," *Journal of the Korean Mathematical Society*, vol. 38, no. 3, pp. 645–656, 2001.
- [21] B. Bouikhalene, E. Elhoucien, and Th. M. Rassias, "On the generalized Hyers-Ulam stability of the quadratic functional equation with a general involution," *Nonlinear Functional Analysis and Applications*, vol. 12, no. 2, pp. 247–262, 2007.
- [22] B. Bouikhalene and E. Elqorachi, "Ulam-Găvruta-Rassias stability of the Pexider functional equation," *International Journal of Applied Mathematics & Statistics*, vol. 7, no. Fe07, pp. 27–39, 2007.
- [23] H. Cao, J.-R. Lv, and J. M. Rassias, "Superstability for generalized module left derivations and generalized module derivations on a Banach module. I," *Journal of Inequalities and Applications*, vol. 2009, Article ID 718020, 10 pages, 2009.

- [24] P. W. Cholewa, "The stability of the sine equation," *Proceedings of the American Mathematical Society*, vol. 88, no. 4, pp. 631–634, 1983.
- [25] M. Eshaghi Gordji and H. Khodaei, "On the generalized Hyers-Ulam-Rassias stability of quadratic functional equations," *Abstract and Applied Analysis*, vol. 2009, Article ID 923476, 11 pages, 2009.
- [26] S.-M. Jung and Z.-H. Lee, "A fixed point approach to the stability of quadratic functional equation with involution," *Fixed Point Theory and Applications*, vol. 2008, Article ID 732086, 11 pages, 2008.
- [27] P. Nakmahachalasint, "On the generalized Ulam-Gavruta-Rassias stability of mixed-type linear and Euler-Lagrange-Rassias functional equations," *International Journal of Mathematics and Mathematical Sciences*, vol. 2007, Article ID 63239, 10 pages, 2007.
- [28] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Journal of Functional Analysis*, vol. 46, no. 1, pp. 126–130, 1982.
- [29] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Bulletin des Sciences Mathématiques. 2e Série*, vol. 108, no. 4, pp. 445–446, 1984.
- [30] J. M. Rassias, "On a new approximation of approximately linear mappings by linear mappings," *Discussiones Mathematicae*, vol. 7, pp. 193–196, 1985.
- [31] J. M. Rassias, "Solution of a problem of Ulam," *Journal of Approximation Theory*, vol. 57, no. 3, pp. 268–273, 1989.
- [32] J. M. Rassias, "Solution of a stability problem of Ulam," *Discussiones Mathematicae*, vol. 12, pp. 95–103, 1992.
- [33] J. M. Rassias, "Complete solution of the multi-dimensional problem of Ulam," *Discussiones Mathematicae*, vol. 14, pp. 101–107, 1994.
- [34] J. M. Rassias, "On the stability of a multi-dimensional Cauchy type functional equation," in *Geometry, Analysis and Mechanics*, pp. 365–376, World Scientific, River Edge, NJ, USA, 1994.
- [35] K. Ravi, M. Arunkumar, and J. M. Rassias, "Ulam stability for the orthogonally general Euler-Lagrange type functional equation," *International Journal of Mathematics and Statistics*, vol. 3, no. A08, pp. 36–46, 2008.
- [36] M. B. Savadkouhi, M. E. Gordji, J. M. Rassias, and N. Ghobadipour, "Approximate ternary Jordan derivations on Banach ternary algebras," *Journal of Mathematical Physics*, vol. 50, no. 4, Article ID 042303, 9 pages, 2009.
- [37] H. Stetkaer, "Functional equations on abelian groups with involution," *Aequationes Mathematicae*, vol. 54, no. 1-2, pp. 144–172, 1997.
- [38] J. Brzdęk and J. Sikorska, "A conditional exponential functional equation and its stability," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 6, pp. 2923–2934, 2010.
- [39] K. Ciepliński, "On a generalized Pexider equation and the Pexider difference," in *Iteration Theory (ECIT '06)*, vol. 351 of *Grazer Mathematische Berichte*, pp. 27–36, Institut für Mathematik, Karl-Franzens-Universität Graz, Graz, Austria, 2007.
- [40] J. B. Diaz and B. Margolis, "A fixed point theorem of the alternative, for contractions on a generalized complete metric space," *Bulletin of the American Mathematical Society*, vol. 74, no. 2, pp. 305–309, 1968.
- [41] J. Sikorska, "On two conditional Pexider functional equations and their stabilities," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 7, pp. 2673–2684, 2009.
- [42] J. Sikorska, "On a pexiderized conditional exponential functional equation," *Acta Mathematica Hungarica*, vol. 125, no. 3, pp. 287–299, 2009.
- [43] J. Sikorska, "Exponential functional equation on spheres," *Applied Mathematics Letters*, vol. 23, no. 2, pp. 156–160, 2010.
- [44] D. H. Hyers, G. Isac, and Th. M. Rassias, *Topics in Nonlinear Analysis & Applications*, World Scientific, River Edge, NJ, USA, 1997.
- [45] L. Cădariu and V. Radu, "Fixed points and the stability of Jensen's functional equation," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 1, article 4, 7 pages, 2003.
- [46] J. Brzdęk and A. Pietrzyk, "A note on stability of the general linear equation," *Aequationes Mathematicae*, vol. 75, no. 3, pp. 267–270, 2008.