

Research Article

Second Moment Convergence Rates for Uniform Empirical Processes

You-You Chen and Li-Xin Zhang

Department of Mathematics, Zhejiang University, Hangzhou 310027, China

Correspondence should be addressed to You-You Chen, cyyooo@gmail.com

Received 21 May 2010; Revised 3 August 2010; Accepted 19 August 2010

Academic Editor: Andrei Volodin

Copyright © 2010 Y.-Y. Chen and L.-X. Zhang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let $\{U_1, U_2, \dots, U_n\}$ be a sequence of independent and identically distributed $U[0, 1]$ -distributed random variables. Define the uniform empirical process as $\alpha_n(t) = n^{-1/2} \sum_{i=1}^n (I\{U_i \leq t\} - t)$, $0 \leq t \leq 1$, $\|\alpha_n\| = \sup_{0 \leq t \leq 1} |\alpha_n(t)|$. In this paper, we get the exact convergence rates of weighted infinite series of $E\|\alpha_n\|^2 I\{\|\alpha_n\| \geq \varepsilon(\log n)^{1/\beta}\}$.

1. Introduction and Main Results

Let $\{X, X_n; n \geq 1\}$ be a sequence of independent and identically distributed (i.i.d.) random variables with zero mean. Set $S_n = \sum_{i=1}^n X_i$ for $n \geq 1$, and $\log x = \ln(x \vee e)$. Hsu and Robbins [1] introduced the concept of complete convergence. They showed that

$$\sum_{n=1}^{\infty} P\{|S_n| \geq \varepsilon n\} < \infty, \quad \varepsilon > 0 \quad (1.1)$$

if $EX = 0$ and $EX^2 < \infty$. The converse part was proved by the study of Erdős in [2]. Obviously, the sum in (1.1) tends to infinity as $\varepsilon \searrow 0$. Many authors studied the exact rates in terms of ε (cf. [3–5]). Chow [6] studied the complete convergence of $E\{|S_n - \varepsilon n^\alpha\}_+$, $\varepsilon > 0$. Recently, Liu and Lin [7] introduced a new kind of complete moment convergence which is interesting, and got the precise rate of it as follows.

Theorem A. Suppose that $\{X, X_n; n \geq 1\}$ is a sequence of i.i.d. random variables, then

$$\lim_{\varepsilon \searrow 0} \frac{1}{-\log \varepsilon} \sum_{n=1}^{\infty} \frac{1}{n^2} \text{ES}_n^2 I\{|S_n| \geq \varepsilon n\} = 2\sigma^2 \quad (1.2)$$

holds, if and only if $\text{EX} = 0$, $\text{EX}^2 = \sigma^2$, and $\text{EX}^2 \log^+ |X| < \infty$.

Other than partial sums, many authors investigated precise rates in some different cases, such as U -statistics (cf. [8, 9]) and self-normalized sums (cf. [10, 11]). Zhang and Yang [12] extended the precise asymptotic results to the uniform empirical process. We suppose U_1, U_2, \dots, U_n is the sample of $U[0, 1]$ random variables and $E_n(t)$ is the empirical distribution function of it. Denote the uniform empirical process by $\alpha_n(t) = \sqrt{n}(E_n(t) - t)$, $0 \leq t \leq 1$, and the norm of a function $f(t)$ on $[0, 1]$ by $\|f\| = \sup_{0 \leq t \leq 1} |f(t)|$. Let $B(t)$, $t \in [0, 1]$ be the Brownian bridge. We present one result of Zhang and Yang [12] as follows.

Theorem B. For any $\delta > -1$, one has

$$\lim_{\varepsilon \searrow 0} \varepsilon^{2\delta+2} \sum_{n=1}^{\infty} \frac{(\log n)^\delta}{n} P\left\{\|\alpha_n\| \geq \varepsilon \sqrt{\log n}\right\} = \frac{\text{E}\|B\|^{2\delta+2}}{\delta+1}. \quad (1.3)$$

Inspired by the above conclusions, we consider second moment convergence rates for the uniform empirical process in the law of iterated logarithm and the law of the logarithm. Throughout this paper, let C denote a positive constant whose values can be different from one place to another. $[x]$ will denote the largest integer $\leq x$. The following two theorems are our main results.

Theorem 1.1. For $0 < \beta \leq 2$, $\delta > 2/\beta - 1$, one has

$$\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta-2/\beta}}{n} \text{E}\|\alpha_n\|^2 I\left\{\|\alpha_n\| \geq \varepsilon (\log n)^{1/\beta}\right\} = \frac{\beta \text{E}\|B\|^{\beta(\delta+1)}}{\beta(\delta+1)-2}. \quad (1.4)$$

Theorem 1.2. For $0 < \beta \leq 2$, $\delta > 2/\beta - 1$, one has

$$\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta-2/\beta}}{n \log n} \text{E}\|\alpha_n\|^2 I\left\{\|\alpha_n\| \geq \varepsilon (\log \log n)^{1/\beta}\right\} = \frac{\beta \text{E}\|B\|^{\beta(\delta+1)}}{\beta(\delta+1)-2}. \quad (1.5)$$

Remark 1.3. It is well known that $P\{\|B\| \geq x\} = 2 \sum_{k=1}^{\infty} (-1)^{k+1} e^{-2k^2 x^2}$, $x > 0$ (see Csörgő and Révész [13, page 43]). Therefore, by Fubini's theorem we have

$$\begin{aligned} E\|B\|^{\beta(\delta+1)} &= \beta(\delta+1) \int_0^{\infty} x^{\beta(\delta+1)-1} P\{\|B\| \geq x\} dx \\ &= 2\beta(\delta+1) \int_0^{\infty} x^{\beta(\delta+1)-1} \sum_{k=1}^{\infty} (-1)^{k+1} e^{-2k^2 x^2} dx \\ &= \frac{\beta(\delta+1)\Gamma(\beta(\delta+1)/2)}{2^{\beta(\delta+1)/2}} \sum_{k=1}^{\infty} (-1)^{k+1} k^{-\beta(\delta+1)}. \end{aligned} \quad (1.6)$$

Consequently, explicit results of (1.4) and (1.5) can be calculated further.

2. The Proofs

In order to prove Theorem 1.1, we present several propositions first.

Proposition 2.1. For $\beta > 0$, $\delta > -1$, one has

$$\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta}}{n} P\{\|B\| \geq \varepsilon(\log n)^{1/\beta}\} = \frac{E\|B\|^{\beta(\delta+1)}}{\delta+1}. \quad (2.1)$$

Proof. We calculate that

$$\begin{aligned} &\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta}}{n} P\{\|B\| \geq \varepsilon(\log n)^{1/\beta}\} \\ &= \lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)} \int_2^{\infty} \frac{(\log y)^{\delta}}{y} P\{\|B\| \geq \varepsilon(\log y)^{1/\beta}\} dy \\ &= \beta \int_0^{\infty} t^{\beta(\delta+1)-1} P\{\|B\| \geq t\} dt \\ &= \frac{E\|B\|^{\beta(\delta+1)}}{\delta+1}. \end{aligned} \quad (2.2)$$

□

Proposition 2.2. For $\beta > 0$, $\delta > -1$, one has

$$\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta}}{n} \left| P\{\|\alpha_n\| \geq \varepsilon(\log n)^{1/\beta}\} - P\{\|B\| \geq \varepsilon(\log n)^{1/\beta}\} \right| = 0. \quad (2.3)$$

Proof. Following [4], set $A(\varepsilon) = [\exp(M/\varepsilon^\beta)]$, where $M > 1$. Write

$$\begin{aligned} & \sum_{n=2}^{\infty} \frac{(\log n)^\delta}{n} \left| P\{\|\alpha_n\| \geq \varepsilon(\log n)^{1/\beta}\} - P\{\|B\| \geq \varepsilon(\log n)^{1/\beta}\} \right| \\ &= \sum_{n \leq A(\varepsilon)} \frac{(\log n)^\delta}{n} \left| P\{\|\alpha_n\| \geq \varepsilon(\log n)^{1/\beta}\} - P\{\|B\| \geq \varepsilon(\log n)^{1/\beta}\} \right| \\ & \quad + \sum_{n > A(\varepsilon)} \frac{(\log n)^\delta}{n} \left| P\{\|\alpha_n\| \geq \varepsilon(\log n)^{1/\beta}\} - P\{\|B\| \geq \varepsilon(\log n)^{1/\beta}\} \right| \\ &=: I_1 + I_2. \end{aligned} \tag{2.4}$$

It is wellknown that $\alpha_n(\cdot) \xrightarrow{d} B(\cdot)$ (see Csörgő and Révész [13, page 17]). By continuous mapping theorem, we have $\|\alpha_n\| \xrightarrow{d} \|B\|$. As a result, it follows that

$$\Delta_n := \sup_x |P\{\|\alpha_n\| \geq x\} - P\{\|B\| \geq x\}| \rightarrow 0, \quad \text{as } n \rightarrow \infty. \tag{2.5}$$

Using the Toeplitz's lemma (see Stout [14, pages 120-121]), we can get $\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)} I_1 = 0$. For I_2 , it is obvious that

$$\begin{aligned} I_2 &\leq \sum_{n > A(\varepsilon)} \frac{(\log n)^\delta}{n} P\{\|B\| \geq \varepsilon(\log n)^{1/\beta}\} + \sum_{n > A(\varepsilon)} \frac{(\log n)^\delta}{n} P\{\|\alpha_n\| \geq \varepsilon(\log n)^{1/\beta}\} \\ &=: I_3 + I_4. \end{aligned} \tag{2.6}$$

Notice that $A(\varepsilon) - 1 \geq \sqrt{A(\varepsilon)}$, for a small ε . Via the similar argument in [4] we have

$$\begin{aligned} \varepsilon^{\beta(\delta+1)} I_3 &\leq \varepsilon^{\beta(\delta+1)} \sum_{n > A(\varepsilon)} \frac{(\log n)^\delta}{n} P\{\|B\| \geq \varepsilon(\log n)^{1/\beta}\} \\ &\leq C \int_{(M/2)^{1/\beta}}^{\infty} y^{\beta(\delta+1)-1} P\{\|B\| \geq y\} dy \rightarrow 0, \quad \text{as } M \rightarrow \infty. \end{aligned} \tag{2.7}$$

From Kiefer and Wolfowitz [15], we have

$$P\{\|\alpha_n\| \geq x\} \leq Ce^{-Cx^2}. \tag{2.8}$$

Therefore,

$$\begin{aligned}
 \varepsilon^{\beta(\delta+1)} I_4 &\leq C\varepsilon^{\beta(\delta+1)} \sum_{n>A(\varepsilon)} \frac{(\log n)^\delta}{n} \exp\{-C\varepsilon^2(\log n)^{2/\beta}\} \\
 &\leq C\varepsilon^{\beta(\delta+1)} \int_{\sqrt{A(\varepsilon)}}^\infty \frac{(\log x)^\delta}{x} \exp\{-C\varepsilon^2(\log x)^{2/\beta}\} dx \\
 &\leq C \int_{C(M/2)^{2/\beta}}^\infty y^{\beta(\delta+1)/2-1} e^{-y} dy \rightarrow 0, \quad \text{as } M \rightarrow \infty.
 \end{aligned}
 \tag{2.9}$$

From (2.6), (2.7), and (2.9), we get $\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)} I_2 = 0$. Proposition 2.2 has been proved. \square

Proposition 2.3. For $\beta > 0$, $\delta > 2/\beta - 1$, one has

$$\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} \sum_{n=2}^\infty \frac{(\log n)^{\delta-2/\beta}}{n} \int_{\varepsilon(\log n)^{1/\beta}}^\infty 2yP\{\|B\| \geq y\} dy = \frac{2E\|B\|^{\beta(\delta+1)}}{(\delta+1)(\beta(\delta+1)-2)}. \tag{2.10}$$

Proof. The calculation here is analogous to (2.1), so it is omitted here. \square

Proposition 2.4. For $0 < \beta \leq 2$, $\delta > 2/\beta - 1$, one has

$$\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} \sum_{n=2}^\infty \frac{(\log n)^{\delta-2/\beta}}{n} \left| \int_{\varepsilon(\log n)^{1/\beta}}^\infty 2yP\{\|\alpha_n\| \geq y\} dy - \int_{\varepsilon(\log n)^{1/\beta}}^\infty 2yP\{\|B\| \geq y\} dy \right| = 0. \tag{2.11}$$

Proof. Like [4] and Proposition 2.2, we divide the summation into two parts,

$$\begin{aligned}
 &\sum_{n=2}^\infty \frac{(\log n)^{\delta-2/\beta}}{n} \left| \int_{\varepsilon(\log n)^{1/\beta}}^\infty 2yP\{\|\alpha_n\| \geq y\} dy - \int_{\varepsilon(\log n)^{1/\beta}}^\infty 2yP\{\|B\| \geq y\} dy \right| \\
 &= \sum_{n \leq A(\varepsilon)} \frac{(\log n)^{\delta-2/\beta}}{n} \left| \int_{\varepsilon(\log n)^{1/\beta}}^\infty 2yP\{\|\alpha_n\| \geq y\} dy - \int_{\varepsilon(\log n)^{1/\beta}}^\infty 2yP\{\|B\| \geq y\} dy \right| \\
 &\quad + \sum_{n > A(\varepsilon)} \frac{(\log n)^{\delta-2/\beta}}{n} \left| \int_{\varepsilon(\log n)^{1/\beta}}^\infty 2yP\{\|\alpha_n\| \geq y\} dy - \int_{\varepsilon(\log n)^{1/\beta}}^\infty 2yP\{\|B\| \geq y\} dy \right| \\
 &=: J_1 + J_2.
 \end{aligned}
 \tag{2.12}$$

First, consider J_1 ,

$$\begin{aligned}
J_1 &\leq \sum_{n \leq A(\varepsilon)} \frac{(\log n)^{\delta-2/\beta}}{n} \int_{\varepsilon(\log n)^{1/\beta}}^{\infty} 2y |P\{\|\alpha_n\| \geq y\} - P\{\|B\| \geq y\}| dy \\
&\leq \sum_{n \leq A(\varepsilon)} \frac{(\log n)^\delta}{n} \int_0^\infty 2(x+\varepsilon) \left| P\{\|\alpha_n\| \geq (x+\varepsilon)(\log n)^{1/\beta}\} - P\{\|B\| \geq (x+\varepsilon)(\log n)^{1/\beta}\} \right| dx \\
&\leq \sum_{n \leq A(\varepsilon)} \frac{(\log n)^\delta}{n} \left(\int_0^{(\log n)^{-1/\beta} \Delta_n^{-1/4}} 2(x+\varepsilon) \left| P\{\|\alpha_n\| \geq (x+\varepsilon)(\log n)^{1/\beta}\} \right. \right. \\
&\quad \left. \left. - P\{\|B\| \geq (x+\varepsilon)(\log n)^{1/\beta}\} \right| dx \right. \\
&\quad \left. + \int_{(\log n)^{-1/\beta} \Delta_n^{-1/4}}^\infty 2(x+\varepsilon) P\{\|B\| \geq (x+\varepsilon)(\log n)^{1/\beta}\} dx \right. \\
&\quad \left. + \int_{(\log n)^{-1/\beta} \Delta_n^{-1/4}}^\infty 2(x+\varepsilon) P\{\|\alpha_n\| \geq (x+\varepsilon)(\log n)^{1/\beta}\} dx \right) \\
&=: \sum_{n \leq A(\varepsilon)} \frac{(\log n)^\delta}{n} (J_{11} + J_{12} + J_{13}).
\end{aligned} \tag{2.13}$$

Since $n \leq A(\varepsilon)$ means $\varepsilon < (M/\log n)^{1/\beta}$, it follows

$$\begin{aligned}
(\log n)^{2/\beta} J_{11} &\leq (\log n)^{2/\beta} \int_0^{(\log n)^{-1/\beta} \Delta_n^{-1/4}} 2(x+\varepsilon) \Delta_n dx \\
&\leq (\log n)^{2/\beta} \Delta_n \left((\log n)^{-1/\beta} \Delta_n^{-1/4} + (\log n)^{-1/\beta} M^{1/\beta} \right)^2 \\
&\leq \left(\Delta_n^{1/4} + M^{1/\beta} \Delta_n^{1/2} \right)^2 \rightarrow 0, \text{ as } n \rightarrow \infty.
\end{aligned} \tag{2.14}$$

By Lemma 2.1 in Zhang and Yang [12], we have $P\{\|B\| \geq x\} \leq 2e^{-2x^2}$. For J_{12} , it is easy to get

$$\begin{aligned}
(\log n)^{2/\beta} J_{12} &\leq (\log n)^{2/\beta} \int_{\varepsilon(\log n)^{1/\beta} + \Delta_n^{-1/4}}^\infty (\log n)^{-2/\beta} \cdot 2y P\{\|B\| \geq y\} dy \\
&\leq C \int_{\Delta_n^{-1/4}}^\infty 2y \exp\{-2y^2\} dy \\
&\leq C \exp\{-2\Delta_n^{-1/2}\} \rightarrow 0, \text{ as } n \rightarrow \infty.
\end{aligned} \tag{2.15}$$

In the same way, by the inequality $P\{\|\alpha_n\| \geq x\} \leq Ce^{-Cx^2}$, we can get

$$(\log n)^{2/\beta} J_{13} \leq C \exp\{-C\Delta_n^{-1/2}\} \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (2.16)$$

Put the three parts together, we get that $(\log n)^{2/\beta}(J_{11} + J_{12} + J_{13}) \rightarrow 0$ uniformly in ε as $n \rightarrow \infty$. Using Toeplitz's lemma again, we have $\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} J_1 = 0$.

In the sequel, we verify $\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} J_2 = 0$. It is easy to see that

$$\begin{aligned} J_2 &\leq \sum_{n>A(\varepsilon)} \frac{(\log n)^{\delta-2/\beta}}{n} \int_{\varepsilon(\log n)^{1/\beta}}^{\infty} 2xP\{\|B\| \geq x\} dx \\ &\quad + \sum_{n>A(\varepsilon)} \frac{(\log n)^{\delta-2/\beta}}{n} \int_{\varepsilon(\log n)^{1/\beta}}^{\infty} 2xP\{\|\alpha_n\| \geq x\} dx \\ &=: J_{21} + J_{22}. \end{aligned} \quad (2.17)$$

We estimate J_{22} first, by noticing $0 < \beta \leq 2$ and (2.8), it follows

$$\begin{aligned} J_{22} &\leq \sum_{n>A(\varepsilon)} \frac{(\log n)^{\delta-2/\beta}}{n} \int_n^{\infty} 2\varepsilon(\log y)^{1/\beta} P\{\|\alpha_n\| \geq \varepsilon(\log y)^{1/\beta}\} \frac{\varepsilon}{\beta y} (\log y)^{1/\beta-1} dy \\ &\leq C \int_{A(\varepsilon)}^{\infty} \frac{(\log x)^{\delta-2/\beta}}{x} \int_x^{\infty} \frac{\varepsilon^2 (\log y)^{2/\beta-1}}{y} \exp\{-C\varepsilon^2 (\log y)^{2/\beta}\} dy dx \\ &\leq C \int_{A(\varepsilon)}^{\infty} \frac{\varepsilon^2 (\log y)^{2/\beta-1}}{y} \exp\{-C\varepsilon^2 (\log y)^{2/\beta}\} (\log y)^{\delta-2/\beta+1} dy \\ &\leq C\varepsilon^2 \int_{A(\varepsilon)}^{\infty} \frac{(\log y)^{\delta}}{y} \exp\{-C\varepsilon^2 \log y\} dy \\ &\leq C\varepsilon^2 \frac{\log^{\delta}(A(\varepsilon))}{(A(\varepsilon))^{C\varepsilon^2}} \leq C\varepsilon^{2-\beta\delta}. \end{aligned} \quad (2.18)$$

Therefore, we get $\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} J_{22} = 0$. So far, we only need to prove $\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} J_{21} = 0$. Use the inequality $P\{\|B\| \geq x\} \leq 2e^{-2x^2}$ again and follow the proof of J_{22} , we can get this result. The proof of the proposition is completed now. \square

Proof of Theorem 1.1. According to Fubini's theorem, it is easy to get

$$EXI\{X \geq a\} = aP\{X \geq a\} + \int_a^{\infty} P\{X \geq x\} dx, \quad (2.19)$$

for $a > 0$. Therefore, we have

$$\begin{aligned} \mathbb{E}\|\alpha_n\|^2 I\{\|\alpha_n\| \geq \varepsilon(\log n)^{1/\beta}\} &= \varepsilon^2 (\log n)^{2/\beta} P\{\|\alpha_n\| \geq \varepsilon(\log n)^{1/\beta}\} \\ &+ \int_{\varepsilon(\log n)^{1/\beta}}^{\infty} 2yP\{\|\alpha_n\| \geq y\} dy. \end{aligned} \quad (2.20)$$

From Proposition 2.1–2.4, we have

$$\begin{aligned} &\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta-2/\beta}}{n} \mathbb{E}\|\alpha_n\|^2 I\{\|\alpha_n\| \geq \varepsilon(\log n)^{1/\beta}\} \\ &= \lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta}}{n} P\{\|\alpha_n\| \geq \varepsilon(\log n)^{1/\beta}\} \\ &+ \lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} \sum_{n=2}^{\infty} \frac{(\log n)^{\delta-2/\beta}}{n} \int_{\varepsilon(\log n)^{1/\beta}}^{\infty} 2yP\{\|\alpha_n\| \geq y\} dy \\ &= \frac{\beta \mathbb{E}\|B\|^{\beta(\delta+1)}}{\beta(\delta+1)-2}. \end{aligned} \quad (2.21)$$

a □

Proof of Theorem 1.2. From (2.19), we have

$$\begin{aligned} &\varepsilon^{\beta(\delta+1)-2} \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta-2/\beta}}{n \log n} \mathbb{E}\|\alpha_n\|^2 I\{\|\alpha_n\| \geq \varepsilon(\log \log n)^{1/\beta}\} \\ &= \varepsilon^{\beta(\delta+1)-2} \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta-2/\beta}}{n \log n} \int_{\varepsilon(\log \log n)^{1/\beta}}^{\infty} 2yP\{\|\alpha_n\| \geq y\} dy \\ &+ \varepsilon^{\beta(\delta+1)} \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta}}{n \log n} P\{\|\alpha_n\| \geq \varepsilon(\log \log n)^{1/\beta}\}. \end{aligned} \quad (2.22)$$

Via the similar argument in Proposition 2.1 and 2.2,

$$\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)} \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta}}{n \log n} P\{\|\alpha_n\| \geq \varepsilon \|\log \log n\|^{1/\beta}\} = \frac{\mathbb{E}\|B\|^{\beta(\delta+1)}}{\delta+1}. \quad (2.23)$$

Also, by the analogous proof of Proposition 2.3 and 2.4,

$$\lim_{\varepsilon \searrow 0} \varepsilon^{\beta(\delta+1)-2} \sum_{n=3}^{\infty} \frac{(\log \log n)^{\delta-2/\beta}}{n \log n} \int_{\varepsilon(\log \log n)^{1/\beta}}^{\infty} 2yP\{\|\alpha_n\| \geq y\} dy = \frac{2\mathbb{E}\|B\|^{\beta(\delta+1)}}{(\delta+1)(\beta(\delta+1)-2)}. \quad (2.24)$$

Combine (2.22), (2.23), and (2.24) together, we get the result of Theorem 1.2. □

Acknowledgment

This work was supported by NSFC (No. 10771192) and ZJNSF (No. J20091364).

References

- [1] P. L. Hsu and H. Robbins, "Complete convergence and the law of large numbers," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 33, pp. 25–31, 1947.
- [2] P. Erdős, "On a theorem of Hsu and Robbins," *Annals of Mathematical Statistics*, vol. 20, pp. 286–291, 1949.
- [3] R. Chen, "A remark on the tail probability of a distribution," *Journal of Multivariate Analysis*, vol. 8, no. 2, pp. 328–333, 1978.
- [4] A. Gut and A. Spătaru, "Precise asymptotics in the Baum-Katz and Davis laws of large numbers," *Journal of Mathematical Analysis and Applications*, vol. 248, no. 1, pp. 233–246, 2000.
- [5] C. C. Heyde, "A supplement to the strong law of large numbers," *Journal of Applied Probability*, vol. 12, pp. 173–175, 1975.
- [6] Y. S. Chow, "On the rate of moment convergence of sample sums and extremes," *Bulletin of the Institute of Mathematics*, vol. 16, no. 3, pp. 177–201, 1988.
- [7] W. Liu and Z. Lin, "Precise asymptotics for a new kind of complete moment convergence," *Statistics & Probability Letters*, vol. 76, no. 16, pp. 1787–1799, 2006.
- [8] K.-A. Fu, "Asymptotics for the moment convergence of U-statistics in LIL," *Journal of Inequalities and Applications*, vol. 2010, Article ID 350517, 8 pages, 2010.
- [9] J. G. Yan and C. Su, "Precise asymptotics of U-statistics," *Acta Mathematica Sinica*, vol. 50, no. 3, pp. 517–526, 2007 (Chinese).
- [10] T.-X. Pang, L.-X. Zhang, and J. F. Wang, "Precise asymptotics in the self-normalized law of the iterated logarithm," *Journal of Mathematical Analysis and Applications*, vol. 340, no. 2, pp. 1249–1262, 2008.
- [11] Q.-P. Zang, "A limit theorem for the moment of self-normalized sums," *Journal of Inequalities and Applications*, vol. 2009, Article ID 957056, 10 pages, 2009.
- [12] Y. Zhang and X.-Y. Yang, "Precise asymptotics in the law of the iterated logarithm and the complete convergence for uniform empirical process," *Statistics & Probability Letters*, vol. 78, no. 9, pp. 1051–1055, 2008.
- [13] M. Csörgő and P. Révész, *Strong Approximations in Probability and Statistics*, Probability and Mathematical Statistics, Academic Press, New York, NY, USA, 1981.
- [14] W. F. Stout, *Almost Sure Convergence*, Academic Press, New York, NY, USA, 1974, Probability and Mathematical Statistics, Vol. 2.
- [15] J. Kiefer and J. Wolfowitz, "On the deviations of the empiric distribution function of vector chance variables," *Transactions of the American Mathematical Society*, vol. 87, pp. 173–186, 1958.