

Research Article

Superstability of Some Pexider-Type Functional Equation

Gwang Hui Kim

Department of Mathematics, Kangnam University, Yongin, Gyeonggi 446-702, Republic of Korea

Correspondence should be addressed to Gwang Hui Kim, ghkim@kangnam.ac.kr

Received 27 August 2010; Revised 18 October 2010; Accepted 19 October 2010

Academic Editor: Andrei Volodin

Copyright © 2010 Gwang Hui Kim. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We will investigate the superstability of the sine functional equation from the following Pexider-type functional equation $f(x+y)+g(x-y)=\lambda \cdot h(x)k(y)$ (λ : constant), which can be considered the mixed functional equation of the sine and cosine functions, the mixed functional equation of the hyperbolic sine and hyperbolic cosine functions, and the exponential-type functional equations.

1. Introduction

In 1940, Ulam [1] conjectured the stability problem. Next year, this problem was affirmatively solved by Hyers [2], which is through the following.

Let X and Y be Banach spaces with norm $\|\cdot\|$, respectively. If $f: X \rightarrow Y$ satisfies

$$\|f(x+y)-f(x)-f(y)\| \leq \varepsilon, \quad \forall x, y \in X, \quad (1.1)$$

then there exists a unique additive mapping $A: X \rightarrow Y$ such that

$$\|f(x)-A(x)\| \leq \varepsilon, \quad \forall x \in X. \quad (1.2)$$

The above result was generalized by Bourgin [3] and Aoki [4] in 1949 and 1950. In 1978 and 1982, Hyers' result was improved by Th. M. Rassias [5] and J. M. Rassias [6] which is that the condition bounded by the constant is replaced to the condition bounded by two variables, and thereafter it was improved moreover by Găvruta [7] to the condition bounded by the function.

In 1979, Baker et al. [8] showed that if f is a function from a vector space to R satisfying

$$|f(x+y) - f(x)f(y)| \leq \varepsilon, \quad (1.3)$$

then either f is bounded or satisfies the exponential functional equation

$$f(x+y) = f(x)f(y). \quad (1.4)$$

This method is referred to as the superstability of the functional equation (1.4).

In this paper, let $(G, +)$ be a uniquely 2 divisible Abelian group, \mathbb{C} the field of complex numbers, and \mathbb{R} the field of real numbers, \mathbb{R}_+ the set of positive reals. Whenever we only deal with (C), $(G, +)$ needs the Abelian which is not 2-divisible.

We may assume that f, g, h and k are nonzero functions, λ, ε is a nonnegative real constant, and $\varphi : G \rightarrow \mathbb{R}_+$ is a mapping.

In 1980, the superstability of the cosine functional equation (also referred the d'Alembert functional equation)

$$f(x+y) + f(x-y) = 2f(x)f(y), \quad (C)$$

was investigated by Baker [9] with the following result: let $\varepsilon > 0$. If $f : G \rightarrow \mathbb{C}$ satisfies

$$|f(x+y) + f(x-y) - 2f(x)f(y)| \leq \varepsilon, \quad (1.5)$$

then either $|f(x)| \leq (1 + \sqrt{1 + 2\varepsilon})/2$ for all $x \in G$ or f is a solution of (C). Badora [10] in 1998, and Badora and Ger [11] in 2002 under the condition $|f(x+y) + f(x-y) - 2f(x)f(y)| \leq \varepsilon$, $\varphi(x)$ or $\varphi(y)$, respectively. Also the stability of the d'Alembert functional equation is founded in papers [12–16].

In the present work, the stability question regarding a Pexider-type trigonometric functional equation as a generalization of the cosine equation (C) is investigated.

To be systematic, we first list all functional equations that are of interest here.

$$f(x+y) + g(x-y) = \lambda h(x)h(y) \quad (P_{fghh}^\lambda)$$

$$f(x+y) + g(x-y) = \lambda f(x)h(y), \quad (P_{fgh}^\lambda)$$

$$f(x+y) + g(x-y) = \lambda h(x)f(y), \quad (P_{fghf}^\lambda)$$

$$\begin{aligned}
f(x+y) + g(x-y) &= \lambda g(x)h(y), & (P_{fgh}^\lambda) \\
f(x+y) + g(x-y) &= \lambda h(x)g(y), & (P_{fgh}^\lambda) \\
f(x+y) + g(x-y) &= \lambda f(x)g(y), & (P_{fgf}^\lambda) \\
f(x+y) + g(x-y) &= \lambda g(x)f(y), & (P_{fgf}^\lambda) \\
f(x+y) + g(x-y) &= \lambda f(x)f(y), & (P_{fgf}^\lambda) \\
f(x+y) + g(x-y) &= \lambda g(x)g(y), & (P_{fgg}^\lambda) \\
f(x+y) + f(x-y) &= \lambda g(x)h(y), & (P_{fgh}^\lambda) \\
f(x+y) + f(x-y) &= \lambda g(x)g(y), & (P_{fgg}^\lambda) \\
f(x+y) + f(x-y) &= \lambda f(x)g(y), & (C_{fg}^\lambda) \\
f(x+y) + f(x-y) &= \lambda g(x)f(y), & (C_{gf}^\lambda) \\
f(x+y) + f(x-y) &= \lambda f(x)f(y), & (C^\lambda) \\
f(x+y) + g(x-y) &= 2h(x)k(y), & (P_{fghk}) \\
f(x+y) + g(x-y) &= 2h(x)h(y), & (P_{fghh}) \\
f(x+y) + g(x-y) &= 2f(x)h(y), & (P_{fghh}) \\
f(x+y) + g(x-y) &= 2h(x)f(y), & (P_{fghf}) \\
f(x+y) + g(x-y) &= 2g(x)h(y), & (P_{fggh}) \\
f(x+y) + g(x-y) &= 2h(x)g(y), & (P_{fghg}) \\
f(x+y) + g(x-y) &= 2f(x)g(y), & (P_{fgfg}) \\
f(x+y) + g(x-y) &= 2g(x)f(y), & (P_{fggf}) \\
f(x+y) + g(x-y) &= 2f(x)f(y), & (P_{fgff}) \\
f(x+y) + g(x-y) &= 2g(x)g(y), & (P_{fggg}) \\
f(x+y) + f(x-y) &= 2f(x)g(y), & (C_{fg}) \\
f(x+y) + f(x-y) &= 2g(x)f(y), & (C_{gf}) \\
f(x+y) + f(x-y) &= 2g(x)g(y), & (C_{gg}) \\
f(x+y) + f(x-y) &= 2g(x)h(y), & (C_{gh}) \\
f(x+y) + f(x-y) &= 2f(x). & (J_x)
\end{aligned}$$

The hyperbolic cosine function, hyperbolic sine function, hyperbolic trigonometric function, some exponential functions, and Jensen equation satisfy the above mentioned

equations; therefore, they can also be called the *hyperbolic* cosine (sine, trigonometric) functional equation, exponential functional equation, and Jensen equation, respectively.

For example,

$$\begin{aligned}
 \cosh(x+y) + \cosh(x-y) &= 2 \cosh(x) \cosh(y), \\
 \cosh(x+y) - \cosh(x-y) &= 2 \sinh(x) \sinh(y), \\
 \sinh(x+y) + \sinh(x-y) &= 2 \sinh(x) \cosh(y), \\
 \sinh(x+y) - \sinh(x-y) &= 2 \cosh(x) \sinh(y), \\
 \sinh^2\left(\frac{x+y}{2}\right) - \sinh^2\left(\frac{x-y}{2}\right) &= \sinh(x) \sinh(y), \\
 ca^{x+y} + ca^{x-y} &= 2 \frac{ca^x}{2} (a^y + a^{-y}) = 2ce^x \frac{a^y + a^{-y}}{2}, \\
 e^{x+y} + e^{x-y} &= 2 \frac{e^x}{2} (e^y + e^{-y}) = 2e^x \cosh(y), \\
 (n(x+y) + c) + (n(x-y) + c) &= 2(nx + c) : \text{for } f(x) = nx + c,
 \end{aligned} \tag{1.6}$$

where a and c are constants.

The equation (C_{fg}) is referred to as the Wilson equation. In 2001, Kim and Kannappan [13] investigated the superstability related to the d'Alembert (C) and the Wilson functional equations (C_{fg}) , (C_{gf}) under the condition bounded by constant. Kim has also improved the superstability of the generalized cosine type-functional equations (C_{gg}) , and (P_{fgfg}) , (P_{fggf}) in papers [14, 15, 17].

In particular, author Kim and Lee [18] investigated the superstability of (S) from the functional equation (C_{gh}) under the condition bounded by function, that is

(1) if $f, g, h : G \rightarrow C$ satisfies

$$|f(x+y) + f(x-y) - 2g(x)h(y)| \leq \varphi(x), \tag{1.7}$$

then either h is bounded or g satisfies (S);

(2) if $f, g, h : G \rightarrow C$ satisfies

$$|f(x+y) + f(x-y) - 2g(x)h(y)| \leq \varphi(y), \tag{1.8}$$

then either g is bounded or h satisfies (S).

In 1983, Cholewa [19] investigated the superstability of the sine functional equation

$$f(x)f(y) = f\left(\frac{x+y}{2}\right)^2 - f\left(\frac{x-y}{2}\right)^2, \tag{S}$$

under the condition bounded by constant. Namely, if an unbounded function $f : G \rightarrow \mathbb{C}$ satisfies

$$\left| f(x)f(y) - f\left(\frac{x+y}{2}\right)^2 + f\left(\frac{x-y}{2}\right)^2 \right| \leq \varepsilon, \quad (1.9)$$

then it satisfies (S).

In Kim's work [20, 21], the superstability of sine functional equation from the generalized sine functional equations

$$f(x)g(y) = f\left(\frac{x+y}{2}\right)^2 - f\left(\frac{x-y}{2}\right)^2, \quad (S_{fg})$$

$$g(x)f(y) = f\left(\frac{x+y}{2}\right)^2 - f\left(\frac{x-y}{2}\right)^2, \quad (S_{gf})$$

$$g(x)h(y) = f\left(\frac{x+y}{2}\right)^2 - f\left(\frac{x-y}{2}\right)^2 \quad (S_{gh})$$

was treated under the conditions bounded by constant and functions.

The aim of this paper is to investigate the transferred superstability for the sine functional equation from the following Pexider type functional equations:

$$f(x+y) + g(x-y) = \lambda \cdot h(x)k(y), \quad \lambda : \text{constant} \quad (P_{fghk}^\lambda)$$

on the abelian group. Furthermore, the obtained results can be extended to the Banach space.

Consequently, as corollaries, we can obtain 29×4 stability results concerned with the sine functional equation (S) and the Wilson-type equations (C_{fg}^λ) from 29 functional equations of the P^λ, C^λ, P , and C types from a selection of functions f, g, h, k in the order of variables $x+y, x-y, x, y$.

2. Superstability of the Sine Functional Equation from the Equation (P_{fghk}^λ)

In this section, we will investigate the superstability related to the d'Alembert-type equation (C^λ) and Wilson-type equation (C_{fg}^λ) , of the sine functional equation (S) from the Pexider type functional equation (P_{fghk}^λ) .

Theorem 2.1. Suppose that $f, g, h, k : G \rightarrow \mathbb{C}$ satisfy the inequality

$$|f(x+y) + g(x-y) - \lambda \cdot h(x)k(y)| \leq \varphi(x), \quad \forall x, y \in G. \quad (2.1)$$

If k fails to be bounded, then

(i) h satisfies (S) under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$; and

(ii) In addition, if k satisfies (C^λ) , then h and k are solutions of $(C_{fg}^\lambda) := h(x+y) + h(x-y) = \lambda h(x)k(y)$.

Proof. Let k be unbounded solution of the inequality (3.12). Then, there exists a sequence $\{y_n\}$ in G such that $0 \neq |k(y_n)| \rightarrow \infty$ as $n \rightarrow \infty$.

(i) Taking $y = y_n$ in the inequality (3.12), dividing both sides by $|\lambda k(y_n)|$, and passing to the limit as $n \rightarrow \infty$, we obtain

$$h(x) = \lim_{n \rightarrow \infty} \frac{f(x+y_n) + g(x-y_n)}{\lambda \cdot k(y_n)}, \quad x \in G. \quad (2.2)$$

Replace y by $y + y_n$ and $-y + y_n$ in (3.12), we have

$$\begin{aligned} & |f(x + (y + y_n)) + g(x - (y + y_n)) - \lambda \cdot h(x)k(y + y_n) \\ & f(x + (-y + y_n)) + g(x - (-y + y_n)) - \lambda \cdot h(x)k(-y + y_n)| \leq 2\varphi(x) \end{aligned} \quad (2.3)$$

so that

$$\begin{aligned} & \left| \frac{f((x+y) + y_n) + g((x+y) - y_n)}{\lambda \cdot k(y_n)} \right. \\ & \left. + \frac{f((x-y) + y_n) + g((x-y) - y_n)}{\lambda \cdot k(y_n)} - \lambda \cdot h(x) \cdot \frac{k(y+y_n) + k(-y+y_n)}{\lambda \cdot k(y_n)} \right| \leq \frac{2\varphi(x)}{\lambda \cdot |k(y_n)|} \end{aligned} \quad (2.4)$$

for all $x, y, y_n \in G$.

We conclude that, for every $y \in G$, there exists a limit function

$$l_k(y) := \lim_{n \rightarrow \infty} \frac{k(y+y_n) + k(-y+y_n)}{\lambda \cdot k(y_n)}, \quad (2.5)$$

where the function $l_k : G \rightarrow \mathbb{C}$ satisfies the equation

$$h(x+y) + h(x-y) = \lambda \cdot h(x)l_k(y), \quad \forall x, y \in G. \quad (2.6)$$

Applying the case $h(0) = 0$ in (2.6), it implies that h is odd. Keeping this in mind, by means of (2.6), we infer the equality

$$\begin{aligned} h(x+y)^2 - h(x-y)^2 &= \lambda \cdot h(x)l_k(y) [h(x+y) - h(x-y)] \\ &= h(x) [h(x+2y) - h(x-2y)] \\ &= h(x) [h(2y+x) + h(2y-x)] \\ &= \lambda \cdot h(x)h(2y)l_k(x). \end{aligned} \quad (2.7)$$

Putting $y = x$ in (2.6), we get the equation

$$h(2x) = \lambda \cdot h(x)l_k(x), \quad x \in G. \quad (2.8)$$

This, in return, leads to the equation

$$h(x+y)^2 - h(x-y)^2 = h(2x)h(2y) \quad (2.9)$$

valid for all $x, y \in G$ which, in the light of the unique 2divisibility of G , states nothing else but (S).

In the particular case $f(-x) = -g(x)$, it is enough to show that $h(0) = 0$. Suppose that this is not the case.

Putting $x = 0$ in (3.12), due to $h(0) \neq 0$ and $f(-x) = -g(x)$, we obtain the inequality

$$|k(y)| \leq \frac{\varphi(0)}{\lambda \cdot |h(0)|}, \quad y \in G. \quad (2.10)$$

This inequality means that k is globally bounded, which is a contradiction. Thus, since the claimed $h(0) = 0$ holds, we know that h satisfies (S).

(ii) In the case k satisfies (C^λ) , the limit l_k states nothing else but k , so, from (2.6), h and k validate (C_{fg}^λ) . \square

Theorem 2.2. Suppose that $f, g, h, k : G \rightarrow \mathbb{C}$ satisfy the inequality

$$|f(x+y) + g(x-y) - \lambda \cdot h(x)k(y)| \leq \varphi(y) \quad \forall x, y \in G. \quad (2.11)$$

If h fails to be bounded, then

(i) k satisfies (S) under one of the cases $k(0) = 0$ or $f(x) = -g(x)$

(ii) in addition, if h satisfies (C^λ) , then k and h are solutions of the equation of $(C_{gf}^\lambda) := k(x+y) + k(x-y) = \lambda h(x)k(y)$.

Proof. (i) Taking $x = x_n$ in the inequality (2.11), dividing both sides by $|\lambda \cdot h(x_n)|$, and passing to the limit as $n \rightarrow \infty$, we obtain that

$$k(y) = \lim_{n \rightarrow \infty} \frac{f(x_n+y) + g(x_n-y)}{\lambda \cdot h(x_n)}, \quad x \in G. \quad (2.12)$$

Replace x by $x_n + x$ and $x_n - x$ in (2.11) divide by $\lambda \cdot h(x_n)$; then it gives us the existence of the limit function

$$l_h(x) := \lim_{n \rightarrow \infty} \frac{h(x_n+x) + h(x_n-x)}{\lambda \cdot h(x_n)}, \quad (2.13)$$

where the function $l_h : G \rightarrow \mathbb{C}$ satisfies the equation

$$k(x+y) + k(-x+y) = \lambda \cdot l_h(x)k(y), \quad \forall x, y \in G. \quad (2.14)$$

Applying the case $k(0) = 0$ in (2.14), it implies that k is odd.

A similar procedure to that applied after (2.6) of Theorem 2.1 in (2.14) allows us to show that k satisfies (S).

The case $f(x) = -g(x)$ is also the same as the reason for Theorem 2.1.

(ii) In the case h satisfies (C^λ) , the limit l_h states nothing else but h , so, from (2.14), k and h validate (C_{fg}^λ) . \square

The following corollaries follow immediately from the Theorems 2.1 and 2.2.

Corollary 2.3. Suppose that $f, g, h, k : G \rightarrow \mathbb{C}$ satisfy the inequality

$$|f(x+y) + g(x-y) - \lambda \cdot h(x)k(y)| \leq \min\{\phi(x), \phi(y)\}, \quad \forall x, y \in G. \quad (2.15)$$

(a) If k fails to be bounded, then

- (i) h satisfies (S) under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$, and
- (ii) in addition, if h satisfies (C^λ) , then h and k are solutions of $(C_{fg}^\lambda) := h(x+y) + h(x-y) = \lambda h(x)k(y)$.

(b) If h fails to be bounded, then

- (iii) k satisfies (S) under one of the cases $k(0) = 0$ or $f(x) = -g(x)$, and
- (iv) in addition, if h satisfies (C^λ) , then h and k are solutions of $(C_{gf}^\lambda) := k(x+y) + k(x-y) = \lambda h(x)k(y)$.

Corollary 2.4. Suppose that $f, g, h, k : G \rightarrow \mathbb{C}$ satisfy the inequality

$$|f(x+y) + g(x-y) - \lambda \cdot h(x)k(y)| \leq \varepsilon, \quad \forall x, y \in G. \quad (2.16)$$

(a) If k fails to be bounded, then

- (i) h satisfies (S) under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$, and
- (ii) in addition, if k satisfies (C^λ) , then h and k are solutions of $(C_{fg}^\lambda) := h(x+y) + h(x-y) = \lambda h(x)k(y)$.

(b) If h fails to be bounded, then

- (iii) k satisfies (S) under one of the cases $k(0) = 0$ or $f(x) = -g(x)$, and
- (iv) in addition, if h satisfies (C^λ) , then h and k are solutions of $(C_{gf}^\lambda) := k(x+y) + k(x-y) = \lambda h(x)k(y)$.

3. Applications in the Reduced Equations

3.1. Corollaries of the Equations Reduced to Three Unknown Functions

Replacing k by one of the functions f, g, h in all the results of the Section 2 and exchanging each functions f, g, h in the above equations, we then obtain P^λ, C^λ types 14 equations.

We will only illustrate the results for the cases of $(P_{fghh}^\lambda), (P_{fgfh}^\lambda)$ in the obtained equations. The other cases are similar to these; thus their illustrations will be omitted.

Corollary 3.1. Suppose that $f, g, h : G \rightarrow \mathbb{C}$ satisfy the inequality

$$|f(x+y) + g(x-y) - \lambda \cdot h(x)h(y)| \leq \begin{cases} \varphi(x) & \text{or} \\ \varphi(y) & \text{or} \\ \min\{\varphi(x), \varphi(y)\} & \text{or} \\ \varepsilon \end{cases} \quad \forall x, y \in G. \quad (3.1)$$

If h fails to be bounded, then, under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$, h satisfies (S).

Corollary 3.2. Suppose that $f, g, h : G \rightarrow \mathbb{C}$ satisfy the inequality

$$|f(x+y) + g(x-y) - \lambda \cdot f(x)h(y)| \leq \varphi(x), \quad \forall x, y \in G. \quad (3.2)$$

If h fails to be bounded, then

- (i) f satisfies (S) under one of the cases $f(0) = 0$ or $f(-x) = -g(x)$, and
- (ii) in addition, if h satisfies (C^λ) , then f and h are solutions of $(C_{fg}) := f(x+y) + f(x-y) = \lambda \cdot f(x)h(y)$.

Corollary 3.3. Suppose that $f, g, h : G \rightarrow \mathbb{C}$ satisfy the inequality

$$|f(x+y) + g(x-y) - \lambda \cdot f(x)h(y)| \leq \varphi(y), \quad \forall x, y \in G. \quad (3.3)$$

If f fails to be bounded, then

- (i) h satisfies (S) under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$, and
- (ii) in addition, if f satisfies (C^λ) , then h and f are solutions of $(C_{gf}) := h(x+y) + h(x-y) = \lambda \cdot f(x)h(y)$.

Corollary 3.4. Suppose that $f, g, h : G \rightarrow \mathbb{C}$ satisfy the inequality

$$|f(x+y) + g(x-y) - \lambda \cdot f(x)h(y)| \leq \min\{\varphi(x), \varphi(y)\}, \quad \forall x, y \in G. \quad (3.4)$$

(a) If h fails to be bounded, then

- (i) f satisfies (S) under one of the cases $f(0) = 0$ or $f(-x) = -g(x)$, and
- (ii) in addition, if h satisfies (C^λ) , then f and h are solutions of $(C_{fg}^\lambda) := f(x+y) + f(x-y) = \lambda \cdot f(x)h(y)$.

(b) If f fails to be bounded, then

- (i) h satisfies (S) under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$, and
- (ii) in addition, if f satisfies (C^λ) , then h and f are solutions of $(C_{gf}^\lambda) := h(x+y) + h(x-y) = \lambda \cdot f(x)h(y)$.

Corollary 3.5. Suppose that $f, g, h : G \rightarrow \mathbb{C}$ satisfy the inequality

$$|f(x+y) + g(x-y) - \lambda \cdot f(x)h(y)| \leq \varepsilon, \quad \forall x, y \in G. \quad (3.5)$$

(a) If h fails to be bounded, then

- (i) f satisfies (S) under one of the cases $f(0) = 0$ or $f(-x) = -g(x)$, and
- (ii) in addition, if h satisfies (C^λ) , then f and h are solutions of $(C_{fg}^\lambda) := f(x+y) + f(x-y) = \lambda \cdot f(x)h(y)$.

(b) If f fails to be bounded, then

- (i) h satisfies (S) under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$, and
- (ii) in addition, if f satisfies (C^λ) , then h and f are solutions of $(C_{gf}^\lambda) := h(x+y) + h(x-y) = \lambda \cdot f(x)h(y)$.

Remark 3.6. As the above corollaries, we obtain the stability results of $12 \times 4(\varphi(x), \varphi(y), \min\{\varphi(x), \varphi(y)\}, \varepsilon)$ numbers for 12 equations by choosing f, g, h , and λ , namely, which are the following: $(P_{fghf}^\lambda), (P_{fghh}^\lambda), (P_{fghg}^\lambda), (P_{fgfg}^\lambda), (P_{fggf}^\lambda), (P_{fgff}^\lambda), (P_{fggg}^\lambda), (P_{ffgh}^\lambda), (P_{ffgg}^\lambda), (C_{fg}^\lambda), (C_{gf}^\lambda)$, and (C^λ) .

3.2. Applications of the Case $\lambda = 2$ in (P_{fghk}^λ)

Let us apply the case $\lambda = 2$ in (P_{fghk}^λ) and all P^λ -type equations considered in the Sections 2 and Sec 3.1. Then, we obtain the P -type equations

$$f(x+y) + g(x-y) = 2 \cdot h(x)k(y), \quad (P_{fghk})$$

and $(P_{fghh}^\lambda), (P_{fghf}^\lambda), (P_{fghg}^\lambda), (P_{fgfg}^\lambda), (P_{fggf}^\lambda), (P_{fgff}^\lambda), (P_{fggg}^\lambda)$, and C - and J -type $(C_{fg}), (C_{gf}), (C_{gg}), (C_{gh}), (C),$ and (J_x) , which are concerned with the (hyperbolic) cosine, sine, exponential functions, and Jensen equation.

In papers (Aczél [22], Aczél and Dhombres [23], Kannappan [24, 25], and Kim and Kannappan [13]), we can find that the Wilson equation and the sine equations can be represented by the composition of a homomorphism. By applying these results, we also obtain, additionally, the explicit solutions of the considered functional equations.

Corollary 3.7. *Suppose that $f, g, h, k : G \rightarrow \mathbb{C}$ satisfy the inequality*

$$|f(x+y) + g(x-y) - 2h(x)k(y)| \leq \varphi(x) \quad \forall x, y \in G. \quad (3.6)$$

If k fails to be bounded, then

(i) *h satisfies (S) under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$, and h is of the form*

$$h(x) = A(x) \quad \text{or} \quad h(x) = c(E(x) - E^*(x)), \quad (3.7)$$

where $A : G \rightarrow \mathbb{C}$ is an additive function, $c \in \mathbb{C}$, $E : G \rightarrow \mathbb{C}^$ is a homomorphism and $E^* = 1/E(x)$,*

(ii) *in addition, if k satisfies (C), then h and k are solutions of (C_{fg}) and h, k are given by*

$$k(x) = \frac{E(x) + E^*(x)}{2}, \quad h(x) = c(E(x) - E^*(x)) + \frac{d(E(x) + E^*(x))}{2}, \quad (3.8)$$

where $c, d \in \mathbb{C}$, E and E^ are as in (i).*

Proof. The proof of the Corollary is enough from Theorem 2.1 except for the solution. However, they are immediate from the following:

- (i) appealing to the solutions of (S) in [2, page 153] (see also [24, 25]), the explicit shapes of h are as stated in the statement of the theorem. This completes the proof of the Corollary,
- (ii) the given explicit solutions are taken from [24, 25] (page 148) (see also [22, 23]). \square

Corollary 3.8. *Suppose that $f, g, h, k : G \rightarrow \mathbb{C}$ satisfy the inequality*

$$|f(x+y) + g(x-y) - 2h(x)k(y)| \leq \varphi(y) \quad \forall x, y \in G. \quad (3.9)$$

If h fails to be bounded, then

(i) *k satisfies (S) under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$, and k is of the form*

$$k(x) = A(x) \quad \text{or} \quad k(x) = c(E(x) - E^*(x)), \quad (3.10)$$

where $A : G \rightarrow \mathbb{C}$ is an additive function, $c \in \mathbb{C}$, $E : G \rightarrow \mathbb{C}^$ is a homomorphism and $E^* = 1/E(x)$, and*

(ii) in addition, if h satisfies (C), then k and h are solutions of (C_{fg}) and k, h are given by

$$h(x) = \frac{E(x) + E^*(x)}{2}, \quad k(x) = c(E(x) - E^*(x)) + \frac{d(E(x) + E^*(x))}{2}, \quad (3.11)$$

where $c, d \in \mathbb{C}$, E and E^* are as in (i).

Corollary 3.9. Suppose that $f, g, h, k : G \rightarrow \mathbb{C}$ satisfy the inequality

$$|f(x+y) + g(x-y) - 2h(x)k(y)| \leq \begin{cases} \min\{\varphi(x), \varphi(y)\} \\ \varepsilon \end{cases} \quad \forall x, y \in G. \quad (3.12)$$

(a) If k fails to be bounded, then

- (i) h satisfies (S) under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$, and h is of the form $h(x) = A(x)$ or $h(x) = c(E(x) - E^*(x))$, and where $A : G \rightarrow \mathbb{C}$ is an additive function, $c \in \mathbb{C}$, $E : G \rightarrow \mathbb{C}^*$ is a homomorphism and $E^* = 1/E(x)$.
- (ii) in addition, if k satisfies (C), then h and k are solutions of (C_{fg}) and h, k are given by

$$k(x) = \frac{E(x) + E^*(x)}{2}, \quad h(x) = c(E(x) - E^*(x)) + \frac{d(E(x) + E^*(x))}{2}, \quad (3.13)$$

where $c, d \in \mathbb{C}$, E and E^* are as in (i).

(b) If h fails to be bounded, then

- (i) k satisfies (S) under one of the cases $h(0) = 0$ or $f(-x) = -g(x)$, and k is of the form

$$k(x) = A(x) \quad \text{or} \quad k(x) = c(E(x) - E^*(x)), \quad (3.14)$$

where $A : G \rightarrow \mathbb{C}$ is an additive function, $c \in \mathbb{C}$, $E : G \rightarrow \mathbb{C}^*$ is a homomorphism and $E^* = 1/E(x)$, and

- (ii) in addition, if h satisfies (C), then k and h are solutions of (C_{fg}) and k, h are given by

$$h(x) = \frac{E(x) + E^*(x)}{2}, \quad k(x) = c(E(x) - E^*(x)) + \frac{d(E(x) + E^*(x))}{2}, \quad (3.15)$$

where $c, d \in \mathbb{C}$, E and E^* are as in (i).

Remark 3.10. Applying the case $\lambda = 2$ in the first paragraph of the Section 3.2 implies the above 15 equations. Therefore, as the above Corollaries 3.7, 3.8, and 3.9, we can obtain additionally the stability results of $14 \times 4(\varphi(x), \varphi(y), \min\{\varphi(x), \varphi(y)\}, \varepsilon)$ numbers for the other 14 equations. Some, which excepted the explicit solutions represented by composition of a homomorphism, in the obtained results are found in papers [7, 11, 13–15, 17].

4. Extension to the Banach Space

In all the results presented in Sections 2 and 3, the range of functions on the Abelian group can be extended to the semisimple commutative Banach space. We will represent just for the main equation (P_{fghk}^λ) .

Theorem 4.1. *Let $(E, \|\cdot\|)$ be a semisimple commutative Banach space. Assume that $f, g, h, k : G \rightarrow E$ satisfy one of each inequalities*

$$\|f(x+y) + g(x-y) - \lambda \cdot h(x)k(y)\| \leq \varphi(x), \quad (4.1)$$

$$\|f(x+y) + g(x-y) - \lambda \cdot h(x)k(y)\| \leq \varphi(y) \quad (4.2)$$

for all $x, y \in G$. For an arbitrary linear multiplicative functional $x^* \in E^*$.

(a) case (4.1)

Suppose that $x^* \circ k$ fails to be bounded, then

- (i) h satisfies (S) under one of the cases $x^* \circ h(0) = 0$ or $x^* \circ f(-x) = -x^* \circ g(x)$, and
- (ii) in addition, if k satisfies (C^λ) , then h and k are solutions of (C_{fg}^λ) .

(b) Case (4.2)

Suppose that $x^* \circ h$ fails to be bounded, then

- (iii) k satisfies (S) under one of the cases $x^* \circ k(0) = 0$ or $x^* \circ f(-x) = -x^* \circ g(x)$, and
- (iv) in addition, if $x^* \circ h$ satisfies (C^λ) , then h and k are solutions of (C_{fg}^λ) .

Proof. For (i) of (a), assume that (4.1) holds and arbitrarily fixes a linear multiplicative functional $x^* \in E^*$. As is well known, we have $\|x^*\| = 1$; hence, for every $x, y \in G$, we have

$$\begin{aligned} \varphi(x) &\geq \|f(x+y) + g(x-y) - \lambda \cdot h(x)k(y)\| \\ &= \sup_{\|y^*\|=1} |y^*(f(x+y) + g(x-y) - \lambda \cdot h(x)k(y))| \\ &\geq |x^*(f(x+y)) + x^*(g(x-y)) - \lambda \cdot x^*(h(x))x^*(k(y))|, \end{aligned} \quad (4.3)$$

which states that the superpositions $x^* \circ f$, $x^* \circ g$, $x^* \circ h$, and $x^* \circ k$ yield a solution of inequality (2.1) in Theorem 2.1. Since, by assumption, the superposition $x^* \circ k$ with $x^* \circ h(0) = 0$ is unbounded, an appeal to Theorem 2.1 shows that the two results hold.

First, the superposition $x^* \circ h$ solves (S), that is

$$(x^* \circ h)\left(\frac{x+y}{2}\right)^2 - (x^* \circ h)\left(\frac{x-y}{2}\right)^2 = (x^* \circ h)(x)(x^* \circ h)(y). \quad (4.4)$$

Since x^* is a linear multiplicative functional, we get

$$x^* \left(h \left(\frac{x+y}{2} \right)^2 - h \left(\frac{x-y}{2} \right)^2 - h(x)h(y) \right) = 0. \quad (4.5)$$

Hence an unrestricted choice of x^* implies that

$$h \left(\frac{x+y}{2} \right)^2 - h \left(\frac{x-y}{2} \right)^2 - h(x)h(y) \in \bigcap \{ \ker x^* : x^* \in E^* \}. \quad (4.6)$$

Since the space E is semisimple, $\bigcap \{ \ker x^* : x^* \in E^* \} = 0$, which means that h satisfies the claimed equation (S).

For second case $(x^* \circ f)(-x) = -(x^* \circ g)(x)$, it is enough to show that $(x^* \circ h)(0) = 0$, which can be easily check as Theorem 2.1. Hence, the proof (i) of (a) is completed.

For (ii) of (a), as (i) of (a), an appeal to Theorem 2.1 shows that if $x^* \circ k$ satisfies (C^λ) , then $x^* \circ h$ and $x^* \circ k$ are solutions of the Wilson-type equation

$$(x^* \circ h)(x+y) + (x^* \circ h)(x-y) = \lambda(x^* \circ h)(x)(x^* \circ k)(y). \quad (4.7)$$

This means by a linear multiplicativity of x^* that

$$\mathfrak{D}C_{hk}^\lambda(x, y) := h(x+y) + h(x-y) - \lambda h(x)k(y) \quad (4.8)$$

falls into the kernel of x^* . As the above process, since x^* is a linear multiplicative, we obtain

$$\mathfrak{D}C_{hk}^\lambda(x, y) = 0, \quad \forall x, y \in G \quad (4.9)$$

as claimed.

(b) the case (4.2) also runs along the proof of case (4.1). \square

Theorem 4.2. Let $(E, \|\cdot\|)$ be a semisimple commutative Banach space. Assume that $f, g, h, k : G \rightarrow E$ satisfy one of each inequalities

$$\|f(x+y) + g(x-y) - \lambda \cdot h(x)k(y)\| \leq \begin{cases} \min\{\varphi(x), \varphi(y)\} & \text{or} \\ \varepsilon \end{cases} \quad \forall x, y \in G. \quad (4.10)$$

for all $x, y \in G$. For an arbitrary linear multiplicative functional $x^* \in E^*$,

(a) suppose that $x^* \circ k$ fails to be bounded, then

- (i) h satisfies (S) under one of the cases $x^* \circ h(0) = 0$ or $x^* \circ f(-x) = -x^* \circ g(x)$, and
- (ii) in addition, if k satisfies (C^λ) , then h and k are solutions of (C_{fg}^λ) .

(b) Suppose that $x^* \circ h$ fails to be bounded, then

(iii) k satisfies (S) under one of the cases $x^* \circ k(0) = 0$ or $x^* \circ f(-x) = -x^* \circ g(x)$, and

(iv) in addition, if $x^* \circ h$ satisfies (C^λ) , then h and k are solutions of (C_{fg}^λ) .

Remark 4.3. As in the Remark 3.10, we can apply all results of the Sections 2 and 3 to the Banach space.

Namely, we obtain the stability results of $14 \times 4(\varphi(x), \varphi(y), \min\{\varphi(x), \varphi(y)\}, \varepsilon)$ numbers for the other 14 equations except for (P_{fghk}^λ) . Some of them are found in papers [7, 11, 13–15, 17, 18].

Acknowledgment

This work was supported by a Kangnam University research grant in 2009.

References

- [1] S. M. Ulam, *Problems in Modern Mathematics*, Science Editions John Wiley & Sons, New York, NY, USA, 1964.
- [2] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [3] D. G. Bourgin, "Approximately isometric and multiplicative transformations on continuous function rings," *Duke Mathematical Journal*, vol. 16, pp. 385–397, 1949.
- [4] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.
- [5] Th. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [6] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Journal of Functional Analysis*, vol. 46, no. 1, pp. 126–130, 1982.
- [7] P. Găvruta, "On the stability of some functional equations," in *Stability of Mappings of Hyers-Ulam Type*, Hadronic Press Collection of Original Articles, pp. 93–98, Hadronic Press, Palm Harbor, Fla, USA, 1994.
- [8] J. Baker, J. Lawrence, and F. Zorzitto, "The stability of the equation $f(x+y) = f(x)f(y)$," *Proceedings of the American Mathematical Society*, vol. 74, no. 2, pp. 242–246, 1979.
- [9] J. A. Baker, "The stability of the cosine equation," *Proceedings of the American Mathematical Society*, vol. 80, no. 3, pp. 411–416, 1980.
- [10] R. Badora, "On the stability of the cosine functional equation," *Rocznik Naukowo-Dydaktyczny. Prace Matematyczne*, no. 15, pp. 5–14, 1998.
- [11] R. Badora and R. Ger, "On some trigonometric functional inequalities," in *Functional Equations—Results and Advances*, vol. 3, pp. 3–15, Kluwer Academic Publishers, Dodrecht, The Netherlands, 2002.
- [12] B. Bouikhalene, E. Elqorachi, and J. M. Rassias, "The superstability of d'Alembert's functional equation on the Heisenberg group," *Applied Mathematics Letters*, vol. 23, no. 1, pp. 105–109, 2010.
- [13] G. H. Kim and Pl. Kannappan, "On the stability of the generalized cosine functional equations," *Annales Academiæ Paedagogicæ Cracoviensis—Studia Mathematica*, vol. 1, pp. 49–58, 2001.
- [14] G. H. Kim, "The stability of pexiderized cosine functional equations," *Korean Journal of Mathematics*, vol. 16, no. 1, pp. 103–114, 2008.
- [15] G. H. Kim, "The stability of d'Alembert and Jensen type functional equations," *Journal of Mathematical Analysis and Applications*, vol. 325, no. 1, pp. 237–248, 2007.
- [16] L. Székelyhidi, "The stability of d'Alembert-type functional equations," *Acta Universitatis Szegediensis. Acta Scientiarum Mathematicarum*, vol. 44, no. 3-4, pp. 313–320, 1982.
- [17] G. H. Kim, "On the superstability of the Pexider type trigonometric functional equation," *Journal of Inequalities and Applications*, vol. 2010, Article ID 897123, 14 pages, 2010.
- [18] G. H. Kim and Y. H. Lee, "The superstability of the Pexider type trigonometric functional equation," *Australian Journal of Mathematical Analysis*. In press.

- [19] P. W. Cholewa, "The stability of the sine equation," *Proceedings of the American Mathematical Society*, vol. 88, no. 4, pp. 631–634, 1983.
- [20] G. H. Kim, "A stability of the generalized sine functional equations," *Journal of Mathematical Analysis and Applications*, vol. 331, no. 2, pp. 886–894, 2007.
- [21] G. H. Kim, "On the stability of the generalized sine functional equations," *Acta Mathematica Sinica*, vol. 25, no. 1, pp. 29–38, 2009.
- [22] J. Aczél, *Lectures on Functional Equations and Their Applications*, Mathematics in Science and Engineering, Vol. 19, Academic Press, New York, NY, USA, 1966.
- [23] J. Aczél and J. Dhombres, *Functional Equations in Several Variables*, vol. 31 of *Encyclopedia of Mathematics and Its Applications*, Cambridge University Press, Cambridge, UK, 1989.
- [24] P. Kannappan, "The functional equation $f(xy) + f(xy^{-1}) = 2f(x)f(y)$ for groups," *Proceedings of the American Mathematical Society*, vol. 19, pp. 69–74, 1968.
- [25] P. Kannappan, *Functional Equations and Inequalities with Applications*, Springer Monographs in Mathematics, Springer, New York, NY, USA, 2009.