

## Research Article

# Some Shannon-McMillan Approximation Theorems for Markov Chain Field on the Generalized Bethe Tree

Kangkang Wang<sup>1</sup> and Decai Zong<sup>2</sup>

<sup>1</sup> School of Mathematics and Physics, Jiangsu University of Science and Technology, Zhenjiang 212003, China

<sup>2</sup> College of Computer Science and Engineering, Changshu Institute of Technology, Changshu 215500, China

Correspondence should be addressed to Wang Kangkang, wkk.cn@126.com

Received 26 September 2010; Accepted 7 January 2011

Academic Editor: Józef Banaś

Copyright © 2011 W. Kangkang and D. Zong. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A class of small-deviation theorems for the relative entropy densities of arbitrary random field on the generalized Bethe tree are discussed by comparing the arbitrary measure  $\mu$  with the Markov measure  $\mu_Q$  on the generalized Bethe tree. As corollaries, some Shannon-McMillan theorems for the arbitrary random field on the generalized Bethe tree, Markov chain field on the generalized Bethe tree are obtained.

## 1. Introduction and Lemma

Let  $T$  be a tree which is infinite, connected and contains no circuits. Given any two vertices  $x \neq y \in T$ , there exists a unique path  $x = x_1, x_2, \dots, x_m = y$  from  $x$  to  $y$  with  $x_1, x_2, \dots, x_m$  distinct. The distance between  $x$  and  $y$  is defined to  $m - 1$ , the number of edges in the path connecting  $x$  and  $y$ . To index the vertices on  $T$ , we first assign a vertex as the “root” and label it as  $O$ . A vertex is said to be on the  $n$ th level if the path linking it to the root has  $n$  edges. The root  $O$  is also said to be on the 0th level.

*Definition 1.1.* Let  $T$  be a tree with root  $O$ , and let  $\{N_n, n \geq 1\}$  be a sequence of positive integers.  $T$  is said to be a generalized Bethe tree or a generalized Cayley tree if each vertex on the  $n$ th level has  $N_{n+1}$  branches to the  $n + 1$ th level. For example, when  $N_1 = N + 1 \geq 2$  and  $N_n = N$  ( $n \geq 2$ ),  $T$  is rooted Bethe tree  $T_{B,N}$  on which each vertex has  $N + 1$  neighboring




Figure 1: Bethe tree  $T_{B,2}$ .

vertices (see Figure 1,  $T_{B,2}$ ), and when  $N_n = N \geq 1$  ( $n \geq 1$ ),  $T$  is rooted Cayley tree  $T_{C,N}$  on which each vertex has  $N$  branches to the next level.

In the following, we always assume that  $T$  is a generalized Bethe tree and denote by  $T^{(n)}$  the subgraph of  $T$  containing the vertices from level 0 (the root) to level  $n$ . We use  $(n, j)$  ( $1 \leq j \leq N_1 \cdots N_n, n \geq 1$ ) to denote the  $j$ th vertex at the  $n$ th level and denote by  $|B|$  the number of vertices in the subgraph  $B$ . It is easy to see that, for  $n \geq 1$ ,

$$|T^{(n)}| = \sum_{m=0}^n N_0 \cdots N_m = 1 + \sum_{m=1}^n N_1 \cdots N_m. \quad (1.1)$$

Let  $S = \{s_0, s_1, s_2, \dots\}$ ,  $\Omega = S^T$ ,  $\omega = \omega(\cdot) \in \Omega$ , where  $\omega(\cdot)$  is a function defined on  $T$  and taking values in  $S$ , and let  $F$  be the smallest Borel field containing all cylinder sets in  $\Omega$ . Let  $X = \{X_t, t \in T\}$  be the coordinate stochastic process defined on the measurable space  $(\Omega, F)$ ; that is, for any  $\omega = \{\omega(t), t \in T\}$ , define

$$X_t(\omega) = \omega(t), \quad t \in T. \quad (1.2)$$

$$X^{T^{(n)}} \triangleq \{X_t, t \in T^{(n)}\}, \quad \mu(X^{T^{(n)}} = x^{T^{(n)}}) = \mu(x^{T^{(n)}}). \quad (1.3)$$

Now we give a definition of Markov chain fields on the tree  $T$  by using the cylinder distribution directly, which is a natural extension of the classical definition of Markov chains (see [1]).

*Definition 1.2.* Let  $Q = Q(j | i)$ . One has a strictly positive stochastic matrix on  $S$ ,  $q = (q(s_0), q(s_1), q(s_2) \dots)$  a strictly positive distribution on  $S$ , and  $\mu_Q$  a measure on  $(\Omega, F)$ . If

$$\begin{aligned} \mu_Q(x_{0,1}) &= q(x_{0,1}), \\ \mu_Q(x^{T^{(n)}}) &= q(x_{0,1}) \prod_{m=0}^{n-1} \prod_{i=1}^{N_0 \cdots N_m} \prod_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} q(x_{m+1,j} | x_{m,i}), \quad n \geq 1. \end{aligned} \quad (1.4)$$

Then  $\mu_Q$  will be called a Markov chain field on the tree  $T$  determined by the stochastic matrix  $Q$  and the distribution  $q$ .

Let  $\mu$  be an arbitrary probability measure defined as (1.3), denote

$$f_n(\omega) = -\frac{1}{|T^{(n)}|} \log \mu(X^{T^{(n)}}). \tag{1.5}$$

$f_n(\omega)$  is called the entropy density on subgraph  $T^{(n)}$  with respect to  $\mu$ . If  $\mu = \mu_Q$ , then by (1.4), (1.5) we have

$$f_n(\omega) = -\frac{1}{|T^{(n)}|} \left[ \log q(X_{0,1}) + \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \log q(X_{m+1,j} | X_{m,i}) \right]. \tag{1.6}$$

The convergence of  $f_n(\omega)$  in a sense ( $L_1$  convergence, convergence in probability, or almost sure convergence) is called the Shannon-McMillan theorem or the entropy theorem or the asymptotic equipartition property (AEP) in information theory. The Shannon-McMillan theorem on the Markov chain has been studied extensively (see [2, 3]). In the recent years, with the development of the information theory scholars get to study the Shannon-McMillan theorems for the random field on the tree graph (see [4]). The tree models have recently drawn increasing interest from specialists in physics, probability and information theory. Berger and Ye (see [5]) have studied the existence of entropy rate for G-invariant random fields. Recently, Ye and Berger (see [6]) have also studied the ergodic property and Shannon-McMillan theorem for PPG-invariant random fields on trees. But their results only relate to the convergence in probability. Yang et al. [7–9] have recently studied a.s. convergence of Shannon-McMillan theorems, the limit properties and the asymptotic equipartition property for Markov chains indexed by a homogeneous tree and the Cayley tree, respectively. Shi and Yang (see [10]) have investigated some limit properties of random transition probability for second-order Markov chains indexed by a tree.

In this paper, we study a class of Shannon-McMillan random approximation theorems for arbitrary random fields on the generalized Bethe tree by comparison between the arbitrary measure and Markov measure on the generalized Bethe tree. As corollaries, a class of Shannon-McMillan theorems for arbitrary random fields and the Markov chains field on the generalized Bethe tree are obtained. Finally, some limit properties for the expectation of the random conditional entropy are discussed.

**Lemma 1.3.** *Let  $\mu_1$  and  $\mu_2$  be two probability measures on  $(\Omega, \mathcal{F})$ ,  $D \in \mathcal{F}$ , and let  $\{\tau_n, n \geq 0\}$  be a positive-valued stochastic sequence such that*

$$\liminf_n \frac{\tau_n}{|T^{(n)}|} > 0, \quad \mu_1\text{-a.s. } \omega \in D, \tag{1.7}$$

then

$$\limsup_{n \rightarrow \infty} \frac{1}{\tau_n} \log \frac{\mu_2(X^{T^{(n)}})}{\mu_1(X^{T^{(n)}})} \leq 0, \quad \mu_1\text{-a.s. } \omega \in D. \tag{1.8}$$

In particular, let  $\tau_n = |T^{(n)}|$ , then

$$\limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log \frac{\mu_2(X^{T^{(n)}})}{\mu_1(X^{T^{(n)}})} \leq 0, \quad \mu_1\text{-a.s. } \omega \in D. \quad (1.9)$$

*Proof* (see [11]). Let

$$\varphi(\mu | \mu_Q) = \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log \frac{\mu(X^{T^{(n)}})}{\mu_Q(X^{T^{(n)}})}. \quad (1.10)$$

$\varphi(\mu | \mu_Q)$  is called the sample relative entropy rate of  $\mu$  relative to  $\mu_Q$ .  $\varphi(\mu | \mu_Q)$  is also called the asymptotic logarithmic likelihood ratio. By (1.9)

$$\varphi(\mu | \mu_Q) \geq \liminf_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log \frac{\mu(X^{T^{(n)}})}{\mu_Q(X^{T^{(n)}})} \geq 0, \quad \mu\text{-a.s.} \quad (1.11)$$

Hence  $\varphi(\mu | \mu_Q)$  can be look on as a type of measures of the deviation between the arbitrary random fields and the Markov chain fields on the generalized Bethe tree.  $\square$

## 2. Main Results

**Theorem 2.1.** Let  $X = \{X_t, t \in T\}$  be an arbitrary random field on the generalized Bethe tree.  $f_n(\omega)$  and  $\varphi(\mu | \mu_Q)$  are, respectively, defined as (1.5) and (1.10). Denote  $\alpha \geq 0$ ,  $H_m^Q(X_{m+1,j} | X_{m,i})$  the random conditional entropy of  $X_{m+1,j}$  relative to  $X_{m,i}$  on the measure  $\mu_Q$ , that is,

$$H_m^Q(X_{m+1,j} | X_{m,i}) = - \sum_{x_{m+1,j} \in S} q(x_{m+1,j} | X_{m,i}) \log q(x_{m+1,j} | X_{m,i}). \quad (2.1)$$

Let

$$D(c) = \{\omega : \varphi(\mu | \mu_Q) \leq c\}, \quad (2.2)$$

$$b_\alpha = \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} E_Q \left[ \log^2 q(X_{m+1,j} | X_{m,i}) \cdot q(X_{m+1,j} | X_{m,i})^{-\alpha} | X_{m,i} \right] < \infty, \quad (2.3)$$

when  $0 \leq c \leq \alpha^2 b_\alpha / 2$ ,

$$\limsup_{n \rightarrow \infty} \left\{ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right\} \tag{2.4}$$

$$\leq \sqrt{2cb_\alpha}, \quad \mu\text{-a.s. } \omega \in D(c).$$

$$\liminf_{n \rightarrow \infty} \left\{ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right\} \tag{2.5}$$

$$\geq -\sqrt{2cb_\alpha} - c, \quad \mu\text{-a.s. } \omega \in D(c).$$

In particular,

$$\lim_{n \rightarrow \infty} \left[ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right] \tag{2.6}$$

$$= 0, \quad \mu\text{-a.s. } \omega \in D(0),$$

where  $\log$  is the natural logarithmic,  $E_Q$  is expectation with respect to the measure  $\mu_Q$ .

*Proof.* Let  $(\Omega, \mathcal{F}, \mu)$  be the probability space we consider,  $\lambda$  an arbitrary constant. Define

$$E_Q \left[ q(X_{m+1,j} | X_{m,i})^{-\lambda} \mid X_{m,i} = x_{m,i} \right] = \sum_{x_{m+1,j} \in S} q(x_{m+1,j} | x_{m,i})^{1-\lambda}; \tag{2.7}$$

denote

$$\mu_Q(\lambda, x^{T^{(n)}}) = q(x_{0,1}) \prod_{m=0}^{n-1} \prod_{i=1}^{N_0 \cdots N_m} \prod_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \frac{q(x_{m+1,j} | x_{m,i})^{1-\lambda}}{E_Q \left[ q(X_{m+1,j} | X_{m,i})^{-\lambda} \mid X_{m,i} = x_{m,i} \right]}. \tag{2.8}$$

We can obtain by (2.7), (2.8) that in the case  $n \geq 1$ ,

$$\begin{aligned}
 & \sum_{x^{L_n} \in S} \mu_Q(\lambda; x^{T^{(n)}}) \\
 &= \sum_{x^{L_n} \in S} q(x_{0,1}) \prod_{m=0}^{n-1} \prod_{i=1}^{N_0 \cdots N_m} \prod_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \frac{q(x_{m+1,j} | x_{m,i})^{1-\lambda}}{E_Q[q(X_{m+1,j} | X_{m,i})^{-\lambda} | X_{m,i} = x_{m,i}]} \\
 &= \mu_Q(\lambda; x^{T^{(n-1)}}) \sum_{x^{L_n} \in S} \prod_{i=1}^{N_0 \cdots N_{n-1}} \prod_{j=N_n(i-1)+1}^{N_n i} \frac{q(x_{n,j} | x_{n-1,i})^{1-\lambda}}{E_Q[q(X_{n,j} | X_{n-1,i})^{-\lambda} | X_{n-1,i} = x_{n-1,i}]} \\
 &= \mu_Q(\lambda; x^{T^{(n-1)}}) \prod_{i=1}^{N_0 \cdots N_{n-1}} \prod_{j=N_n(i-1)+1}^{N_n i} \sum_{x_{n,j} \in S} \frac{q(x_{n,j} | x_{n-1,i})^{1-\lambda}}{E_Q[q(x_{n,j} | x_{n-1,i})^{-\lambda} | X_{n-1,i} = x_{n-1,i}]} \\
 &= \mu_Q(\lambda; x^{T^{(n-1)}}) \prod_{i=1}^{N_0 \cdots N_{n-1}} \prod_{j=N_n(i-1)+1}^{N_n i} \frac{E_Q[q(X_{n,j} | X_{n-1,i})^{-\lambda} | X_{n-1,i} = x_{n-1,i}]}{E_Q[q(x_{n,j} | x_{n-1,i})^{-\lambda} | X_{n-1,i} = x_{n-1,i}]} \\
 &= \mu_Q(\lambda; x^{T^{(n-1)}}),
 \end{aligned} \tag{2.9}$$

$$\sum_{x^{L_0} \in S} \mu_Q(\lambda; x^{T^{(0)}}) = \sum_{x_{0,1} \in S} q(x_{0,1}) = 1. \tag{2.10}$$

Therefore,  $\mu_Q(\lambda, x^{T^{(n)}})$ ,  $n = 0, 1, 2, \dots$  are a class of consistent distributions on  $S^{T^{(n)}}$ . Let

$$U_n(\lambda, \omega) = \frac{\mu_Q(\lambda, X^{T^{(n)}})}{\mu(X^{T^{(n)}})}, \tag{2.11}$$

then  $\{U_n(\lambda, \omega), \mathcal{F}_n, n \geq 1\}$  is a nonnegative supermartingale which converges almost surely (see [12]). By Doob's martingale convergence theorem we have

$$\lim_{n \rightarrow \infty} U_n(\lambda, \omega) = U_\infty(\lambda, \omega) < \infty. \quad \mu\text{-a.s.} \tag{2.12}$$

Hence by (1.3), (1.9), (2.9), and (2.11) we get

$$\limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log U_n(\lambda, \omega) \leq 0. \quad \mu\text{-a.s.} \tag{2.13}$$

By (1.4), (2.8), and (2.11), we have

$$\begin{aligned} & \frac{1}{|T^{(n)}|} \log U_n(\lambda, \omega) \\ &= \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \left[ -\lambda \log q(X_{m+1,j} | X_{m,i}) - \log E_Q \left( q(X_{m+1,j} | X_{m,i})^{-\lambda} | X_{m,i} \right) \right] \\ & \quad + \frac{1}{|T^{(n)}|} \log \frac{\mu_Q(X^{T^{(n)}})}{\mu(X^{T^{(n)}})}. \end{aligned} \tag{2.14}$$

By (1.10), (2.2), (2.13), and (2.14) we have

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \left[ -\lambda \log q(X_{m+1,j} | X_{m,i}) - \log E_Q \left( q(X_{m+1,j} | X_{m,i})^{-\lambda} | X_{m,i} \right) \right] \\ & \leq \varphi(\mu | \mu_Q) \leq c, \quad \mu\text{-a.s. } \omega \in D(c). \end{aligned} \tag{2.15}$$

By (2.15) we have

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} (-\lambda) \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\ & \leq \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \left[ \log E_Q \left( q(X_{m+1,j} | X_{m,i})^{-\lambda} | X_{m,i} \right) \right. \\ & \quad \left. - E_Q(-\lambda \log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \right] + c, \quad \mu\text{-a.s. } \omega \in D(c). \end{aligned} \tag{2.16}$$

By the inequality

$$x^{-\lambda} - 1 + \lambda \log x \leq \left( \frac{1}{2} \right) \lambda^2 (\log x)^2 x^{-|\lambda|}, \quad 0 \leq x \leq 1, \tag{2.17}$$

$\log x \leq x - 1$  ( $x \geq 0$ ) and (2.16), (2.17), (2.3), we have in the case of  $|\lambda| < \alpha$ ,

$$\begin{aligned}
& \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} (-\lambda) \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\
& \leq \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \left[ E_Q \left( q(X_{m+1,j} | X_{m,i})^{-\lambda} | X_{m,i} \right) - 1 \right. \\
& \qquad \qquad \qquad \left. - E_Q \left( -\lambda \log q(X_{m+1,j} | X_{m,i}) | X_{m,i} \right) \right] + c \\
& \leq \limsup_{n \rightarrow \infty} \frac{1}{2|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} E_Q \left[ \lambda^2 \log^2(q(X_{m+1,j} | X_{m,i})) \right. \\
& \qquad \qquad \qquad \left. \cdot q(X_{m+1,j} | X_{m,i})^{-|\lambda|} | X_{m,i} \right] + c \\
& \leq \limsup_{n \rightarrow \infty} \frac{\lambda^2}{2|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} E_Q \left[ \log^2(q(X_{m+1,j} | X_{m,i})) \cdot q(X_{m+1,j} | X_{m,i})^{-\alpha} | X_{m,i} \right] \\
& \qquad \qquad \qquad + c = \left( \frac{1}{2} \right) \lambda^2 b_\alpha + c. \quad \mu\text{-a.s. } \omega \in D(c).
\end{aligned} \tag{2.18}$$

When  $0 < \lambda < \alpha$ , we get by (2.18)

$$\begin{aligned}
& \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} - \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\
& \leq \left( \frac{1}{2} \right) \lambda b_\alpha + \frac{c}{\lambda}, \quad \mu\text{-a.s. } \omega \in D(c).
\end{aligned} \tag{2.19}$$

Let  $g(\lambda) = (1/2)\lambda b_\alpha + c/\lambda$ , in the case  $0 < c \leq (\alpha^2 b_\alpha)/2$ , then it is obvious  $g(\lambda)$  attains, at  $\lambda = \sqrt{(2c)/b_\alpha}$ , its smallest value  $g(\sqrt{(2c)/b_\alpha}) = \sqrt{2cb_\alpha}$  on the interval  $(0, \alpha)$ . We have

$$\begin{aligned}
& \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} - \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\
& \leq \sqrt{2cb_\alpha}, \quad \mu\text{-a.s. } \omega \in D(c).
\end{aligned} \tag{2.20}$$

When  $c = 0$ , we select  $0 < \lambda_i < \alpha$  such that  $\lambda_i \rightarrow 0$  ( $i \rightarrow \infty$ ). Hence for all  $i$ , it follows from (2.19) that

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} - \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\ & \leq 0, \quad \mu\text{-a.s. } \omega \in D(0). \end{aligned} \tag{2.21}$$

It is easy to see that (2.20) also holds if  $c = 0$  from (2.21).

Analogously, when  $-\alpha < \lambda < 0$ , it follows from (2.18) if  $0 \leq c \leq (\alpha^2 b_\alpha)/2$ ,

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} - \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\ & \geq -\sqrt{2cb_\alpha}, \quad \mu\text{-a.s. } \omega \in D(c). \end{aligned} \tag{2.22}$$

Setting  $\lambda = 0$  in (2.14), by (2.14) we have

$$\limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log U_n(0, \omega) = \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log \frac{\mu_Q(X^{T^{(n)}})}{\mu(X^{T^{(n)}})} \leq 0, \quad \mu\text{-a.s.} \tag{2.23}$$

Noticing

$$H_m^Q(X_{m+1,j} | X_{m,i}) = E_Q[-\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}]. \tag{2.24}$$

By (1.4), (1.5), (2.20), and (2.23), we obtain

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \left[ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j}, X_{m,i}) \right] \\ & \leq \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log \frac{\mu_Q(X^{T^{(n)}})}{\mu(X^{T^{(n)}})} \\ & \quad + \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \\ & \quad - \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\ & \leq \sqrt{2cb_\alpha}, \quad \mu\text{-a.s. } \omega \in D(c). \end{aligned} \tag{2.25}$$

Hence (2.4) follows from (2.25). By (1.4), (1.5), (1.10), (2.2), and (2.22), we have

$$\begin{aligned}
 & \liminf_{n \rightarrow \infty} \left[ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(\omega) \right] \\
 & \geq \liminf_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log \left[ \frac{\mu_Q(X^{T^{(n)}})}{\mu(X^{T^{(n)}})} \right] \\
 & \quad + \liminf_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \\
 & \quad - \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\
 & \geq -\varphi(\mu | \mu_Q) - \sqrt{2cb_\alpha} \geq -\sqrt{2cb_\alpha} - c, \quad \mu\text{-a.s. } \omega \in D(c).
 \end{aligned} \tag{2.26}$$

Therefore (2.5) follows from (2.26). Set  $c = 0$  in (2.4) and (2.5), (2.6) holds naturally. □

**Corollary 2.2.** *Let  $X = \{X_t, t \in T\}$  be the Markov chains field determined by the measure  $\mu_Q$  on the generalized Bethe tree  $T$ .  $f_n(\omega)$ ,  $b_\alpha$  are, respectively, defined as (1.6) and (2.3), and  $H_m^Q(X_{m+1,j} | X_{m,i})$  is defined by (2.1). Then*

$$\lim_{n \rightarrow \infty} \left\{ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right\} = 0. \quad \mu_Q\text{-a.s.} \tag{2.27}$$

*Proof.* We take  $\mu \equiv \mu_Q$ , then  $\varphi(\mu | \mu_Q) \equiv 0$ . It implies that (2.2) always holds when  $c = 0$ . Therefore  $D(0) = \Omega$  holds. Equation(2.27) follows from (2.3) and (2.6). □

### 3. Some Shannon-McMillan Approximation Theorems on the Finite State Space

**Corollary 3.1.** *Let  $X = \{X_t, t \in T\}$  be an arbitrary random field which takes values in the alphabet  $S = \{s_1, \dots, s_N\}$  on the generalized Bethe tree.  $f_n(\omega)$ ,  $\varphi(\mu | \mu_Q)$  and  $D(c)$  are defined as (1.5), (1.10), and (2.2). Denote  $0 \leq \alpha < 1$ ,  $0 \leq c \leq 2N\alpha^2 / [(1 - \alpha)e]^2$ .  $H_m^Q(X_{m+1,j} | X_{m,i})$  is defined as above. Then*

$$\begin{aligned}
 & \limsup_{n \rightarrow \infty} \left[ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right] \\
 & \leq \frac{2e^{-1}}{(1 - \alpha)} \sqrt{2cN}, \quad \mu\text{-a.s. } \omega \in D(c),
 \end{aligned} \tag{3.1}$$

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \left[ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right] \\ & \geq -\frac{2e^{-1}}{1-\alpha} \sqrt{2cN} - c, \quad \mu\text{-a.s. } \omega \in D(c). \end{aligned} \tag{3.2}$$

*Proof.* Set  $0 < \alpha < 1$  we consider the function

$$\phi(x) = (\log x)^2 x^{1-\alpha}, \quad 0 < x \leq 1, \quad 0 < \alpha < 1 \quad (\text{Set } \phi(0) = 0). \tag{3.3}$$

Then

$$\phi'(x) = x^{-\alpha} \left[ 2(\log x) + (\log x)^2 (1 - \alpha) \right]. \tag{3.4}$$

Let  $\phi'(x) = 0$  thus  $x = e^{2/(\alpha-1)}$ . Accordingly it can be obtained that

$$\max\{\phi(x), 0 \leq x \leq 1\} = \phi\left(e^{2/(\alpha-1)}\right) = \left(\frac{2}{\alpha-1}\right)^2 e^{-2}. \tag{3.5}$$

By (2.3) and (3.5) we have

$$\begin{aligned} b_\alpha &= \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} E_Q \left[ \log^2 q(X_{m+1,j} | X_{m,i}) \cdot q(X_{m+1,j} | X_{m,i})^{-\alpha} | X_{m,i} \right] \\ &= \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \sum_{x_{m+1,j} \in S} \log^2 q(x_{m+1,j} | X_{m,i}) \cdot q(x_{m+1,j} | X_{m,i})^{1-\alpha} \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \sum_{x_{m+1,j} \in S_1} \left(\frac{2}{\alpha-1}\right)^2 e^{-2} \\ &= N \left(\frac{2}{\alpha-1}\right)^2 e^{-2} \cdot \limsup_{n \rightarrow \infty} \frac{|T^{(n)}| - 1}{|T^{(n)}|} = N \left(\frac{2}{\alpha-1}\right)^2 e^{-2} < \infty. \end{aligned} \tag{3.6}$$

Therefore, (2.3) holds naturally. By (2.18) and (3.6) we have

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} (-\lambda) \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\ & \leq N\lambda^2 \frac{2e^{-2}}{(\alpha-1)^2} + c, \quad \mu\text{-a.s. } \omega \in D(c). \end{aligned} \tag{3.7}$$

In the case of  $0 < \lambda < \alpha$ , by (3.7) we have

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} - \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\ & \leq N\lambda \frac{2e^{-2}}{(\alpha-1)^2} + \frac{c}{\lambda}, \quad \mu\text{-a.s. } \omega \in D(c). \end{aligned} \quad (3.8)$$

Let  $g(\lambda) = 2\lambda N e^{-2}/(\alpha-1)^2 + c/\lambda$ , in the case  $0 < c \leq 2N\alpha^2/[(1-\alpha)e]^2$ , then it is obvious  $g(\lambda)$  attains, at  $\lambda = (1-\alpha)e\sqrt{c/2N}$ , its smallest value  $g((1-\alpha)e\sqrt{c/2N}) = 2e^{-1}\sqrt{2cN}/(1-\alpha)$  on the interval  $(0, \alpha)$ . That is

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} - \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\ & \leq \frac{2e^{-1}}{(1-\alpha)} \sqrt{2cN}, \quad \mu\text{-a.s. } \omega \in D(c). \end{aligned} \quad (3.9)$$

By the similar means of reasoning (2.21), it can be concluded that (3.9) also holds when  $c = 0$ . According to the methods of proving (2.4), (3.1) follows from (1.5), (2.23), and (3.9). Similarly, when  $-\alpha < \lambda < 0$ ,  $0 \leq c \leq 2N\alpha^2/[(1-\alpha)e]^2$ , by (3.7) we have

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} - \{ \log q(X_{m+1,j} | X_{m,i}) - E_Q(\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}) \} \\ & \geq -\frac{2e^{-1}}{(1-\alpha)} \sqrt{2cN}, \quad \mu\text{-a.s. } \omega \in D(c). \end{aligned} \quad (3.10)$$

Imitating the proof of (2.5), (3.2) follows from (1.5), (1.10), (2.2), and (3.10).  $\square$

**Corollary 3.2** (see [9]). *Let  $X = \{X_t, t \in T\}$  be the Markov chains field determined by the measure  $\mu_Q$  on the generalized Bethe tree  $T$ .  $f_n(\omega)$  is defined as (1.6), and  $H_m^Q(X_{m+1,j} | X_{m,i})$  is defined as (2.1). Then*

$$\lim_{n \rightarrow \infty} \left\{ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right\} = 0. \quad \mu_Q\text{-a.s.} \quad (3.11)$$

*Proof.* By (3.1) and (3.2) in Corollary 3.1, we obtain that when  $c = 0$ ,

$$\lim_{n \rightarrow \infty} \left\{ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right\} = 0, \quad \mu\text{-a.s. } \omega \in D(0). \tag{3.12}$$

Set  $\mu \equiv \mu_Q$ , then  $\varphi(\mu | \mu_Q) \equiv 0$ . It implies (2.2) always holds when  $c = 0$ . Therefore  $D(0) = \Omega$  holds. Equation (3.11) follows from (3.12). □

**Corollary 3.3.** *Under the assumption of Corollary 3.1, if  $\mu \ll \mu_Q$ , then*

$$\lim_{n \rightarrow \infty} \left\{ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right\} = 0. \quad \mu\text{-a.s.} \tag{3.13}$$

*Proof.* It can be obtained that  $\varphi(\mu | \mu_Q) \equiv 0, \mu$  a.s. holds if  $\mu \ll \mu_Q$  (see Gray 1990 [13]), therefore  $\mu(D(0)) = 1$ . Equation (3.13) follows from (3.12).

Let  $X = \{X_t, t \in T\}$  be a Markov chains field on the generalized Bethe tree with the initial distribution and the joint distribution with respect to the measure  $\mu_P$  as follows:

$$\mu_P(x_{0,1}) = p(x_{0,1}), \tag{3.14}$$

$$\mu_P(x^{T^{(n)}}) = p(x_{0,1}) \prod_{m=0}^{n-1} \prod_{i=1}^{N_0 \cdots N_m} \prod_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} p(x_{m+1,j} | x_{m,i}), \quad n \geq 1, \tag{3.15}$$

where  $P = p(j | i)$  is a strictly positive stochastic matrix on  $S$ ,  $p = (p(s_0), p(s_1), p(s_2) \dots)$  is a strictly positive distribution. Therefore, the entropy density of  $X = \{X_t, t \in T\}$  with respect to the measure  $\mu_P$  is

$$f_n(\omega) = -\frac{1}{|T^{(n)}|} \left[ \log p(X_{0,1}) + \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \log p(X_{m+1,j} | X_{m,i}) \right]. \tag{3.16}$$

Let the initial distribution and joint distribution of  $X = \{X_t, t \in T\}$  with respect to the measure  $\mu_Q$  be defined as (1.4) and (1.5), respectively. □

We have the following conclusion.

**Corollary 3.4.** *Let  $X = \{X_t, t \in T\}$  be a Markov chains field on the generalized Bethe tree  $T$  whose initial distribution and joint distribution with respect to the measure  $\mu_P$  and  $\mu_Q$  are defined by (3.14), (3.15) and (1.4), (1.5), respectively.  $f_n(\omega)$  is defined as (3.16). If*

$$\sum_{h \in S} \sum_{l \in S} \frac{[p(l | h) - q(l | h)]^+}{q(l | h)} \leq c, \tag{3.17}$$

then

$$\begin{aligned} \limsup_{n \rightarrow \infty} \left[ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right] \\ \leq \frac{2e^{-1}}{1-\alpha} \sqrt{2cN}, \quad \mu_P\text{-a.s.} \end{aligned} \quad (3.18)$$

$$\begin{aligned} \liminf_{n \rightarrow \infty} \left[ f_n(\omega) - \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} H_m^Q(X_{m+1,j} | X_{m,i}) \right] \\ \geq -\frac{2e^{-1}}{1-\alpha} \sqrt{2cN} - c. \quad \mu_P\text{-a.s.} \end{aligned} \quad (3.19)$$

*Proof.* Let  $\mu = \mu_P$  in Corollary 3.1, and by (1.5), (3.15) we get (3.16). By the inequalities  $\log x \leq x - 1$  ( $x > 0$ ),  $a \leq [a]^+$ , (3.17), and (1.10), we obtain

$$\begin{aligned} & \varphi(\mu_P | \mu_Q) \\ &= \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log \frac{\mu_P(X^{T^{(n)}})}{\mu_Q(X^{T^{(n)}})} \\ &= \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log \frac{p(X_{0,1}) \prod_{m=0}^{n-1} \prod_{i=1}^{N_0 \cdots N_m} \prod_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} p(X_{m+1,j} | X_{m,i})}{q(X_{0,1}) \prod_{m=0}^{n-1} \prod_{i=1}^{N_0 \cdots N_m} \prod_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} q(X_{m+1,j} | X_{m,i})} \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \log \frac{p(X_{0,1})}{q(X_{0,1})} + \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \log \frac{p(X_{m+1,j} | X_{m,i})}{q(X_{m+1,j} | X_{m,i})} \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \sum_{h \in S} \sum_{l \in S} \delta_h(X_{m,i}) \delta_l(X_{m+1,j}) \log \frac{p(l|h)}{q(l|h)} \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{|T^{(n)}|} \sum_{m=0}^{n-1} \sum_{i=1}^{N_0 \cdots N_m} \sum_{j=N_{m+1}(i-1)+1}^{N_{m+1}i} \sum_{h \in S} \sum_{l \in S} \delta_h(X_{m,i}) \delta_l(X_{m+1,j}) \left[ \frac{p(l|h)}{q(l|h)} - 1 \right] \\ &\leq \sum_{h \in S} \sum_{l \in S} \limsup_{n \rightarrow \infty} \frac{|T^{(n)}| - 1}{|T^{(n)}|} \frac{p(l|h) - q(l|h)}{q(l|h)} \\ &\leq \sum_{h \in S} \sum_{l \in S} \frac{[p(l|h) - q(l|h)]^+}{q(l|h)}. \end{aligned} \quad (3.20)$$

By (3.17) and (3.20) we have

$$\varphi(\mu_P | \mu_Q) \leq c, \quad \text{a.s.} \quad (3.21)$$

It follows from (2.2) and (3.21) that  $D(c) = \Omega$ ; therefore (3.18), (3.19) follow from (3.1), (3.2).  $\square$

### 4. Some Limit Properties for Expectation of Random Conditional Entropy on the Finite State Space

**Lemma 4.1** (see [8]). Let  $X^{T^{(n)}} = \{X_t, t \in T^{(n)}\}$  be a Markov chains field defined on a Bethe tree  $T_{B,N}$ ,  $S_n(k, \omega)$  be the number of  $k$  in the set of random variables  $X^{T^{(n)}} = \{X_t, t \in T^{(n)}\}$ . then for all  $k \in S$ ,

$$\lim_n \frac{S_n(k, \omega)}{|T^{(n)}|} = \pi(k) \quad \mu_Q\text{-a.s.} \tag{4.1}$$

where  $\pi = (\pi(1), \dots, \pi(N))$  is the stationary distribution determined by  $Q$ .

**Theorem 4.2.** Let  $X^{T^{(n)}} = \{X_t, t \in T^{(n)}\}$  be a Markov chains field defined on a Bethe tree  $T_{B,N}$ , and let  $H_m^Q(X_{m+1,j} | X_{m,i})$  be defined as above. Then

$$\begin{aligned} \lim_n \frac{1}{|T^{(n)}|} & \left[ \sum_{i=1}^{N+1} H_0^Q(X_{1,i} | X_{0,1}) + \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} H_m^Q(X_{m+1,j} | X_{m,i}) \right] \\ & = - \sum_{k \in S} \sum_{l \in S} \pi(k) q(l | k) \log q(l | k), \quad \mu_Q\text{-a.s.} \end{aligned} \tag{4.2}$$

*Proof.* Noticing now  $N_1 = N + 1$ , for all  $n \geq 2$ ,  $N_n = N$ , that therefore we have

$$\begin{aligned} & \sum_{i=1}^{N+1} H_0^Q(X_{1,i} | X_{0,1}) + \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} H_m^Q(X_{m+1,j} | X_{m,i}) \\ & = \sum_{i=1}^{N+1} - E_Q [\log q(X_{1,i} | X_{0,1}) | X_{0,1}] \\ & \quad + \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} - E_Q [\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}] \\ & = - \sum_{i=1}^{N+1} \sum_{x_{1,i} \in S} q(x_{1,i} | X_{0,1}) \log q(x_{1,i} | X_{0,1}) \\ & \quad - \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} \sum_{x_{m+1,j} \in S} q(x_{m+1,j} | X_{m,i}) \log q(x_{m+1,j} | X_{m,i}) \end{aligned}$$

$$\begin{aligned}
 &= - \sum_{i=1}^{N+1} \sum_{k \in S} \sum_{l \in S} \delta_k(X_{0,1}) q(l | k) \log q(l | k) \\
 &\quad - \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} \sum_{k \in S} \sum_{l \in S} \delta_k(X_{m,i}) q(l | k) \log q(l | k) \\
 &= - \sum_{k \in S} \sum_{l \in S} q(l | k) \log q(l | k) \left[ \sum_{i=1}^{N+1} \delta_k(X_{0,1}) + \sum_{m=0}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} \delta_k(X_{m,i}) \right] \\
 &= - \sum_{k \in S} \sum_{l \in S} q(l | k) \log q(l | k) [NS_{n-1}(k, \omega) + \delta_k(X_{0,1})].
 \end{aligned} \tag{4.3}$$

Noticing that  $\lim_{n \rightarrow \infty} (|T^{(n)}|/|T^{(n-1)}|) = N$ , by (4.3) we have

$$\begin{aligned}
 &\lim_n \frac{1}{|T^{(n)}|} \left[ \sum_{i=1}^{N+1} H_0^Q(X_{1,i} | X_{0,1}) + \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} H_m^Q(X_{m+1,j} | X_{m,i}) \right] \\
 &= - \lim_n \frac{1}{|T^{(n)}|} \sum_{k \in S} \sum_{l \in S} q(l | k) \log q(l | k) [NS_{n-1}(k, \omega) + \delta_k(X_{0,1})] \\
 &= - \sum_{k \in S} \sum_{l \in S} q(l | k) \log q(l | k) \lim_n \frac{1}{|T^{(n-1)}|} S_{n-1}(k, \omega) \\
 &= - \sum_{k \in S} \sum_{l \in S} \pi(k) q(l | k) \log q(l | k).
 \end{aligned} \tag{4.4}$$

Equation(4.2) follows from (4.4). □

**Theorem 4.3.** Let  $X^{T^{(n)}} = \{X_t, t \in T^{(n)}\}$  be a Markov chains field defined on a Bethe tree  $T_{B,N}$ ,  $H_m^Q(X_{m+1,j} | X_{m,i})$  defined as above. Then

$$\begin{aligned}
 &\lim_n \frac{1}{|T^{(n)}|} \left\{ \sum_{i=1}^{N+1} E_Q \left[ H_0^Q(X_{1,i} | X_{0,1}) \right] + \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} E_Q \left[ H_m^Q(X_{m+1,j} | X_{m,i}) \right] \right\} \\
 &= - \sum_{k \in S} \sum_{l \in S} q(k) q(l | k) \log q(l | k). \quad \mu_Q\text{-a.s.}
 \end{aligned} \tag{4.5}$$

*Proof.* By the definition of  $H_m^Q(X_{m+1,j} | X_{m,i})$  and properties of conditional expectation, we have

$$\begin{aligned}
 & \sum_{i=1}^{N+1} E_Q \left[ H_0^Q(X_{1,i} | X_{0,1}) \right] + \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} E_Q \left[ H_m^Q(X_{m+1,j} | X_{m,i}) \right] \\
 &= \sum_{i=1}^{N+1} - E_Q \{ E_Q [\log q(X_{1,i} | X_{0,1}) | X_{0,1}] \} \\
 & \quad + \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} - E_Q \{ E_Q [\log q(X_{m+1,j} | X_{m,i}) | X_{m,i}] \} \\
 &= \sum_{i=1}^{N+1} - E_Q [\log q(X_{1,i} | X_{0,1})] \\
 & \quad + \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} - E_Q [\log q(X_{m+1,j} | X_{m,i})] \\
 &= - \sum_{i=1}^{N+1} \sum_{x_{0,1} \in S} \sum_{x_{1,i} \in S} q(x_{0,1}, x_{1,i}) \log q(x_{1,i} | x_{0,1}) \\
 & \quad - \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} \sum_{x_{m,i} \in S} \sum_{x_{m+1,j} \in S} q(x_{m,i}, x_{m+1,j}) \log q(x_{m+1,j} | x_{m,i}) \\
 &= - \sum_{i=1}^{N+1} \sum_{k \in S} \sum_{l \in S} q(k) q(l | k) \log q(l | k) \\
 & \quad - \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} \sum_{k \in S} \sum_{l \in S} q(k) q(l | k) \log q(l | k) \\
 &= - \sum_{k \in S} \sum_{l \in S} q(k) q(l | k) \log q(l | k) \left( |T^{(n)}| - 1 \right).
 \end{aligned} \tag{4.6}$$

Accordingly we have by (4.6)

$$\begin{aligned}
 & \lim_n \frac{1}{|T^{(n)}|} \left\{ \sum_{i=1}^{N+1} E_Q \left[ H_0^Q(X_{1,i} | X_{0,1}) \right] + \sum_{m=1}^{n-1} \sum_{i=1}^{(N+1)N^{m-1}} \sum_{j=N(i-1)+1}^{Ni} E_Q \left[ H_m^Q(X_{m+1,j} | X_{m,i}) \right] \right\} \\
 &= - \sum_{k \in S} \sum_{l \in S} q(k) q(l | k) \log q(l | k) \lim_n \frac{|T^{(n)}| - 1}{|T^{(n)}|} \\
 &= - \sum_{k \in S} \sum_{l \in S} q(k) q(l | k) \log q(l | k), \quad \mu_Q\text{-a.s.}
 \end{aligned} \tag{4.7}$$

Therefore (4.5) also holds. □

## Acknowledgments

The work is supported by the Project of Higher Schools' Natural Science Basic Research of Jiangsu Province of China (09KJD110002). The author would like to thank the referee for his valuable suggestions. Correspondence author: K. Wang, main research interest is strong limit theory in probability theory. D. Zong main research interest is intelligent algorithm.

## References

- [1] K. L. Chung, *A Course in Probability Theory*, Academic Press, New York, NY, USA, 2nd edition, 1974.
- [2] L. Wen and Y. Weiguo, "An extension of Shannon-McMillan theorem and some limit properties for nonhomogeneous Markov chains," *Stochastic Processes and their Applications*, vol. 61, no. 1, pp. 129–145, 1996.
- [3] W. Liu and W. Yang, "Some extensions of Shannon-McMillan theorem," *Journal of Combinatorics, Information & System Sciences*, vol. 21, no. 3-4, pp. 211–223, 1996.
- [4] Z. Ye and T. Berger, *Information Measures for Discrete Random Fields*, Science Press, Beijing, China, 1998.
- [5] T. Berger and Z. X. Ye, "Entropic aspects of random fields on trees," *IEEE Transactions on Information Theory*, vol. 36, no. 5, pp. 1006–1018, 1990.
- [6] Z. Ye and T. Berger, "Ergodicity, regularity and asymptotic equipartition property of random fields on trees," *Journal of Combinatorics, Information & System Sciences*, vol. 21, no. 2, pp. 157–184, 1996.
- [7] W. Yang, "Some limit properties for Markov chains indexed by a homogeneous tree," *Statistics & Probability Letters*, vol. 65, no. 3, pp. 241–250, 2003.
- [8] W. Yang and W. Liu, "Strong law of large numbers and Shannon-McMillan theorem for Markov chain fields on trees," *IEEE Transactions on Information Theory*, vol. 48, no. 1, pp. 313–318, 2002.
- [9] W. Yang and Z. Ye, "The asymptotic equipartition property for nonhomogeneous Markov chains indexed by a homogeneous tree," *IEEE Transactions on Information Theory*, vol. 53, no. 9, pp. 3275–3280, 2007.
- [10] Z. Shi and W. Yang, "Some limit properties of random transition probability for second-order nonhomogeneous markov chains indexed by a tree," *Journal of Inequalities and Applications*, vol. 2009, Article ID 503203, 2009.
- [11] W. Liu and W. Yang, "Some strong limit theorems for Markov chain fields on trees," *Probability in the Engineering and Informational Sciences*, vol. 18, no. 3, pp. 411–422, 2004.
- [12] J. L. Doob, *Stochastic Processes*, John Wiley & Sons, New York, NY, USA, 1953.
- [13] R. M. Gray, *Entropy and Information Theory*, Springer, New York, NY, USA, 1990.