

# A REFINEMENT OF THE POINCARÉ INEQUALITY FOR KOLMOGOROV OPERATORS ON $\mathbb{R}^d$

YASUHIRO FUJITA

Received 28 June 2004

We give a refinement of the Poincaré inequality for Kolmogorov operators on  $\mathbb{R}^d$ . This refinement yields some regularity result of the corresponding semigroups.

## 1. Introduction

Let  $\{P_t\}$  be the semigroup on  $B_b(\mathbb{R}^d)$  associated with the Kolmogorov operator

$$L_0 = \frac{1}{2}\Delta + F(x) \cdot D. \quad (1.1)$$

Here we denote by  $B_b(\mathbb{R}^d)$  the Banach space of all Borel and bounded functions, endowed with the supremum norm. We assume a suitable dissipative assumption on the function  $F = (F_1, \dots, F_d)$  such that there exists a unique invariant probability measure  $\nu$  on  $\mathbb{R}^d$  associated with  $\{P_t\}$ . Let  $H^1(\nu)$  and  $H^2(\nu)$  be the Sobolev spaces with the norms

$$\|\varphi\|_{H^1(\nu)} = \left[ \int_{\mathbb{R}^d} [|\varphi|^2 + |D\varphi|^2] d\nu \right]^{1/2}, \quad (1.2)$$

$$\|\varphi\|_{H^2(\nu)} = \left[ \int_{\mathbb{R}^d} [|\varphi|^2 + |D\varphi|^2 + |D^2\varphi|^2] d\nu \right]^{1/2}, \quad (1.3)$$

respectively. It is well known that the Poincaré inequality with respect to  $\nu$  is the following:

$$\int_{\mathbb{R}^d} (\varphi - \bar{\varphi})^2 d\nu \leq \frac{1}{2\alpha} \int_{\mathbb{R}^d} |D\varphi|^2 d\nu, \quad \varphi \in H^1(\nu), \quad (1.4)$$

where  $\alpha > 0$  is a constant determined by  $F$ , and  $\bar{\varphi} = \int_{\mathbb{R}^d} \varphi d\nu$ . The Poincaré inequality (1.4) is so important that it implies existence of a *spectral gap* or, equivalently, *exponential*

convergence of equilibrium of the semigroup  $\{P_t\}$  such that

$$\int_{\mathbb{R}^d} |P_t\varphi - \bar{\varphi}|^2 d\nu \leq e^{-2\alpha t} \int_{\mathbb{R}^d} |\varphi|^2 d\nu, \quad t \geq 0, \varphi \in L^2(\nu) \quad (1.5)$$

(cf. [2, Proposition 3.12]).

The aim of this paper is to give a refinement of the Poincaré inequality (1.4) such that

$$\int_{\mathbb{R}^d} (\varphi - \bar{\varphi})^2 d\nu + \frac{1}{2\alpha} \int_0^\infty dt \int_{\mathbb{R}^d} |D^2 P_t \varphi|^2 d\nu \leq \frac{1}{2\alpha} \int_{\mathbb{R}^d} |D\varphi|^2 d\nu, \quad \varphi \in H^1(\nu). \quad (1.6)$$

When  $F(x) = -\alpha x$  in (1.1) (i.e.,  $\{P_t\}$  is the Ornstein-Uhlenbeck semigroup), inequality (1.6) is reduced to an equality. Furthermore, we will show that inequality (1.6) yields the regularity result such that  $P_t\varphi - \bar{\varphi} \in L^2((0, \infty), dt; H^2(\nu))$  for  $\varphi \in H^1(\nu)$ . This regularity result corresponds to the well-known regularity result such that  $P_t\varphi - \bar{\varphi} \in L^2((0, \infty), dt; H^1(\nu))$  for  $\varphi \in L^2(\nu)$  (cf. (1.5) and (3.18)).

In the proof of the Poincaré inequality (1.4), the following inequality was used for  $\varphi \in C_b^1(\mathbb{R}^d)$ :

$$|DP_t\varphi(x)|^2 \leq e^{-2\alpha t} P_t(|D\varphi|^2)(x), \quad (t, x) \in (0, \infty) \times \mathbb{R}^d \quad (1.7)$$

(cf. [2, Proposition 2.8]). In our proof of inequality (1.6), we will also use (1.7). However, we will derive another differential inequality so as not to lose the term  $|D^2 P_t\varphi(x)|^2$ . For this purpose, it is crucial to assume that the Kolmogorov operator  $L_0$  has the form of (1.1). It seems hard for the author to apply our proof directly to a more general Kolmogorov operator such as  $(1/2)\text{tr}[C(x)D^2] + F(x) \cdot D$ .

The contents of this paper are as follows. In Section 2, we will state the main results. They will be proved in Section 3.

## 2. Main results

First of all, we recall the results about invariant probability measures on  $\mathbb{R}^d$  (for details, see [2]). Following [1, Hypothesis 1.1], we make the following assumptions on  $F = (F_1, \dots, F_d)$  of (1.1).

(A)  $F \in C^4(\mathbb{R}^d; \mathbb{R}^d)$ , and there exist

$$m \geq 0 \text{ such that } \sup_{x \in \mathbb{R}^d} \frac{|D^\beta F(x)|}{1 + |x|^{2m+1-\beta}} < +\infty, \quad \beta = 0, 1, 2, 3, 4, \quad (2.1)$$

$$\alpha > 0 \text{ such that } DF(x)y \cdot y \leq -\alpha|y|^2, \quad x, y \in \mathbb{R}^d,$$

$$a, \gamma, c > 0 \text{ such that } (F(x+y) - F(x)) \cdot y \leq -a|y|^{2m+2} + c(|x|^\gamma + 1), \quad x, y \in \mathbb{R}^d.$$

By [1, Proposition 1.2.2], the stochastic differential equation

$$d\xi(t, x) = F(\xi(t, x))dt + dw(t), \quad \xi(0, x) = x, \quad (2.2)$$

admits a unique strong solution  $(\xi(t, x))$ , where  $(w(t))$  is a  $d$ -dimensional standard Brownian motion on a probability space. Then we can define the semigroup  $\{P_t\}$  on  $B_b(\mathbb{R}^d)$  by

$$P_t\varphi(x) = \mathbb{E}[\varphi(\xi(t, x))]. \tag{2.3}$$

By [2, Proposition 2.7], there exists a unique probability measure  $\nu$  on  $\mathbb{R}^d$  satisfying the following: for any uniformly continuous and bounded function  $\chi$  on  $\mathbb{R}^d$ , we have

$$\int_{\mathbb{R}^d} P_t\chi d\nu = \int_{\mathbb{R}^d} \chi d\nu, \quad t \geq 0. \tag{2.4}$$

Such a probability measure  $\nu$  on  $\mathbb{R}^d$  is called the invariant probability measure for  $\{P_t\}$ . Using this invariant probability measure  $\nu$ , we can extend  $\{P_t\}$  to a strongly continuous semigroup of contractions on  $L^p(\nu)$  for every  $p \geq 1$ . We also denote by  $\{P_t\}$  this extended strongly continuous semigroup. The generator  $(L, \text{dom}_p(L))$  of  $\{P_t\}$  in  $L^p(\nu)$  is the closure of the Kolmogorov operator  $(L_0, C_0^\infty(\mathbb{R}^d))$ , where  $L_0$  is the operator defined by (1.1), and  $C_0^\infty(\mathbb{R}^d)$  is the space of  $C^\infty$ -functions with compact supports. An important example of  $L$  is the Ornstein-Uhlenbeck operator corresponding to the case  $F(x) = -\alpha x$ .

Next, we define the Sobolev spaces  $H^1(\nu)$  and  $H^2(\nu)$ . The operators  $(D, C_0^\infty(\mathbb{R}^d))$  and  $(D^2, C_0^\infty(\mathbb{R}^d))$  are closable in  $L^p(\nu)$  for every  $p \geq 1$ . We also denote their closures by  $(D, \text{dom}_p(D))$  and  $(D^2, \text{dom}_p(D^2))$ , respectively. Then, we can define the Sobolev spaces  $H^1(\nu)$  and  $H^2(\nu)$  by  $H^1(\nu) = \text{dom}_2(D)$  and  $H^2(\nu) = \text{dom}_2(D^2)$ , respectively. They become Hilbert spaces with the norms defined by (1.2) and (1.3), respectively. Then, the Poincaré inequality (1.4) holds for the constant  $\alpha$  of (2.1).

Now, we state the main results of this paper.

**THEOREM 2.1.** *Assume (2.1). Then, for every  $\varphi \in H^1(\nu)$ ,*

$$P_t\varphi \in H^2(\nu), \quad t\text{-a.e. on } (0, \infty), \tag{2.5}$$

$$P_t\varphi - \bar{\varphi} \in L^2((0, \infty), dt; H^2(\nu)), \tag{2.6}$$

$$\int_{\mathbb{R}^d} (\varphi - \bar{\varphi})^2 d\nu + \frac{1}{2\alpha} \int_0^\infty dt \int_{\mathbb{R}^d} |D^2 P_t\varphi|^2 d\nu \leq \frac{1}{2\alpha} \int_{\mathbb{R}^d} |D\varphi|^2 d\nu. \tag{2.7}$$

$$\text{When } F(x) = -\alpha x, \text{ inequality (2.7) is reduced to an equality.} \tag{2.8}$$

Results (2.5) and (2.6) give a regularity result of  $P_t\varphi$  for  $\varphi \in H^1(\nu)$ . On the other hand, results (2.7) and (2.8) give refinements of the Poincaré inequality.

### 3. Proof of Theorem 2.1

In this section, we prove Theorem 2.1. For  $\varphi \in C_0^\infty(\mathbb{R}^d)$ , we set

$$\eta(t, x) = P_t\varphi(x), \quad (t, x) \in [0, \infty) \times \mathbb{R}^d. \tag{3.1}$$

First, we give two lemmas.

LEMMA 3.1. Assume (2.1). If  $\varphi \in C_0^\infty(\mathbb{R}^d)$ , then

$$(D\eta)_t, D^\beta \eta \quad (\beta = 0, 1, 2, 3) \text{ are continuous on } [0, \infty) \times \mathbb{R}^d, \quad (3.2)$$

$$(D\eta)_t = D\eta_t \quad \text{on } [0, \infty) \times \mathbb{R}^d. \quad (3.3)$$

*Proof.* Since  $F \in C^4(\mathbb{R}^d; \mathbb{R}^d)$  and  $\varphi \in C_0^\infty(\mathbb{R}^d)$ , it follows from the theory in [1, Chapter 1] that

$$D^\beta \eta \text{ is continuous on } [0, \infty) \times \mathbb{R}^d \quad \text{for } \beta = 0, 1, 2, 3. \quad (3.4)$$

Since  $\eta$  of (3.1) satisfies the Kolmogorov equation

$$\eta_t = \frac{1}{2} \Delta \eta + F \cdot D\eta \quad \text{on } [0, \infty) \times \mathbb{R}^d, \quad (3.5)$$

we have, for any  $R, T > 0$ ,

$$D\eta(t+h, x) - D\eta(t, x) = \int_t^{t+h} DL\eta(s, x) ds, \quad 0 \leq t \leq T, |x| < R, \quad (3.6)$$

where  $h \in \mathbb{R}$  is chosen such that  $t+h \geq 0$ . By (3.4) and (3.6), we conclude that  $D\eta(t, x)$  is differentiable with respect to  $t$  for  $|x| < R$  and

$$(D\eta)_t(t, x) = DL\eta(t, x) = D\eta_t(t, x), \quad 0 \leq t \leq T, |x| < R. \quad (3.7)$$

Since  $R, T > 0$  are arbitrary, (3.3) follows. By (3.4) and (3.7),  $(D\eta)_t$  is continuous on  $[0, \infty) \times \mathbb{R}^d$ . The proof is complete.  $\square$

LEMMA 3.2. Assume that (2.1) holds and  $\varphi \in C_0^\infty(\mathbb{R}^d)$ . Let

$$\chi(t, x) = |D\eta(t, x)|^2 = \sum_{j=1}^d |D_j \eta(t, x)|^2, \quad (t, x) \in [0, \infty) \times \mathbb{R}^d. \quad (3.8)$$

Then,

$$\chi_t, D^\beta \chi \quad (\beta = 0, 1, 2) \text{ are continuous on } [0, \infty) \times \mathbb{R}^d, \quad (3.9)$$

$$|D^2 \eta|^2 + \chi_t \leq L\chi - 2\alpha\chi \quad \text{on } [0, \infty) \times \mathbb{R}^d. \quad (3.10)$$

When  $F(x) = -\alpha x$ , inequality (3.10) is reduced to an equality.

*Proof.* We obtain (3.9) from (3.2). Differentiating equation (3.5) with respect to  $x_j$ , we have, by (3.3),

$$(D_j \eta)_t = \frac{1}{2} \Delta (D_j \eta) + \sum_{i=1}^d F_i [D_i (D_j \eta)] + \sum_{i=1}^d (D_j F_i) (D_i \eta) \quad \text{on } [0, \infty) \times \mathbb{R}^d. \quad (3.11)$$

On the other hand, we note that

$$\frac{1}{2}D_i\chi = \sum_{j=1}^d (D_j\eta)[D_i(D_j\eta)], \quad 1 \leq i \leq d, \quad (3.12)$$

$$\frac{1}{2}\Delta\chi = |D^2\eta|^2 + \sum_{j=1}^d (D_j\eta)[\Delta(D_j\eta)], \quad (3.13)$$

$$\sum_{i,j=1}^d (D_jF_i)(D_i\eta)(D_j\eta) \leq -\alpha\chi. \quad (3.14)$$

Here we used (2.1) in (3.14). Inequality (3.14) is reduced to an equality when  $F(x) = -\alpha x$ . Then, by (3.11)–(3.14), we obtain on  $[0, \infty) \times \mathbb{R}^d$

$$\begin{aligned} \frac{1}{2}\chi_t &= \sum_{j=1}^d (D_j\eta)(D_j\eta)_t \\ &= \frac{1}{2} \sum_{j=1}^d (D_j\eta)[\Delta(D_k\eta)] \\ &\quad + \sum_{i,j=1}^d F_i(D_j\eta)[D_i(D_j\eta)] + \sum_{i,j=1}^d (D_jF_i)(D_i\eta)(D_j\eta) \\ &\leq \frac{1}{2} \left( \frac{1}{2}\Delta\chi - |D^2\eta|^2 \right) + \frac{1}{2}F \cdot D\chi - \alpha\chi. \end{aligned} \quad (3.15)$$

Thus, (3.10) follows. It is easy to see that inequality (3.10) is reduced to an equality when  $F(x) = -\alpha x$ . The proof is complete.  $\square$

Now, we prove Theorem 2.1.

*Proof of Theorem 2.1.*

*Step 1.* In this step, we will show Theorem 2.1 under the assumption that  $\varphi \in C_0^\infty(\mathbb{R}^d)$ . We choose  $0 < T < +\infty$  arbitrarily. Integrating (3.10) over  $[0, T] \times \mathbb{R}^d$  with respect to  $dt \times d\nu$ , we have

$$\begin{aligned} &\int_0^T dt \int_{\mathbb{R}^d} |D^2\eta(t, \cdot)|^2 d\nu + \int_{\mathbb{R}^d} [ |D\eta(T, \cdot)|^2 - |D\varphi(\cdot)|^2 ] d\nu \\ &\leq \int_0^T dt \int_{\mathbb{R}^d} L\chi(t, \cdot) d\nu - 2\alpha \int_0^T dt \int_{\mathbb{R}^d} |D\eta(t, \cdot)|^2 d\nu. \end{aligned} \quad (3.16)$$

By Lemma 3.2, inequality (3.16) is reduced to an equality when  $F(x) = -\alpha x$ . Since  $\nu$  is the invariant probability measure for  $\{P_t\}$  as in (2.4), we have

$$\int_{\mathbb{R}^d} L\chi(t, \cdot) d\nu = 0, \quad t \geq 0. \quad (3.17)$$

On the other hand, by [2, Corollary 3.6], we have

$$\int_0^T dt \int_{\mathbb{R}^d} |DP_t \varphi|^2 d\nu = \int_{\mathbb{R}^d} [|\varphi|^2 - |P_T \varphi|^2] d\nu. \quad (3.18)$$

Thus, we obtain by (3.16)–(3.18)

$$\begin{aligned} & \int_0^T dt \int_{\mathbb{R}^d} |D^2 P_t \varphi|^2 d\nu + 2\alpha \int_{\mathbb{R}^d} [|\varphi|^2 - |P_T \varphi|^2] d\nu \\ & \leq \int_{\mathbb{R}^d} |D\varphi|^2 d\nu - \int_{\mathbb{R}^d} |DP_T \varphi|^2 d\nu, \quad T > 0. \end{aligned} \quad (3.19)$$

Now, let  $T$  tend to positive infinity in (3.19). Using (1.7) and the ergodic property

$$\lim_{T \rightarrow \infty} P_T \varphi(x) = \bar{\varphi}, \quad x \in \mathbb{R}^d \quad (3.20)$$

(cf. [2, (3.11)]), we have obtained (2.7). Then, by (1.5) and (3.18), we have (2.5) and (2.6). Since inequality (3.19) is reduced to the equality when  $F(x) = -\alpha x$ , it is not difficult to see (2.8).

*Step 2.* In this step, we conclude Theorem 2.1. Let  $\varphi \in H^1(\nu)$ . Since  $C_0^\infty(\mathbb{R}^d)$  is dense in  $H^1(\nu)$ , we can choose  $\{\varphi_n\} \subset C_0^\infty(\mathbb{R}^d)$  such that  $\varphi_n \rightarrow \varphi$  in  $H^1(\nu)$ . By Step 1, we see that

$$\int_0^\infty dt \int_{\mathbb{R}^d} |D^2 P_t \varphi_m - D^2 P_t \varphi_n|^2 d\nu \leq \int_{\mathbb{R}^d} |D\varphi_m - D\varphi_n|^2 d\nu. \quad (3.21)$$

Thus,  $\{D^2 P_t \varphi_n\}$  is a Cauchy sequence in  $L^2((0, \infty) \times \mathbb{R}^d, dt \times d\nu; \mathbb{R}^{d^2})$ . Hence, we find an element  $f \in L^2((0, \infty) \times \mathbb{R}^d, dt \times d\nu; \mathbb{R}^{d^2})$  such that

$$D^2 P_t \varphi_n(\cdot) \longrightarrow f(\cdot, \cdot) \quad \text{in } L^2((0, \infty) \times \mathbb{R}^d, dt \times d\nu; \mathbb{R}^{d^2}). \quad (3.22)$$

By the Fubini theorem, we see that  $f(t, \cdot) \in L^2(\mathbb{R}^d, \nu; \mathbb{R}^{d^2})$ ,  $t$ -a.e. On the other hand, by (3.22), we find a subsequence  $\{n_j\}$  such that

$$\int_{\mathbb{R}^d} |D^2 P_t \varphi_{n_j}(\cdot) - f(t, \cdot)|^2 d\nu \longrightarrow 0, \quad t\text{-a.e.} \quad (3.23)$$

This means that

$$D^2 P_t \varphi_{n_j}(\cdot) \longrightarrow f(t, \cdot) \quad \text{in } L^2(\mathbb{R}^d, \nu; \mathbb{R}^{d^2}), \quad t\text{-a.e.} \quad (3.24)$$

Since  $P_t \varphi_{n_j} \in H^2(\nu)$  ( $= \text{dom}_2(D^2)$ ) and  $D^2$  is a closed operator in  $L^2(\nu)$ , we obtain

$$P_t \varphi \in H^2(\nu), \quad f(t, \cdot) = D^2 P_t \varphi(\cdot), \quad t\text{-a.e.} \quad (3.25)$$

Then we obtain (2.5). Next, by (3.24), (3.25), Step 1, and Fatou's lemma, we have

$$\begin{aligned} \int_0^\infty dt \int_{\mathbb{R}^d} |D^2 P_t \varphi|^2 d\nu &\leq \liminf_{n \rightarrow \infty} \int_0^\infty dt \int_{\mathbb{R}^d} |D^2 P_t \varphi_{n_j}|^2 d\nu \\ &\leq \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^d} |D \varphi_{n_j}|^2 d\nu \\ &= \int_{\mathbb{R}^d} |D \varphi|^2 d\nu. \end{aligned} \quad (3.26)$$

Hence, by (1.5) and (3.18), we obtain (2.6). Finally, by (3.22) and (3.25), we conclude that

$$D^2 P_n \varphi_n(\cdot) \longrightarrow D^2 P \varphi(\cdot) \quad \text{in } L^2((0, \infty) \times \mathbb{R}^d, dt \times d\nu; \mathbb{R}^{d^2}). \quad (3.27)$$

Therefore, (2.7) follows from Step 1. By (3.27) and Step 1, it is easy to see (2.8). The proof is complete.  $\square$

## References

- [1] S. Cerrai, *Second Order PDE's in Finite and Infinite Dimension. A Probabilistic Approach*, Lecture Notes in Mathematics, vol. 1762, Springer-Verlag, Berlin, 2001.
- [2] G. Da Prato and B. Goldys, *Elliptic operators on  $\mathbb{R}^d$  with unbounded coefficients*, J. Differential Equations **172** (2001), no. 2, 333–358, *Erratum*, J. Differential Equations, **184** (2002), no. 2, p. 620.

Yasuhiro Fujita: Department of Mathematics, Toyama University, Toyama 930-8555, Japan  
*E-mail address:* yfujita@sci.toyama-u.ac.jp