

CLASSES OF ELLIPTIC MATRICES

ANTONIO TARSIA

Received 12 December 2005; Revised 20 February 2006; Accepted 21 February 2006

The equivalence between some conditions concerning elliptic matrices is shown, namely, the Cordes condition, a generalized form of Campanato's condition, and a generalized form of a condition of Buică.

Copyright © 2006 Antonio Tarsia. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let Ω be an open bounded set in \mathbb{R}^n , $n > 2$, with a sufficiently regular boundary, and let $A(x) = \{a_{ij}(x)\}_{i,j=1,\dots,n}$ be a real matrix, with coefficients $a_{ij} \in L^\infty(\Omega)$. We consider the following problem:

$$\begin{aligned} u &\in H^{2,2} \cap H_0^{1,2}(\Omega), \\ \sum_{i,j=1}^n a_{ij}(x) D_{ij} u(x) &= f(x), \quad \text{a.e. } x \in \Omega. \end{aligned} \tag{1.1}$$

If $f \in L^2(\Omega)$, it is known (see the counterexamples in [6]) that problem (1.1) is not well posed with the only hypothesis of uniform ellipticity on the matrix $A(x)$: there exists a positive constant $\bar{\nu}$ such that

$$\sum_{i,j=1}^n a_{ij}(x) \eta_i \eta_j \geq \bar{\nu} \|\eta\|_n^2, \quad \text{a.e. in } \Omega, \quad \forall \eta = (\eta_1, \dots, \eta_n) \in \mathbb{R}^n. \tag{1.2}$$

It is therefore essential, in order to be able to solve Problem (1.1), to assume some hypotheses on $A(x)$ stronger than (1.2). In this paper we consider some of these ones and compare them. More precisely, we will consider the following *conditions* and show that they are equivalent.

2 Classes of elliptic matrices

Condition 1.1 (the Cordes condition, see [5, 8]). $\|A(x)\|_{\mathbb{R}^{n^2}} \neq 0$, a.e. in Ω , and there exists $\varepsilon \in (0, 1)$ such that

$$\frac{(\sum_{i,j=1}^n a_{ii}(x))^2}{\sum_{i,j=1}^n a_{ij}^2(x)} \geq n - 1 + \varepsilon, \quad \text{a.e. in } \Omega. \quad (1.3)$$

Condition 1.2 (Condition A_{xp}). There exist four real constants $\sigma, \gamma, \delta, p$ with $\sigma > 0, \gamma > 0, \delta \geq 0, \gamma + \delta < 1, p \geq 1$, and a function $a(x) \in L^\infty(\Omega)$, with $a(x) \geq \sigma$ a.e. in Ω , such that

$$\left| \sum_{i=1}^n \xi_{ii} - a(x) \sum_{i,j=1}^n a_{ij}(x) \xi_{ij} \right|^p \leq \gamma \|\xi\|_{n^2}^p + \delta \left| \sum_{i=1}^n \xi_{ii} \right|^p \quad (1.4)$$

for all $\xi = \{\xi_{ij}\}_{i,j=1,\dots,n} \in \mathbb{R}^{n^2}$, a.e. in Ω .

When $p = 1$, the above *condition* will be simply denoted by *Condition A_x* ; it was defined in [10], where it has also been shown to be equivalent to the *Cordes condition*. If $a(x)$ is constant on Ω , *Condition A_x* is the formulation for linear operators of Campanato's *condition A*, (see [4]), which was defined for nonlinear operators. A particular version of *Condition A_{xp}* , that is, with $p = 2$ and (x) constant, is stated in [7] for nonlinear operators.

Condition 1.3 (Condition B_x). There exist four real positive real constants σ, c_1, c_2, c_3 and a function $\beta \in L^\infty(\Omega)$ such that

- (i) $0 < c_1 - c_2 - c_3 < 1$,
- (ii) $\beta(x) \geq \sigma$ a.e. in Ω ,

and moreover

$$\beta(x) \sum_{i,j=1}^n a_{ij}(x) \xi_{ij} \sum_{i=1}^n \xi_{ii} \geq c_1 \left(\sum_{i=1}^n \xi_{ii} \right)^2 - c_2 \left| \sum_{i=1}^n \xi_{ii} \right| \|\xi\|_{n^2} - c_3 \|\xi\|_{n^2}^2 \quad (1.5)$$

for all $\xi = \{\xi_{ij}\}_{i,j=1,\dots,n} \in \mathbb{R}^{n^2}$, a.e. in Ω .

If $\beta(x)$ is constant on Ω , we will denote this *condition* as *Condition B*; it has been defined by Buičă in [2].

The importance of *Conditions A_{xp}* or B_x is in the fact that they allow to show in a relatively simple manner, by means of *near operators theory* (see [4, 9]) or *weakly near operators theory* (see [1–3]), that problem (1.1) is well posed. The usefulness of showing the equivalence among these *conditions* is due to the fact that to verify whether a matrix satisfies *Condition A_{xp}* or B_x is very complicated, even if $n = 2$, while to verify whether it satisfies the *Cordes condition* is much simpler.

2. A procedure of decomposition for matrices

In this section we consider a short procedure of decomposition of the matrices A and I which has been developed in [10]. We set

$$\begin{aligned}\Omega_0 &= \{x \in \Omega : \text{there exists } b(x) \in \mathbb{R} \text{ such that } b(x)A(x) = I\}; \\ \Omega_1 &= \Omega \setminus \Omega_0.\end{aligned}\tag{2.1}$$

Remark 2.1. Set $M = \sup_{\Omega} \|A(x)\|$, $\bar{\nu} = \inf_{\Omega} \|A(x)\|$, accordingly $n\bar{\nu} \leq (A(x) | I) \leq nM$. Then, for each $x \in \Omega_0$, we obtain $1/M \leq b(x) \leq 1/\bar{\nu}$.

We can assume $\text{meas}\Omega_1 > 0$, since otherwise as we will see in the following it is easy to show the equivalence between the above *conditions*. We set for all $x \in \Omega_1$: $W(x) = \{B(x) : B(x) = sI + rA(x), s, r \in \mathbb{R}\}$; $\Sigma_x = W(x) \cap S(I, 1)$ (where $S(I, 1) = \{B : \|B - I\|_{\mathbb{R}^{n^2}} < 1\}$).

Let $v_1, w_2 \in W(x)$ be the projections of I on the lines through the zero vector of \mathbb{R}^{n^2} and tangent to Σ_x . Moreover let v_2 be the projection of I on the line through the zero vector of \mathbb{R}^{n^2} and perpendicular to v_1 , and let w_1 be the projection of I on the line through the zero vector of \mathbb{R}^{n^2} and perpendicular to w_2 . In this manner we find two systems of orthogonal vectors $\{v_1, v_2\}$, $\{w_1, w_2\}$, with $v_i = v_i(x)$, $w_i = w_i(x)$, $i = 1, 2$. Each of them is a basis in the plane $W(x)$. Then $I = v_1 + v_2 = w_1 + w_2$, and there are L^∞ functions $a_i = a_i(x)$ and $b_i = b_i(x)$, $i = 1, 2$, such that

$A(x) = a_1(x)v_1(x) + a_2(x)v_2(x) = b_1(x)w_1(x) + b_2(x)w_2(x)$. (As $\|v_1\| = \|w_2\| = \sqrt{n-1}$ and $\|v_2\| = \|w_1\| = 1$, then for $i = 1, 2$, $a_i^2 \leq a_1^2(n-1) + a_2^2 = (a_1v_1 + a_2v_2 | a_1v_1 + a_2v_2) = (A(x) | A(x)) = \|A(x)\|^2$; here if $B = \{b_{ij}\}_{i,j=1,\dots,n}$ and $C = \{c_{ij}\}_{i,j=1,\dots,n}$, we set $(B | C) = \sum_{i,j=1}^n b_{ij}c_{ij}$.) Set

$$\begin{aligned}Q_v(x, \nu, \tau) &= \{\xi \in \mathbb{R}^{n^2} : \xi = s\nu_1 + t\nu_2, 0 < \nu \leq s, t \leq \tau\}, \\ Q_w(x, \nu, \tau) &= \{\xi \in \mathbb{R}^{n^2} : \xi = s\nu_1 + t\nu_2, 0 < \nu \leq s, t \leq \tau\}, \\ R(x, \nu_0, \tau_0) &= \{\xi \in \mathbb{R}^{n^2} : \xi = s\nu_2 + t\nu_1, 0 < \nu_0 \leq s, t \leq \tau_0\}, \\ C(\Sigma_x) &= \{\nu : \nu \in W(x) \text{ such that } \exists z \in \Sigma_x, \exists t > 0 \text{ for which } \nu = tz\}, \\ C_\rho(x) &= \{\nu : \nu \in C(\Sigma_x) : \exists t > 0 \text{ such that } \|I - t\nu\| < \rho\}, \quad 0 < \rho < 1.\end{aligned}\tag{2.2}$$

The following propositions are proved in [10].

PROPOSITION 2.2. For all $\tau, \nu > 0$ with $\nu \leq \tau$, $\exists \tau_0, \nu_0, 0 < \tau_0 < \nu_0$, such that for all $x \in \Omega_1$,

$$Q_v(x, \nu, \tau) \cap Q_w(x, \nu, \tau) \subset R(x, \nu_0, \tau_0).\tag{2.3}$$

PROPOSITION 2.3. For all $\tau_0, \nu_0, 0 < \tau_0 < \nu_0$, there exists $\rho \in (0, 1)$ such that for all $x \in \Omega_1$,

$$R(x, \nu_0, \tau_0) \subset C_\rho(x).\tag{2.4}$$

3. Condition B_x

PROPOSITION 3.1. Condition A_x and Condition B_x are equivalent.

4 Classes of elliptic matrices

Proof. We assume that A satisfies *Condition* A_x . It follows (from (1.4) with $p = 1$) by squaring both members

$$(I | \xi)^2 - 2a(x)(A | \xi)(I | \xi) \leq \gamma^2 \|\xi\|^2 + 2\gamma\delta |(I | \xi)| \|\xi\| + \delta^2 (I | \xi)^2 \quad (3.1)$$

then

$$2a(x)(A | \xi)(I | \xi) \geq (1 - \delta^2)(I | \xi)^2 - 2\gamma\delta |(I | \xi)| \|\xi\| - \gamma^2 \|\xi\|^2. \quad (3.2)$$

This is *Condition* B_x with $b(x) = 2a(x)$, $c_1 = 1 - \delta^2$, $c_2 = 2\gamma\delta$, $c_3 = \gamma^2$. \square

Conversely, we set $\mathbf{A}(x) = \beta(x)A(x)$ and assume that *Condition* B holds for \mathbf{A} , then we will show that \mathbf{A} also satisfies *Condition* A_x . To this purpose we write *Condition* B in the following form: there exist four real positive constants M , c_1 , c_2 , c_3 with $0 < c_1 - c_2 - c_3 < 1$, $\sup_{x \in \Omega} \|\mathbf{A}(x)\| \leq M$ such that

$$(\mathbf{A}(x) | \xi)(I | \xi) \geq c_1 (I | \xi)^2 - c_2 |(I | \xi)| \|\xi\| - c_3 \|\xi\|^2, \quad (3.3)$$

for all $\xi \in \mathbb{R}^n$, a.e. in Ω . Then we obtain the thesis by using the decomposition of \mathbf{A} and I stated in Section 2. For this we distinguish two cases: $x \in \Omega_0$ and $x \in \Omega_1$.

If $x \in \Omega_0$, that is, there exists $b(x)$ such that $b(x)\mathbf{A}(x) = I$, then *Condition* A_x is trivially true (take in (1.4) $a(x) = b(x)$).

Instead, if $x \in \Omega_1$, with $\text{meas}\Omega_1 > 0$, we observe that (3.3) holds in particular for $\xi \in W(x)$. So we can write ξ as a linear combination of the basis $\{v_1(x), v_2(x)\}$. Now, let $t_1, t_2 \in \mathbb{R}$ be such that $\xi = t_1 v_1(x) + t_2 v_2(x)$, accordingly $\|\xi\|^2 = (\xi | \xi) = t_1^2(n-1) + t_2^2$, then

$$\begin{aligned} (\mathbf{A} | \xi) &= (a_1(x)v_1 + a_2(x)v_2 | t_1 v_1 + t_2 v_2) = a_1 t_1(n-1) + a_2 t_2, \\ (I | \xi) &= (v_1 + v_2 | t_1 v_1 + t_2 v_2) = t_1(n-1) + t_2. \end{aligned} \quad (3.4)$$

Now, (3.4) and the above remarks yield the following form of *Condition* B : for each $\xi \in W(x)$,

$$\begin{aligned} (\mathbf{A} | \xi)(I | \xi) &= [a_1 t_1(n-1) + a_2 t_2][t_1(n-1) + t_2] \\ &\geq c_1 [t_1(n-1) + t_2]^2 - c_2 [t_1(n-1) + t_2] \sqrt{t_1^2(n-1) + t_2^2} - c_3 [t_1^2(n-1) + t_2^2]. \end{aligned} \quad (3.5)$$

Put

$$\begin{aligned} F(t_1, t_2) &= [a_1 t_1(n-1) + a_2 t_2][t_1(n-1) + t_2] - c_1 [t_1(n-1) + t_2]^2 \\ &\quad + c_2 [t_1(n-1) + t_2] \sqrt{t_1^2(n-1) + t_2^2} + c_3 [t_1^2(n-1) + t_2^2]. \end{aligned} \quad (3.6)$$

Remark that

$$F(t_1, t_2) \geq 0, \quad \forall (t_1, t_2) \in \mathbb{R}^2 \text{ (by (3.5)).} \quad (3.7)$$

In particular

$$F\left(\frac{1}{\sqrt{n-1}}, 0\right) = a_1(n-1) - c_1(n-1) + c_2\sqrt{n-1} + c_3 \geq 0 \quad (3.8)$$

from which

$$a_1(x) \geq c_1 - \frac{c_2}{\sqrt{n-1}} - \frac{c_3}{n-1} \geq c_1 - c_2 - c_3 > 0. \quad (3.9)$$

While the inequality $F(0, 1) = a_2(x) - c_1 + c_2 + c_3 \geq 0$ implies $a_2(x) \geq c_1 - c_2 - c_3 > 0$.

In the same way, by taking the system of orthogonal vectors $\{w_1, w_2\}$ as basis of $W(x)$, it follows that

$$b_i(x) \geq c_1 - c_2 - c_3 > 0, \quad i = 1, 2, x \in \Omega_1. \quad (3.10)$$

So we have shown (see Section 2) that $\mathbf{A}(x) \in Q_v(x, \nu, \tau) \cap Q_w(x, \nu, \tau)$. This implies, by Proposition 2.2, $\mathbf{A}(x) \in R(x, \nu_0, \tau_0)$, then by Proposition 2.3, $\mathbf{A}(x) \in C_\rho(x)$, which is equivalent to say that *Condition* A_x is valid with $\delta = 0$.

Taking into account this proposition and the equivalence between the *Cordes condition* and *Condition* A_x , shown in [10], we have the following.

COROLLARY 3.2. *Condition* B_x and the *Cordes condition* are equivalent.

The following example states that *Condition* B is stronger than *Condition* A_x and therefore is also stronger than the *Cordes condition*.

Example 3.3. Let $\Omega = \Omega_1 \cup \Omega_2$, where $\Omega_1 = \{(x_1, x_2) \in \mathbb{R}^2 : 0 < x_1 < 1, 0 < x_2 \leq 1\}$ and $\Omega_2 = \{(x_1, x_2) \in \mathbb{R}^2 : 0 < x_1 < 1, 1 < x_2 < 2\}$, moreover

$$A(x) = \begin{cases} A_1, & \text{if } x \in \Omega_1, \\ A_2, & \text{if } x \in \Omega_2, \end{cases} \quad A_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 200 & -150 \\ -150 & 200 \end{pmatrix}. \quad (3.11)$$

A is uniformly elliptic on Ω and, since $n = 2$, this implies the *Cordes condition* and therefore also *Condition* A_x (see [10]). Nevertheless A does not satisfy *Condition* B . Indeed, we consider $x \in \Omega_1$, then $A(x) = A_1$. We observe that if A_1 satisfied *Condition* B , it would be

$$(A_1 | \xi)(I | \xi) \geq c_1(I | \xi)^2 - c_2|(I | \xi)|\|\xi\| - c_3\|\xi\|^2 \quad (3.12)$$

for each $\xi \in \mathbb{R}^4$, that is,

$$(1 - c_1)(I | \xi)^2 + c_2|(I | \xi)|\|\xi\| + c_3\|\xi\|^2 \geq 0. \quad (3.13)$$

The bilinear form $\Phi(X, Y) = (1 - c_1)X^2 + c_2XY + c_3Y^2$, where $(X, Y) \in \mathbb{R}^2$, is nonnegative if $(1 - c_1)c_3 \geq c_2^2/4$. In particular it must hold $c_1 < 1$. Otherwise if $A(x)$ satisfied *Condition* B on Ω_2 it would be

$$(A_2 | \xi)(I | \xi) \geq c_1(I | \xi)^2 - c_2|(I | \xi)|\|\xi\| - c_3\|\xi\|^2, \quad (3.14)$$

6 Classes of elliptic matrices

where c_1, c_2, c_3 are the above determined constants for the matrix A_1 . Now we consider the matrix

$$\xi = \begin{pmatrix} -1 & 0 \\ -2 & 0 \end{pmatrix}, \quad (3.15)$$

by replacing it in (3.14), we obtain $-100 \geq c_1 - c_2\sqrt{5} - 5c_3$, that is, $c_2(\sqrt{5} - 1) + 4c_3 \geq c_1 - c_2 - c_3 + 100$; that implies (because by hypothesis it holds $c_1 > c_2 + c_3$) $4c_1 > 4(c_2 + c_3) \geq 100$, then $c_1 \geq 25$. This contradicts what we have obtained for A_1 , that is, $c_1 < 1$.

4. Condition A_{xp}

We prove equivalence between the *Cordes condition* and *Condition A_{xp}* in the same way used in [10] for the proof of equivalence between *Condition A* and the *Cordes condition*. The first step is following.

LEMMA 4.1. *Condition A_{xp} with $\delta = 0$ is equivalent to Cordes Condition.*

Proof (see also [10]). We can write *Condition A_{xp}* , if $\delta = 0$, as follows:

$$|(I - a(x)A(x) | \xi) | \leq \gamma^{1/p} \|\xi\| \quad (4.1)$$

for all $\xi \in \mathbb{R}^n$, and $p \geq 1$. This is just *Condition A_x* with $\delta = 0$ and, accordingly to what proved in [10], this is equivalent to the *Cordes condition*. \square

The second step for the achievement of our goal is following.

LEMMA 4.2. *If $A(x)$ satisfies Condition A_{xp} for some function $a(x)$ and some constants σ, γ, δ , then it satisfies the same condition with $\delta = 0$ and possibly different $\sigma, \gamma, a(x)$.*

Proof. We proceed on the line of the proof of [10, Lemma 3.3]. We follow the notations of Section 2. *Condition A_{xp}* , with $\delta \neq 0$, yields *Condition A_{xp}* with $\delta = 0$, by replacing the coefficient $a(x)$ of the first *condition* with a new coefficient $\bar{a}(x)$, defined by

$$\bar{a}(x) = \begin{cases} b(x), & \text{if } x \in \Omega_0, \\ c(x), & \text{if } x \in \Omega_1. \end{cases} \quad (4.2)$$

If $x \in \Omega_0$, then *Condition A_{xp}* with $\delta = 0$ is trivially satisfied. Moreover, by Remark 2.1, $1/M \leq b(x) \leq 1/\bar{\nu}$. Now let $x \in \Omega_1$. We prove the existence of a function $c(x)$ by means of the decomposition of matrices $A(x)$, I stated in Section 2 and replacing the expressions obtained in *Condition A_{xp}* :

$$\begin{aligned} |(I - a(x)A(x) | \xi) |^p &= |(v_1 + v_2 - a(x)(a_1 v_1 + a_2 v_2) | \xi) |^p \\ &= (\text{take } \xi = v_i, i = 1, 2) \\ &= |(v_1 + v_2 - a(x)(a_1 v_1 + a_2 v_2) | v_i) |^p = \left| \|v_i\|^2 - a(x)a_i \|v_i\|^2 \right|^p \\ &= |1 - a(x)a_i|^p \|v_i\|^{2p} \leq \gamma \|v_i\|^p + \delta (v_1 + v_2 | v_i)^p = \gamma \|v_i\|^p + \delta \|v_i\|^{2p}. \end{aligned} \quad (4.3)$$

From this

$$\frac{1}{a(x)} \left(1 - \frac{\sqrt[p]{\gamma + \delta \|v_i\|^p}}{\|v_i\|} \right) \leq a_i \leq \frac{1}{a(x)} \left(1 + \frac{\sqrt[p]{\gamma + \delta \|v_i\|^p}}{\|v_i\|} \right). \quad (4.4)$$

We observe that

$$1 - (\gamma + \delta)^{1/p} \leq 1 - \frac{\sqrt[p]{\gamma + \delta \|v_i\|^p}}{\|v_i\|}, \quad 1 + \frac{\sqrt[p]{\gamma + \delta \|v_i\|^p}}{\|v_i\|} \leq 1 + (\gamma + \delta)^{1/p}. \quad (4.5)$$

Using $\|v_1\| = \sqrt{n-1}$, $v_2 = 1$, we can write

$$\frac{\gamma + \delta \|v_i\|^p}{\|v_i\|^p} \leq \gamma + \delta, \quad i = 1, 2. \quad (4.6)$$

We conclude, from (4.4), by setting

$$M_1 = \sup_{\Omega} a(x), \quad \nu = \frac{1}{M_1} \left[1 - (\gamma + \delta)^{\frac{1}{p}} \right], \quad \tau = \frac{1}{\sigma} \left[1 + (\gamma + \delta)^{1/p} \right] \quad (4.7)$$

for all $x \in \Omega_1$, $A(x) \in Q_\nu(x, \nu, \tau)$. Then by taking $\xi = w_i$ ($i = 1, 2$) in *Condition* A_{xp} , with similar calculations, we obtain for all $x \in \Omega_1$, $A(x) \in Q_w(x, \nu, \tau)$. Then for all $x \in \Omega_1$, $A(x) \in Q_\nu(x, \nu, \tau) \cap Q_w(x, \nu, \tau)$. From Proposition 2.2 it follows that there exist ν_0, τ_0 , with $0 < \nu_0 < \tau_0$, such that $A(x) \in R(x, \nu_0, \tau_0)$. By Proposition 2.3 there exists $\rho \in (0, 1)$ such that $A(x) \in C_\rho(x)$, that is, there exist $c(x) > 0$ and $\rho \in (0, 1)$ such that

$$\|I - c(x)A(x)\| \leq \rho. \quad (4.8)$$

(This inequality also implies $(\sqrt{n}-1)/M < c(x) < (\sqrt{n}+1)/\bar{\nu}$, $x \in \Omega_1$.) \square

From Lemmas 4.1 and 4.2 we have the following.

THEOREM 4.3. *The Cordes condition and Condition A_{xp} are equivalent.*

This theorem and Corollary 3.2 imply the following.

COROLLARY 4.4. *Condition B_x and Condition A_{xp} are equivalent.*

Theorem 4.3 and Corollary 3.2, by the results proved in [10], imply the following.

COROLLARY 4.5. *Let $n = 2$. Then every uniformly elliptic symmetric matrix satisfies Condition A_{xp} and Condition B_x .*

References

- [1] A. Buică, *Some properties preserved by weak nearness*, Seminar on Fixed Point Theory Cluj Napoca 2 (2001), 65–70.
- [2] ———, *Existence of strong solutions of fully nonlinear elliptic equations*, Proceedings of Conference on Analysis and Optimization of Differential Systems, Constanta, September 2002.
- [3] A. Buică and A. Domokos, *Nearness, accretivity, and the solvability of nonlinear equations*, Numerical Functional Analysis and Optimization 23 (2002), no. 5-6, 477–493.

8 Classes of elliptic matrices

- [4] S. Campanato, *A Cordes type condition for nonlinear nonvariational systems*, Rendiconti Accademia Nazionale delle Scienze detta dei XL. Serie V. Memorie di Matematica. Parte I **13** (1989), no. 1, 307–321.
- [5] H. O. Cordes, *Zero order a priori estimates for solutions of elliptic differential equations*, Proceedings of Symposium in Pure Math., vol. 4, American Mathematical Society, Rhode Island, 1961, pp. 157–166.
- [6] O. A. Ladyzhenskaya and N. N. Ural'tseva, *Linear and Quasilinear Elliptic Equations*, Academic Press, New York, 1968.
- [7] A. Maugeri, D. K. Palagachev, and L. G. Softova, *Elliptic and Parabolic Equations with Discontinuous Coefficients*, Wiley-VCH, Berlin, 2000.
- [8] G. Talenti, *Sopra una classe di equazioni ellittiche a coefficienti misurabili*, Annali di Matematica Pura ed Applicata. Serie Quarta **69** (1965), 285–304.
- [9] A. Tarsia, *Some topological properties preserved by nearness between operators and applications to P.D.E.*, Czechoslovak Mathematical Journal **46** (1996), no. 4, 607–624.
- [10] ———, *On Cordes and Campanato conditions*, Archives of Inequalities and Applications **2** (2004), no. 1, 25–39.

Antonio Tarsia: Dipartimento di Matematica “L. Tonelli,” Università di Pisa,
Largo Bruno Pontecorvo 5, 56127 Pisa, Italy
E-mail address: tarsia@dm.unipi.it