

RESEARCH

Open Access

On \ast -class A contractions

Fugen Gao^{*} and Xiaochun Li

^{*}Correspondence:
gaofugen08@126.com
College of Mathematics and
Information Science, Henan Normal
University, Xinxiang, Henan 453007,
China

Abstract

A Hilbert space operator T belongs to \ast -class A if $|T^2| - |T^*|^2 \geq 0$. The famous Fuglede-Putnam theorem is as follows: the operator equation $AX = XB$ implies $A^*X = XB^*$ when A and B are normal operators. In this paper, firstly we prove that if T is a contraction of \ast -class A operators, then either T has a nontrivial invariant subspace or T is a proper contraction and the nonnegative operator $D = |T^2| - |T^*|^2$ is a strongly stable contraction; secondly, we show that if X is a Hilbert-Schmidt operator, A and $(B^*)^{-1}$ are \ast -class A operators such that $AX = XB$, then $A^*X = XB^*$.

MSC: 47B20; 47A63

Keywords: \ast -class A operators; contraction operators; the Fuglede-Putnam theorem

1 Introduction

Let \mathcal{H} be a complex Hilbert space and let \mathbb{C} be the set of complex numbers. Let $B(\mathcal{H})$ denote the C^* -algebra of all bounded linear operators acting on \mathcal{H} . For operators $T \in B(\mathcal{H})$, we shall write $\ker T$ and $\operatorname{ran} T$ for the null space and the range of T , respectively. Also, let $\sigma(T)$ denote the spectrum of T .

Recall that $T \in B(\mathcal{H})$ is called p -hyponormal for $p > 0$ if $(T^*T)^p - (TT^*)^p \geq 0$ [1]; when $p = 1$, T is called hyponormal. And T is called paranormal if $\|Tx\|^2 \leq \|T^2x\|\|x\|$ for all $x \in \mathcal{H}$ [2, 3]. And T is called normaloid if $\|T^n\| = \|T\|^n$ for all $n \in \mathbb{N}$ (equivalently, $\|T\| = r(T)$, the spectral radius of T). In order to discuss the relations between paranormal and p -hyponormal and log-hyponormal operators (T is invertible and $\log T^*T \geq \log TT^*$), Furuta *et al.* [4] introduced a very interesting class of operators: class A defined by $|T^2| - |T|^2 \geq 0$, where $|T| = (T^*T)^{\frac{1}{2}}$ which is called the absolute value of T ; and they showed that the class A is a subclass of paranormal and contains p -hyponormal and log-hyponormal operators. Recently Duggal *et al.* [5] introduced \ast -class A operators (i.e., $|T^2| - |T^*|^2 \geq 0$) and \ast -paranormal operators (i.e., $\|T^*x\|^2 \leq \|T^2x\|\|x\|$ for all $x \in \mathcal{H}$); and they proved that a \ast -class A operator is a generalization of hyponormal operator and \ast -class A operators form a subclass of the class of \ast -paranormal operators.

A contraction is an operator T such that $\|T\| \leq 1$. A contraction T is said to be a proper contraction if $\|Tx\| < \|x\|$ for every nonzero $x \in \mathcal{H}$. A strict contraction is an operator T such that $\|T\| < 1$. A strict contraction is a proper contraction, but a proper contraction is not necessary a strict contraction, although the concepts of strict and proper contractions coincide for compact operators. A contraction T is of class C_0 if $\|T^n x\| \rightarrow 0$ when $n \rightarrow \infty$ for every $x \in \mathcal{H}$ (i.e., T is a strongly stable contraction) and it is said to be of class C_1 if $\lim_{n \rightarrow \infty} \|T^n x\| > 0$ for every nonzero $x \in \mathcal{H}$. Classes C_0 and C_1 are defined by considering T^* instead of T , and we define the class $C_{\alpha\beta}$ for $\alpha, \beta = 0, 1$ by $C_{\alpha\beta} = C_\alpha \cap C_\beta$. An isometry is a contraction for which $\|Tx\| = \|x\|$ for every $x \in \mathcal{H}$.

In this paper, firstly we prove that if T is a contraction of $*$ -class A operators, then either T has a nontrivial invariant subspace or T is a proper contraction and the nonnegative operator $D = |T^2| - |T^*|^2$ is a strongly stable contraction; secondly, we show that if X is a Hilbert-Schmidt operator, A and $(B^*)^{-1}$ are $*$ -class A operators such that $AX = XB$, then $A^*X = XB^*$.

2 On $*$ -class A contractions

Theorem 2.1 *If T is a contraction of $*$ -class A operators, then the nonnegative operator $D = |T^2| - |T^*|^2$ is a contraction whose power sequence $\{D^n\}$ converges strongly to a projection P , and $T^*P = 0$.*

Proof Suppose that T is a contraction of $*$ -class A operators, then $D = |T^2| - |T^*|^2 \geq 0$. Let $R = D^{\frac{1}{2}}$. Then for every $x \in \mathcal{H}$,

$$\begin{aligned} \langle D^{n+1}x, x \rangle &= \|R^{n+1}x\|^2 = \langle DR^n x, R^n x \rangle \\ &= \langle |T^2| R^n x, R^n x \rangle - \langle |T^*|^2 R^n x, R^n x \rangle \\ &= \| |T^2|^{\frac{1}{2}} R^n x \|^2 - \| |T^*| R^n x \|^2 \\ &\leq \| R^n x \|^2 - \| T^* R^n x \|^2 \\ &\leq \| R^n x \|^2 \\ &= \langle D^n x, x \rangle. \end{aligned}$$

Thus R (and so D) is a contraction and $\{D^n\}$ is a decreasing sequence of nonnegative contractions. Hence $\{D^n\}$ converges strongly to a projection P . Moreover,

$$\sum_{n=0}^m \|T^* R^n x\|^2 \leq \sum_{n=0}^m (\|R^n x\|^2 - \|R^{n+1} x\|^2) = \|x\|^2 - \|R^{m+1} x\|^2 \leq \|x\|^2$$

for all nonnegative integers m and every $x \in \mathcal{H}$. Therefore $\|T^* R^n x\| \rightarrow 0$ as $n \rightarrow \infty$, hence we have

$$T^* P x = T^* \lim_{n \rightarrow \infty} D^n x = \lim_{n \rightarrow \infty} T^* R^{2n} x = 0$$

for every $x \in \mathcal{H}$. So that $T^* P = 0$. □

Theorem 2.2 *Let T be a contraction of $*$ -class A operators. If T has no nontrivial invariant subspace, then*

- (i) T is a proper contraction;
- (ii) the nonnegative operator $D = |T^2| - |T^*|^2$ is a strongly stable contraction (so that $D \in C_{00}$).

Proof (i) Suppose that T is a $*$ -class A operator, then $|T^*|^2 \leq |T^2|$. We have

$$\|T^* x\|^2 = \langle |T^*|^2 x, x \rangle \leq \langle |T^2| x, x \rangle = \| |T^2|^{\frac{1}{2}} x \|^2 = \|T^2 x\| \|x\|$$

for every $x \in \mathcal{H}$. By [6] Theorem 3.6, we have that

$$T^*Tx = \|T\|^2x \quad \text{if and only if} \quad \|Tx\| = \|T\|\|x\|.$$

Put $\mathcal{U} = \{x \in \mathcal{H} : \|Tx\| = \|T\|\|x\|\} = \ker(|T|^2 - \|T\|^2)$, which is a subspace of \mathcal{H} . In the following, we shall show that \mathcal{U} is an invariant subspace of T . For every $x \in \mathcal{U}$, we have

$$\begin{aligned} \|T(Tx)\|^2 &\leq \|T\|^2\|Tx\|^2 = \|T\|^4\|x\|^2 = \|\|T\|^2x\|^2 = \|T^*Tx\|^2 \\ &\leq \|T^2Tx\|\|Tx\| = \|T^2Tx\|\|T\|\|x\|, \end{aligned} \quad (2.1)$$

where the second inequality holds since T is a $*$ -class A operator. So, we have that

$$\|T\|^4\|x\|^2 \leq \|T^2Tx\|\|T\|\|x\|,$$

that is, $\|T\|^3\|x\| \leq \|T^2Tx\|$. Hence we have

$$\|T\|^3\|x\| = \|T^2Tx\|. \quad (2.2)$$

By (2.2), we have

$$\|T\|^3\|x\| = \|T^2Tx\| \leq \|T\|\|T(Tx)\|, \quad (2.3)$$

that is, $\|T\|^2\|x\| \leq \|T(Tx)\|$. Hence

$$\|T\|^2\|x\| = \|T(Tx)\|. \quad (2.4)$$

Hence by (2.1) and (2.4), we have

$$\|T^2Tx\|\|T\|\|x\| = \|T(Tx)\|^2.$$

So, we have that $\|T(Tx)\| = \|T\|\|Tx\|$. That is, \mathcal{U} is an invariant subspace of T . Now suppose T is a contraction of $*$ -class A operators. If T is a strict contraction, then it is trivially a proper contraction. If T is not a strict contraction (i.e., $\|T\| = 1$) and T has no nontrivial invariant subspace, then $\mathcal{U} = \{x \in \mathcal{H} : \|Tx\| = \|x\|\} = \{0\}$ (actually, if $\mathcal{U} = \mathcal{H}$, then T is an isometry, and isometries have nontrivial invariant subspaces). Thus, for every nonzero $x \in \mathcal{H}$, $\|Tx\| < \|x\|$, so T is a proper contraction.

(ii) Let T be a contraction of $*$ -class A operators. By Theorem 2.1 we have D is a contraction, $\{D^n\}$ converges strongly to a projection P , and $T^*P = 0$. So, $PT = 0$. Suppose T has no nontrivial invariant subspace. Since $\ker P$ is a nonzero invariant subspace for T whenever $PT = 0$ and $T \neq 0$, it follows that $\ker P = \mathcal{H}$. Hence $P = 0$ and so D^n converges strongly to 0, that is, $D = |T^2| - |T^*|^2$ is a strongly stable contraction. D is self-adjoint, so that $D \in C_{00}$. \square

Since a self-adjoint operator T is a proper contraction if and only if T is a C_{00} -contraction, we have the following corollary by Theorem 2.2.

Corollary 2.3 *Let T be a contraction of $*$ -class A operators. If T has no nontrivial invariant subspace, then both T and the nonnegative operator $D = |T^2| - |T^*|^2$ are proper contractions.*

3 The Fuglede-Putnam theorem for $*$ -class A operators

The famous Fuglede-Putnam theorem is as follows [3, 7, 8].

Theorem 3.1 *Let A and B be normal operators and X be an operator such that $AX = XB$, then $A^*X = XB^*$.*

The Fuglede-Putnam theorem was first proved in the case $A = B$ by Fuglede [7] and then a proof in the general case was given by Putnam [8]. Berberian [9] proved that the Fuglede theorem was actually equivalent to that of Putnam by a nice operator matrix derivation trick. Rosenblum [10] gave an elegant and simple proof of the Fuglede-Putnam theorem by using Liouville's theorem. There were various generalizations of the Fuglede-Putnam theorem to nonnormal operators; we only cite [11–14]. For example, Radjabalipour [13] showed that the Fuglede-Putnam theorem holds for hyponormal operators; Uchiyama and Tanahashi [14] showed that the Fuglede-Putnam theorem holds for p -hyponormal and log-hyponormal operators. If let $X \in B(\mathcal{H})$ be Hilbert-Schmidt class, Mecheri and Uchiyama [15] showed that normality in the Fuglede-Putnam theorem can be replaced by A and B^* class A operators. In this paper, we show that if X is a Hilbert-Schmidt operator, A and $(B^*)^{-1}$ are $*$ -class A operators such that $AX = XB$, then $A^*X = XB^*$.

Let $\mathcal{C}_2(\mathcal{H})$ denote the Hilbert-Schmidt class. For each pair of operators $A, B \in B(\mathcal{H})$, there is an operator $\Gamma_{A,B}$ defined on $\mathcal{C}_2(\mathcal{H})$ via the formula $\Gamma_{A,B}(X) = AXB$ in [11]. Obviously, $\|\Gamma_{A,B}\| \leq \|A\|\|B\|$. The adjoint of $\Gamma_{A,B}$ is given by the formula $\Gamma_{A,B}^*(X) = A^*XB^*$; see details [11].

Let $A \otimes B$ denote the tensor product on the product space $\mathcal{H} \otimes \mathcal{H}$ for non-zero $A, B \in B(\mathcal{H})$. In [5], Duggal *et al.* give a necessary and sufficient condition for $A \otimes B$ to be a $*$ -class A operator.

Lemma 3.2 (see [5]) *Let $A, B \in B(\mathcal{H})$ be non-zero operators. Then $A \otimes B$ belongs to $*$ -class A operators if and only if A and B belong to $*$ -class A operators.*

Theorem 3.3 *Let A and $B \in B(\mathcal{H})$. Then $\Gamma_{A,B}$ is a $*$ -class A operator on $\mathcal{C}_2(\mathcal{H})$ if and only if A and B^* belong to $*$ -class A operators.*

Proof The unitary operator $U : \mathcal{C}_2(\mathcal{H}) \rightarrow \mathcal{H} \otimes \mathcal{H}$ by a map $(x \otimes y)^* \rightarrow x \otimes y$ induces the $*$ -isomorphism $\Psi : B(\mathcal{C}_2(\mathcal{H})) \rightarrow B(\mathcal{H} \otimes \mathcal{H})$ by a map $X \rightarrow UXU^*$. Then we can obtain $\Psi(\Gamma_{A,B}) = A \otimes B^*$; see details [16]. This completes the proof by Lemma 3.2. \square

Lemma 3.4 (see [17]) *Let $T \in B(\mathcal{H})$ be a $*$ -class A operator. If $\lambda \neq 0$ and $(T - \lambda)x = 0$ for some $x \in \mathcal{H}$, then $(T - \lambda)^*x = 0$.*

Now we are ready to extend the Fuglede-Putnam theorem to $*$ -class A operators.

Theorem 3.5 *Let A and $(B^*)^{-1}$ be $*$ -class A operators. If $AX = XB$ for $X \in \mathcal{C}_2(\mathcal{H})$, then $A^*X = XB^*$.*

Proof Let Γ be defined on $C_2(\mathcal{H})$ by $\Gamma Y = AYB^{-1}$. Since A and $(B^{-1})^* = (B^*)^{-1}$ are $*$ -class A operators, we have that Γ is a $*$ -class A operator on $C_2(\mathcal{H})$ by Theorem 3.3. Moreover, we have $\Gamma X = AXB^{-1} = X$ because of $AX = XB$. Hence X is an eigenvector of Γ . By Lemma 3.4 we have $\Gamma^*X = A^*X(B^{-1})^* = X$, that is, $A^*X = XB^*$. The proof is complete. \square

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the writing of the present article. And they also read and approved the final manuscript.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (11071188); the Natural Science Foundation of the Department of Education, Henan Province (2011A110009), Project of Science and Technology Department of Henan province (122300410375) and Key Scientific and Technological Project of Henan Province (122102210132).

Received: 11 January 2013 Accepted: 25 April 2013 Published: 13 May 2013

References

- Aluthge, A: On p -hyponormal operators for $0 < p < 1$. *Integral Equ. Oper. Theory* **13**, 307-315 (1990)
- Furuta, T: On the class of paranormal operators. *Proc. Jpn. Acad.* **43**, 594-598 (1967)
- Furuta, T: *Invitation to Linear Operators*. Taylor & Francis, London (2001)
- Furuta, T, Ito, M, Yamazaki, T: A subclass of paranormal operators including class of log-hyponormal and several classes. *Sci. Math.* **1**(3), 389-403 (1998)
- Duggal, BP, Jeon, IH, Kim, IH: On $*$ -paranormal contractions and properties for $*$ -class A operators. *Linear Algebra Appl.* **436**, 954-962 (2012)
- Kubrusly, CS, Levan, N: Proper contractions and invariant subspace. *Int. J. Math. Math. Sci.* **28**, 223-230 (2001)
- Fuglede, B: A commutativity theorem for normal operators. *Proc. Natl. Acad. Sci. USA* **36**, 35-40 (1950)
- Putnam, CR: On normal operators in Hilbert space. *Am. J. Math.* **73**, 357-362 (1951)
- Berberian, SK: Note on a theorem of Fuglede and Putnam. *Proc. Am. Math. Soc.* **10**, 175-182 (1959)
- Rosenblum, M: On a theorem of Fuglede and Putnam. *J. Lond. Math. Soc.* **33**, 376-377 (1958)
- Berberian, SK: Extensions of a theorem of Fuglede-Putnam. *Proc. Am. Math. Soc.* **71**, 113-114 (1978)
- Furuta, T: On relaxation of normality in the of Fuglede-Putnam theorem. *Proc. Am. Math. Soc.* **77**, 324-328 (1979)
- Radjabalipour, M: An extension of Fuglede-Putnam theorem for hyponormal operators. *Math. Z.* **194**, 117-120 (1987)
- Uchiyama, A, Tanahashi, K: Fuglede-Putnam's theorem for p -hyponormal or log-hyponormal operators. *Glasg. Math. J.* **44**, 397-410 (2002)
- Mecheri, S, Uchiyama, A: An extension of the Fuglede-Putnam's theorem to class A operators. *Math. Inequal. Appl.* **13**(1), 57-61 (2010)
- Brown, A, Pearcy, C: Spectra of tensor products of operators. *Proc. Am. Math. Soc.* **17**, 162-166 (1966)
- Mecheri, S: Isolated points of spectrum of k -quasi- $*$ -class A operators. *Stud. Math.* **208**, 87-96 (2012)

doi:10.1186/1029-242X-2013-239

Cite this article as: Gao and Li: On $*$ -class A contractions. *Journal of Inequalities and Applications* 2013 **2013**:239.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com