

RESEARCH

Open Access

# Local saturation of a positive linear convolution operator

Hee Sun Jung<sup>1\*</sup> and Ryozi Sakai<sup>2</sup>

\*Correspondence:

hsun90@skku.edu

<sup>1</sup>Department of Mathematics Education, Sungkyunkwan University, Seoul, 110-745, Republic of Korea  
Full list of author information is available at the end of the article

## Abstract

Let  $\{H_n(t)\}$  be a sequence of non-negative, even, and continuous functions on  $\mathbb{R}$ . In this paper, we consider a convolution operator  $J_n(f; x) = \int_0^\infty f(t)H_n(t-x) dt$ ,  $f \in L_p(\mathbb{R}^+)$ , and then investigate the local saturation of  $J_n(f; x)$ .

**MSC:** 44A35

**Keywords:** convolution operator; local saturation

## 1 Introduction and theorems

Through this paper we let  $\{H_n(t)\}$ ,  $n = 1, 2, \dots$ , be a sequence of non-negative, even, and continuous functions on  $\mathbb{R} := (-\infty, \infty)$ , and there exist  $M, N > 0$ , and  $T > 0$  such that  $\{H_n(t)\}$  satisfied uniformly

$$\sup_{|t| \geq T} H_n(t) \leq M, \quad n \geq N, \quad (1.1)$$

$$\int_{-\infty}^{\infty} H_n(t) dt = 1, \quad n = 1, 2, \dots, \quad (1.2)$$

$$\int_{-\infty}^{\infty} t^2 H_n(t) dt = \mu_n \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (1.3)$$

and there exist two positive constants  $\alpha, \beta$  with  $2\beta \geq \alpha > 3$  such that uniformly for  $n$ ,

$$\int_{-\infty}^{\infty} |t|^\alpha H_n(t) dt \leq C \mu_n^\beta. \quad (1.4)$$

As an example for  $H_n(t)$  we give the following:

Let

$$H_n(t) = \frac{n}{\sqrt{\pi}} e^{-(nt)^2}.$$

Then it satisfies (1.2),

$$\mu_n = \frac{1}{n^2} \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad \text{and} \quad \int_{-\infty}^{\infty} t^4 H_n(t) dt = \frac{3}{4} \frac{1}{n^4}.$$

Let us denote  $\mathbb{R}^+ := [0, \infty)$ . In what follows we assume that  $H_n(t)$  satisfies the conditions (1.1)-(1.4). Using  $H_n$ , we define the convolution operators for  $f \in L_p(\mathbb{R}^+)$ ,

$$J_n(f; x) := \int_0^\infty f(t)H_n(t-x) dt, \quad n = 1, 2, \dots \quad (1.5)$$

Swetits and Wood [1] studied the operators on a finite interval  $[0, r]$ :

$$K_n(f; x) := \int_0^r f(t)H_n^*(t-x) dt, \quad f \in L_p([0, r]), n = 1, 2, \dots,$$

where  $H_n^*$  is defined on  $[-r, r]$  as  $H_n$ , and then they gave a local saturation theorem. Furthermore, there is a rich bibliography concerning the convergence of positive linear operators on  $[0, \infty)$  (e.g. see [2–4] and the references cited therein).

In this paper we extend [1] to the infinite interval  $\mathbb{R}^+$ . Then we use a similar methods as [1]. For  $1 < p < \infty$  and  $c \geq 0$ , we define  $L_p^2([c, \infty))$  as the space of those functions  $f$  such that  $f \in L_p(\mathbb{R}^+)$  and  $f'$  is a locally absolutely continuous function on  $[c, \infty)$ , with  $f' \in L_p([c, \infty))$  and  $f'' \in L_p([c, \infty))$ . Let  $C_c^2(\mathbb{R}^+)$  be the space of continuous, compactly supported and continuously second differentiable functions on  $\mathbb{R}^+$ . Furthermore, the total variation of a real-valued function  $f$  defined on an interval  $[a, b] \subset \mathbb{R}$  is the quantity

$$V_a^b(f) = \sup_{P \in \mathcal{P}} \sum_{i=0}^{np-1} |f(x_{i+1}) - f(x_i)|.$$

Here the supremum is taken over the set  $\mathcal{P}$  of all partitions  $P = \{x_0, x_1, \dots, x_{np}\}$  of the interval considered. If  $F : [b, \infty) \rightarrow \mathbb{R}$  and  $x \in [b, \infty)$ , we define

$$T_F(x) = \sup \left\{ \sum_1^n |F(x_j) - F(x_{j-1})| : b = x_0 < x_1 < \dots < x_n = x, n \in \mathbb{N} \right\}.$$

If  $T_F(\infty) := \lim_{x \rightarrow \infty} T_F(x)$  is finite, we say that  $F$  is of bounded variation on  $[b, \infty)$ , and  $T_F(\infty)$  is called the total variation of  $F$  on  $[b, \infty)$ . We define  $\text{BV}[b, \infty)$  to be the set of all functions on  $[b, \infty)$  whose total variation on  $[b, \infty)$  is finite.

Then we first give the pointwise convergence theorem.

**Theorem 1.1** (cf. [5]) *Let  $f \in C_c^2(\mathbb{R}^+)$ , and let  $x \in [a, \infty)$ ,  $a > 0$ . Then we have a pointwise convergence as follows:*

$$\lim_{n \rightarrow \infty} \frac{1}{\mu_n} (J_n(f; x) - f(x)) = \frac{1}{2} f''(x). \quad (1.6)$$

Equation (1.6) holds uniformly on  $[a, \infty)$ .

Then the following is a direct convergence theorem.

**Theorem 1.2** *Let  $0 < b < a < \infty$ .*

- (i) *If  $1 < p < \infty$ , then we have for  $f \in L_p^2([b, \infty))$*

$$\|J_n(f; x) - f(x)\|_{L_p([a, \infty))} = O(\mu_n) \quad \text{as } n \rightarrow \infty. \quad (1.7)$$

(ii) If  $p = 1$ , then we have for  $f' \in \text{BV}([b, \infty))$  with  $f \in L_1([0, \infty))$

$$\|J_n(f; x) - f(x)\|_{L_1([a, \infty))} = O(\mu_n) \quad \text{as } n \rightarrow \infty. \quad (1.8)$$

(iii) Let  $1 \leq p < \infty$  and  $f$  be linear on  $[b, \infty)$ . Then we have

$$\|J_n(f; x) - f(x)\|_{L_p([a, \infty))} = o(\mu_n) \quad \text{as } n \rightarrow \infty. \quad (1.9)$$

Finally, we give an inverse theorem as follows.

**Theorem 1.3** Let  $0 < b < a < \infty$ . Let  $f \in L_p(\mathbb{R}^+)$  and  $f' \in L_p(\mathbb{R}^+)$ .

- (i) For  $1 < p < \infty$ , the condition (1.7) implies  $f \in L_p^2([a, \infty))$ .
- (ii) For  $p = 1$ , the condition (1.8) implies  $f' \in \text{BV}([a, \infty))$ .
- (iii) For  $1 \leq p < \infty$ , the condition (1.9) implies that  $f$  is linear on  $[a, \infty)$ .

This paper is organized as follows. In Section 2, we will give some fundamental lemmas in order to prove the main results and we will prove the results in Section 3.

## 2 Fundamental lemmas

Throughout this paper  $C, C_1, C_2, \dots$  denote positive constants independent of  $n, x, t$  or function  $f(x)$ . The same symbol does not necessarily denote the same constant in different occurrences.

To prove the theorems we need some lemmas.

**Lemma 2.1** Let  $\delta > 0$  and  $\ell = 0, 1, 2$ . Then for  $2\beta \geq \alpha > 3$  defined in (1.4)

$$\int_{|u| \geq \delta} |u|^\ell H_n(u) du \leq C \frac{\mu_n^\beta}{\delta^{\alpha-\ell}}. \quad (2.1)$$

*Proof* Let  $\ell = 0, 1, 2$ . Then we have from (1.4)

$$\begin{aligned} \int_{|u| \geq \delta} |u|^\ell H_n(u) du &= \delta^\ell \int_{|u| \geq \delta} \left( \frac{|u|}{\delta} \right)^\ell H_n(u) du \\ &\leq \delta^\ell \int_{|u| \geq \delta} \left( \frac{|u|}{\delta} \right)^\alpha H_n(u) du \leq \delta^{\ell-\alpha} \int_{|u| \geq \delta} |u|^\alpha H_n(u) du \\ &= \delta^{\ell-\alpha} \int_{-\infty}^{\infty} |u|^\alpha H_n(u) du \leq C \frac{\mu_n^\beta}{\delta^{\alpha-\ell}}. \end{aligned} \quad \square$$

**Lemma 2.2** Let  $e_0(x) := 1$ ,  $e_1(x) := x$  and let  $\delta$  be a positive constant. Then for  $n = 1, 2, \dots$ , we have

$$|J_n((\cdot - x); x)| \leq C \frac{\mu_n^\beta}{\delta^{\alpha-1}}, \quad 0 < \delta \leq x \quad (2.2)$$

and

$$|J_n((\cdot - x)^2; x)| \leq \mu_n, \quad x \geq 0. \quad (2.3)$$

Moreover, we have for  $0 < \delta \leq x$

$$|J_n(e_0; x) - e_0| \leq C \frac{\mu_n^\beta}{\delta^\alpha} \quad \text{and} \quad |J_n(e_1; x) - e_1| \leq C \frac{\mu_n^\beta}{\delta^{\alpha-1}}. \quad (2.4)$$

Here,  $\alpha$  and  $\beta$  are defined in (1.4).

*Proof* For  $0 < \delta \leq x$ , we have by (2.1)

$$\begin{aligned} |J_n((\cdot - x); x)| &= \left| \int_0^\infty (t - x) H_n(t - x) dt \right| = \left| \int_{-x}^\infty y H_n(y) dy \right| \\ &= \left| \int_x^\infty y H_n(y) dy \right| \quad \because H_n(t) \text{ is even} \\ &\leq \int_{y \geq \delta} y H_n(y) dy \leq C \frac{\mu_n^\beta}{\delta^{\alpha-1}}. \end{aligned}$$

For (2.3), we have from (1.3)

$$J_n((\cdot - x)^2; x) = \int_0^\infty (t - x)^2 H_n(t - x) dt = \int_{-x}^\infty y^2 H_n(y) dy \leq \mu_n.$$

Since we know for  $0 < \delta \leq x$

$$J_n(e_0; x) = \int_0^\infty H_n(t - x) dt = \int_{-x}^\infty H_n(y) dy = 1 - \int_{-\infty}^{-x} H_n(y) dy = 1 - \int_x^\infty H_n(y) dy,$$

and from (2.1)

$$\left| \int_x^\infty H_n(y) dy \right| \leq C \frac{\mu_n^\beta}{\delta^\alpha}, \quad (2.5)$$

we have for  $0 < \delta \leq x$ ,

$$|J_n(e_0; x) - e_0| \leq C \frac{\mu_n^\beta}{\delta^\alpha}.$$

Next, we give an estimate for  $e_1$ . Since

$$\begin{aligned} J_n(e_1; x) &= \int_0^\infty t H_n(t - x) dt = \int_{-x}^\infty (x + y) H_n(y) dy \\ &= x \left( 1 - \int_{-\infty}^{-x} H_n(y) dy \right) + \int_x^\infty y H_n(y) dy \\ &= x - x \int_x^\infty H_n(y) dy + \int_x^\infty y H_n(y) dy \end{aligned}$$

and by (2.1), for  $0 < \delta \leq x$ ,

$$\left| x \int_x^\infty H_n(y) dy \right| + \left| \int_x^\infty y H_n(y) dy \right| \leq 2 \int_x^\infty y H_n(y) dy \leq C \frac{\mu_n^\beta}{\delta^{\alpha-1}},$$

we have for  $0 < \delta \leq x$ ,

$$|J_n(e_1; x) - e_1| \leq C \frac{\mu_n^\beta}{\delta^{\alpha-1}}.$$

□

Let  $b > 0$  and then we define

$$\chi([0, b]; t) := \begin{cases} 1, & t \in [0, b], \\ 0, & t \notin [0, b]. \end{cases}$$

**Lemma 2.3** Let  $0 < b < a$ ,  $\delta := a - b$  and let  $1 \leq p < \infty$ . If  $f \in L_p(\mathbb{R}^+)$ , then

$$\|J_n(\chi([0, b])f)\|_{L_p[a, \infty)} \leq C \frac{\mu_n^\beta}{\delta^\alpha} \|f\|_{L_p(\mathbb{R}^+)}.$$

*Proof* Let  $p = 1$ . Then we have

$$\begin{aligned} \|J_n(\chi([0, b])f)\|_{L_1([a, \infty))} &= \int_a^\infty \left| \int_0^\infty \chi([0, b]; t) f(t) H_n(t-x) dt \right| dx \\ &\leq \int_a^\infty \int_0^b |f(t)| H_n(t-x) dt dx \\ &\leq \int_0^b |f(t)| \int_a^\infty H_n(t-x) dx dt \\ &\leq \left( \sup_{t \in [0, b]} \int_a^\infty H_n(t-x) dx \right) \|f\|_{L_1(\mathbb{R}^+)}. \end{aligned}$$

For  $t \in [0, b]$  and  $x \geq a$ , we see  $|t-x| \geq a-b =: \delta > 0$ . From (1.4), we have the following:

$$\begin{aligned} \sup_{t \in [0, b]} \int_a^\infty H_n(t-x) dx &\leq \sup_{t \in [0, b]} \int_a^\infty \left( \frac{|t-x|}{\delta} \right)^\alpha H_n(t-x) dx \\ &\leq \frac{1}{\delta^\alpha} \int_{-\infty}^\infty |t|^\alpha H_n(t) dt \leq C \frac{\mu_n^\beta}{\delta^\alpha}. \end{aligned} \tag{2.6}$$

Hence we have

$$\|J_n(\chi([0, b])f)\|_{L_1[a, \infty)} \leq C \frac{\mu_n^\beta}{\delta^\alpha} \|f\|_{L_1(\mathbb{R}^+)}.$$

Let  $1 < p < \infty$ ,  $0 < b < a \leq x$  and  $p+q=pq$ . By Hölder's inequality,

$$\begin{aligned} |J_n(\chi([0, b])f, x)| &= \left| \int_0^\infty \chi([0, b]; t) f(t) H_n(t-x) dt \right| \\ &\leq C \left( \int_0^\infty \chi([0, b]; t) H_n(t-x) dt \right)^{1/q} \\ &\quad \times \left( \int_0^\infty \chi([0, b]; t) |f(t)|^p H_n(t-x) dt \right)^{1/p}. \end{aligned}$$

Here, from (2.1) we have

$$\begin{aligned} \int_0^\infty \chi([0, b]; t) H_n(t-x) dt &= \int_0^b H_n(t-x) dt = \int_{x-b}^x H_n(y) dy \\ &\leq \int_\delta^\infty H_n(y) dy \leq C \frac{\mu_n^\beta}{\delta^\alpha}. \end{aligned} \quad (2.7)$$

Hence,

$$\begin{aligned} \|J_n(\chi([0, b])f)\|_{L_p[a, \infty)} &\leq C \left( \frac{\mu_n^\beta}{\delta^\alpha} \right)^{1/q} \left( \int_a^\infty \int_0^\infty \chi([0, b]; t) |f(t)|^p H_n(t-x) dt dx \right)^{1/p} \\ &\leq C \left( \frac{\mu_n^\beta}{\delta^\alpha} \right)^{1/q} \left( \int_a^\infty \int_0^\infty \chi([0, b]; t) |f(t)|^p H_n(t-x) dt dx \right)^{1/p}. \end{aligned} \quad (2.8)$$

Also, we see by (2.6)

$$\begin{aligned} &\int_a^\infty \int_0^\infty \chi([0, b]; t) |f(t)|^p H_n(t-x) dt dx \\ &\leq \int_0^\infty \int_a^\infty \chi([0, b]; t) |f(t)|^p H_n(t-x) dx dt \\ &\leq \sup_{t \in [0, b]} \int_a^\infty H_n(t-x) dx \int_0^\infty |f(t)|^p dt \leq C \|f\|_{L_p(\mathbb{R}^+)}^p \frac{\mu_n^\beta}{\delta^\alpha}. \end{aligned}$$

Thus, by (2.8) we conclude.  $\square$

### 3 Proof of theorems

*Proof of Theorem 1.1* For  $x \in [a, \infty)$  and  $t \geq 0$ , we set

$$f(t) = f(x) + f'(x)(t-x) + \frac{f''(\eta)}{2}(t-x)^2, \quad x \leq \eta \leq t.$$

Then we see

$$\begin{aligned} J_n(f; x) - f(x) - \frac{1}{2} f''(x) \mu_n &= f(x) (J_n(e_0; x) - e_0) + f'(x) \int_0^\infty (t-x) H_n(t-x) dt \\ &\quad + \frac{1}{2} \left[ \int_0^\infty f''(\eta) (t-x)^2 H_n(t-x) dt - f''(x) \mu_n \right], \quad x \leq \eta \leq t \\ &:= J_1 + J_2 + J_3. \end{aligned}$$

For  $J_1$ , we have from (2.4)

$$|J_1| = |f(x)| O\left(\frac{\mu_n^\beta}{a^\alpha}\right),$$

and for  $J_2$ , we have from (2.2)

$$|J_2| = |f'(x) J_n((\cdot-x); x)| = |f'(x)| O\left(\frac{\mu_n^\beta}{a^{\alpha-1}}\right).$$

Now, we will estimate  $J_3$ . We have

$$\begin{aligned} & \int_0^\infty f''(\eta)(t-x)^2 H_n(t-x) dt - f''(x)\mu_n \\ &= \int_0^\infty (f''(\eta) - f''(x))(t-x)^2 H_n(t-x) dt - f''(x) \int_{-\infty}^0 (t-x)^2 H_n(t-x) dt. \end{aligned}$$

For the second term, we have by (2.1)

$$\left| f''(x) \int_{-\infty}^0 (t-x)^2 H_n(t-x) dt \right| \leq |f''(x)| \int_a^\infty u^2 H_n(u) du \leq C |f''(x)| \frac{\mu_n^\beta}{\alpha^{\alpha-2}}.$$

For a given  $\varepsilon > 0$ , there exists a positive constant  $\delta_1 > 0$  (depending only on  $\varepsilon$ ) such that for  $|x - \eta| < \delta_1$

$$|f''(x) - f''(\eta)| < \varepsilon. \quad (3.1)$$

For the first term, we have using (1.3)

$$\begin{aligned} & \left| \int_{|x-t| \leq \delta_1} (f''(\eta) - f''(x))(t-x)^2 H_n(t-x) dt \right| \\ & \leq \varepsilon \int_{|u| \leq \delta_1} u^2 H_n(u) du \leq \varepsilon \mu_n, \end{aligned}$$

and by (2.1)

$$\begin{aligned} & \left| \int_{|x-t| \geq \delta_1} (f''(\eta) - f''(x))(t-x)^2 H_n(t-x) dt \right| \\ & \leq 2 \|f''\|_{L_\infty(\mathbb{R})} \int_{|u| \geq \delta_1} u^2 H_n(u) du \leq \|f''\|_{L_\infty(\mathbb{R})} O\left(\frac{\mu_n^\beta}{\delta_1^{\alpha-2}}\right). \end{aligned}$$

Then we have for some positive constants  $C_1$ ,  $C_2$ , and  $C_3$

$$\begin{aligned} \left| J_n(f; x) - f(x) - \frac{1}{2} f''(x) \mu_n \right| & \leq C_1 |f(x)| \frac{\mu_n^\beta}{\alpha^\alpha} + C_2 |f'(x)| \frac{\mu_n^\beta}{\alpha^{\alpha-1}} \\ & + \varepsilon \mu_n + C_3 \|f''\|_{L_\infty(\mathbb{R})} \frac{\mu_n^\beta}{\delta_1^{\alpha-2}}. \end{aligned}$$

Therefore, we have for an arbitrary  $\varepsilon > 0$

$$\lim_{n \rightarrow \infty} \frac{1}{\mu_n} \left| J_n(f; x) - f(x) - \frac{1}{2} f''(x) \mu_n \right| \leq \varepsilon.$$

Thus, (1.6) is proved. Moreover, noting (3.1), we see that (1.6) holds uniformly on  $[a, \infty)$ .  $\square$

*Proof of Theorem 1.2* Let  $\delta := \alpha - b > 0$ . (i) We start under the condition  $1 \leq p < \infty$  for the convenience of considering (ii). On the way of the proof we switch over to the assumption

with  $1 < p < \infty$ . Let  $1 < p < \infty$  and let  $\chi := \chi([0, b])$  and let  $\chi_1(t) = 1 - \chi(t)$ . Since

$$f(t) = (\chi f)(t) + (\chi_1 f)(t), \quad f(x) = (\chi_1 f)(x), \quad x \in [a, \infty),$$

we have

$$\|J_n(f) - f\|_{L_p([a, \infty))} \leq \|J_n(\chi f)\|_{L_p([a, \infty))} + \|J_n(\chi_1 f) - \chi_1 f\|_{L_p([a, \infty))} =: K_1 + K_2. \quad (3.2)$$

By Lemma 2.3,

$$K_1 = \|J_n(\chi f)\|_{L_p([a, \infty))} \leq C \|f\|_{L_p(\mathbb{R}^+)} \frac{\mu_n^\beta}{\delta^\alpha}. \quad (3.3)$$

We estimate  $K_2$ . Let  $f \in L_p^2([a, \infty))$ . Then

$$\begin{aligned} J &:= \|J_n((\chi_1 f)(t) - (\chi_1 f)(x), x)\|_{L_p([a, \infty))} \\ &= \left( \int_a^\infty \left| \int_0^\infty (\chi_1(t)f(t) - f(x)) H_n(t-x) dt \right|^p dx \right)^{1/p} \\ &\leq \left( \int_a^\infty \left| \left( \int_0^b + \int_b^\infty \right) (\chi_1(t)f(t) - f(x)) H_n(t-x) dt \right|^p dx \right)^{1/p} \\ &\leq \left( \int_a^\infty |f(x)|^p \left| \int_0^b H_n(t-x) dt \right|^p dx \right)^{1/p} \\ &\quad + \left( \int_a^\infty \left| \int_b^\infty (f(t) - f(x)) H_n(t-x) dt \right|^p dx \right)^{1/p} \\ &=: I_1 + I_2. \end{aligned} \quad (3.4)$$

Here, for the first term, using

$$\int_0^b H_n(t-x) dt \leq C \frac{\mu_n^\beta}{\delta^\alpha},$$

which is shown in (2.7), we have

$$I_1 = \left( \int_a^\infty |f(x)|^p \left| \int_0^b H_n(t-x) dt \right|^p dx \right)^{1/p} \leq C \|f\|_{L_p([a, \infty))} \frac{\mu_n^\beta}{\delta^\alpha}. \quad (3.5)$$

From this we suppose  $1 < p < \infty$ . By

$$f(t) - f(x) = f'(x)(t-x) + \int_x^t (t-u) f''(u) du,$$

we have

$$\begin{aligned} I_2 &= \left( \int_a^\infty \left| \int_b^\infty (f(t) - f(x)) H_n(t-x) dt \right|^p dx \right)^{1/p} \\ &\leq \left( \int_a^\infty \left| \int_b^\infty f'(x)(t-x) H_n(t-x) dt \right|^p dx \right)^{1/p} \end{aligned}$$

$$+ \left( \int_a^\infty \left| \int_b^\infty \int_x^t (t-u) f''(u) du H_n(t-x) dt \right|^p dx \right)^{1/p} \\ =: I_{2,1} + I_{2,2}.$$

Here we note that  $yH_n(y)$  is an odd function. Then we have by (2.1)

$$\left| \int_b^\infty (t-x) H_n(t-x) dt \right| = \left| \int_{b-x}^\infty y H_n(y) dy \right| = \int_{x-b}^\infty y H_n(y) dy \\ \leq \int_{y \geq \delta} y H_n(y) dy \leq C \frac{\mu_n^\beta}{\delta^{\alpha-1}}$$

and the first term  $I_{2,1}$  is estimated as

$$I_{2,1} = \left( \int_a^\infty \left( |f'(x)| \left| \int_b^\infty (t-x) H_n(t-x) dt \right|^p \right)^{1/p} dx \right)^{1/p} \\ \leq C \frac{\mu_n^\beta}{\delta^{\alpha-1}} \left( \int_a^\infty |f'(x)|^p dx \right)^{1/p} \leq C \|f'\|_{L_p([a,\infty))} \frac{\mu_n^\beta}{\delta^{\alpha-1}}. \quad (3.6)$$

Now we set

$$\theta(f'', x) := \sup_{b \leq t, t \neq x} \frac{1}{t-x} \int_x^t |f''(u)| du, \quad a \leq x,$$

and denote the Hardy-Littlewood majorant of  $f''$  at  $x$ . Since  $f'' \in L_p([b, \infty))$  and  $1 < p < \infty$ , we have

$$\|\theta(f'', x)\|_{L_p([a,\infty))} \leq A_p \|f''\|_{L_p([b,\infty))}, \quad (3.7)$$

where  $A_p > 0$  depend only on  $p$  [6, Theorem 1, p.201]. Since

$$\left| \int_x^t (t-u) f''(u) du \right| \leq \pm \int_x^t |t-u| |f''(u)| du \leq (t-x)^2 \frac{1}{t-x} \int_x^t |f''(u)| du,$$

where  $\pm \int_x^t du \geq 0$ , we have by (3.7) and (1.3)

$$I_{2,2} \leq \left( \int_a^\infty \left| \int_b^\infty \left( (t-x)^2 \frac{1}{t-x} \int_x^t |f''(u)| du \right) H_n(t-x) dt \right|^p dx \right)^{1/p} \\ \leq \left( \int_a^\infty |\theta(f'', x)|^p \left| \int_b^\infty (t-x)^2 H_n(t-x) dt \right|^p dx \right)^{1/p} \\ \leq \int_{-\infty}^\infty u^2 H_n(u) du \left( \int_a^\infty |\theta(f'', x)|^p dx \right)^{1/p} \\ \leq A_p \mu_n \|f''\|_{L_p([b,\infty))}. \quad (3.8)$$

Hence, from (3.6) and (3.8), we have for some positive constant  $C > 0$

$$I_2 \leq I_{2,1} + I_{2,2} \leq C (\|f'\|_{L_p([a,\infty))} + \|f''\|_{L_p([b,\infty))}) \mu_n. \quad (3.9)$$

So, from (3.5) and (3.9),

$$J \leq I_1 + I_2 \leq C(\|f\|_{L_p(\mathbb{R}^+)} + \|f'\|_{L_p([a,\infty))} + \|f''\|_{L_p([b,\infty))})\mu_n. \quad (3.10)$$

Now, we see by (3.10) and (2.4)

$$\begin{aligned} K_2 &= \|J_n(\chi_1 f) - (\chi_1 f)\|_{L_p([a,\infty))} \\ &\leq \|J_n((\chi_1 f)(t) - (\chi_1 f)(x), x)\|_{L_p([a,\infty))} + \|(\chi_1 f)(x)(J_n(e_0, x) - e_0)\|_{L_p([a,\infty))} \\ &= J + \|(\chi_1 f)(x)(J_n(e_0, x) - e_0)\|_{L_p([a,\infty))} \\ &\leq C(\|f\|_{L_p(\mathbb{R}^+)} + \|f'\|_{L_p([a,\infty))} + \|f''\|_{L_p([b,\infty))})\mu_n \\ &\quad + \|J_n(e_0, x) - e_0\|_{L_\infty([a,\infty))}\|\chi_1 f\|_{L_p([a,\infty))} \\ &\leq C_1(\|f\|_{L_p(\mathbb{R}^+)} + \|f'\|_{L_p([a,\infty))} + \|f''\|_{L_p([b,\infty))})\mu_n = O(\mu_n). \end{aligned}$$

Consequently, with (3.2) and (3.3) we conclude (i).

(ii) Let  $p = 1$  and  $f' \in \text{BV}[b, \infty)$ . Then we have for  $x, t \in [b, \infty)$ ,

$$f(t) - f(x) = f'(x)(t - x) + \int_x^t (t - u) df'(u). \quad (3.11)$$

On the proof of (i) we recall the part which we assumed as  $p = 1$ . From (3.3) and (3.5) we may only estimate

$$I'_2 := \int_a^\infty \left| \int_b^\infty (f(t) - f(x)) H_n(t - x) dt dx \right| \quad (3.12)$$

(see (3.4)). By (3.11) we see

$$\begin{aligned} I'_2 &\leq \left| \int_a^\infty \int_b^\infty f'(x)(t - x) H_n(t - x) dt dx \right| \\ &\quad + \left| \int_a^\infty \int_b^\infty \int_x^t (t - u) df'(u) H_n(t - x) dt dx \right| \\ &=: I'_{2,1} + I'_{2,2}. \end{aligned} \quad (3.13)$$

Now, we see that

$$\begin{aligned} I'_{2,1} &= \left| \int_a^\infty \int_b^\infty f'(x)(t - x) H_n(t - x) dt dx \right| \\ &= \left| \int_a^\infty f'(x) \int_{x-b}^\infty u H_n(u) du dx \right| \\ &= \left| \int_{a-b}^\infty u H_n(u) \int_a^{u+b} f'(x) dx du \right| \\ &\leq \sup_{a \leq x < \infty} |f'(x)| \left| \int_\delta^\infty (u - \delta) u H_n(u) du \right| \\ &\leq \sup_{[b, \infty)} |f'(x)| O(\mu_n) \end{aligned}$$

by (1.3). We estimate  $I'_{2,2}$ . We fix an arbitrary  $\eta > 0$ . Then we have by means of the substitution  $u = y + x$  with a new variable  $y$

$$\begin{aligned} I'_{2,2} &= \left| \int_a^\infty \int_b^\infty \int_x^t (t-u) df'(u) H_n(t-x) dt dx \right| \\ &= \left| \int_a^\infty \int_b^\infty \int_0^{t-x} (t-x-y) df'(x+y) H_n(t-x) dt dx \right| \\ &\leq 2 \int_a^\infty \int_b^\infty \int_0^{|t-x|} |t-x| |df'(x+y)| H_n(t-x) dt dx \quad \because |t-x-y| \leq 2|t-x| \\ &\leq 2 \int_a^\infty \sum_{j=0}^\infty \int_{j\eta \leq |t-x| \leq (j+1)\eta} \int_0^{(j+1)\eta} |df'(x+y)| |t-x| H_n(t-x) dt dx \\ &\leq 2 \sum_{j=0}^\infty \int_{j\eta \leq |u| \leq (j+1)\eta} |u| H_n(u) du \int_a^\infty \int_0^{(j+1)\eta} |df'(x+y)| dx. \end{aligned}$$

Then

$$\begin{aligned} &\int_a^\infty \int_0^{(j+1)\eta} |df'(x+y)| dx \\ &= \int_a^\infty \int_x^{x+(j+1)\eta} |df'(v)| dx \\ &= \int_{a \leq v \leq a+(j+1)\eta} \int_{a \leq x \leq v} dx |df'(v)| + \int_{a+(j+1)\eta \leq v < \infty} \int_{v-(j+1)\eta \leq x \leq v} dx |df'(v)| \\ &:= A_1 + A_2. \end{aligned}$$

Since we can see

$$A_1 = \int_{a \leq v \leq a+(j+1)\eta} \int_{a \leq x \leq v} dx |df'(v)| = \int_{a \leq v \leq a+(j+1)\eta} (v-a) |df'(v)| \leq (j+1)\eta \text{BV}(f'; \mathbb{R}^+)$$

and

$$A_2 = \int_{a+(j+1)\eta \leq v < \infty} \int_{v-(j+1)\eta \leq x \leq v} dx |df'(v)| \leq (j+1)\eta \text{BV}(f'; \mathbb{R}^+),$$

we have

$$\int_a^\infty \int_0^{(j+1)\eta} |df'(x+y)| dx \leq 2(j+1)\eta \text{BV}(f'; \mathbb{R}^+).$$

Now, we estimate  $\int_{j\eta \leq |u| \leq (j+1)\eta} |u| H_n(u) du$  for a non-negative integer  $j$ . Let  $j = 0$ . Then from (1.2) and (1.3), we have

$$\int_{0 \leq |u| \leq \eta} |u| H_n(u) du \leq \left( \int_{0 \leq |u| \leq \eta} H_n(u) du \right)^{1/2} \left( \int_{0 \leq |u| \leq \eta} u^2 H_n(u) du \right)^{1/2} = O(\mu_n^{1/2}).$$

Let  $j \geq 1$ . Then by (2.1) we have

$$\int_{j\eta \leq |u| \leq (j+1)\eta} |u| H_n(u) du \leq C \frac{\mu_n^\beta}{(j\eta)^{\alpha-1}}.$$

Therefore, we have

$$\begin{aligned}
 I'_{2,2} &\leq \int_{0 \leq |u| \leq \eta} |u| H_n(u) du \int_a^\infty \int_0^\eta |df'(x+y)| dx \\
 &\quad + \sum_{j=1}^{\infty} \int_{j\eta \leq |u| \leq (j+1)\eta} |u| H_n(u) du \int_a^\infty \int_0^{(j+1)\eta} |df'(x+y)| dx \\
 &\leq O(\mu_n^{1/2}) \eta \text{BV}(f'; \mathbb{R}^+) + \sum_{j=1}^{\infty} \frac{1}{(j\eta)^{\alpha-1}} O(\mu_n^\beta) (j+1) \eta \text{BV}(f'; \mathbb{R}^+) \\
 &= O(\mu_n^{1/2}) \eta \text{BV}(f'; \mathbb{R}^+) + O(\mu_n^\beta) \frac{1}{\eta^{\alpha-2}} \text{BV}(f'; \mathbb{R}^+) \quad (\because \alpha > 3).
 \end{aligned}$$

If we let  $\eta = \mu_n^{1/2}$ , then we have  $I'_{2,2} = O(\mu_n) \text{BV}(f'; \mathbb{R}^+)$ , because  $\beta - (\alpha - 2)/2 \geq 1$ . Consequently, (ii) is proved.

(iii) It follows from (2.4). Consequently, for a linear function  $f$  on  $[b, \infty)$

$$\|J_n(f; x) - f(x)\|_{L_p([a, \infty))} = O(\mu_n^\beta) = o(\mu_n).$$

□

*Proof of Theorem 1.3* (i), (ii) hold as follows: Let  $f \in L_p(\mathbb{R}^+)$ . First, we choose  $\psi \in C^2(\mathbb{R}^+)$  with  $\psi'' \in L_q([a, \infty))$  (for any  $1 \leq q \leq \infty$ ) such that

$$\psi(a) = \psi'(a) = \psi''(a) = 0, \quad \lim_{r \rightarrow \infty} \psi^{(i)}(r) = 0, \quad i = 0, 1, 2$$

and

$$\psi(x) = 0 \quad \text{for } x \notin [a, \infty).$$

We use the bilinear functional;

$$A_n(f, \psi) = \frac{1}{\mu_n} \int_0^\infty (J_n(f, x) - f(x)) \psi(x) dx.$$

We will show that for fixed  $\psi$ ,  $A_n(\cdot, \psi)$  is uniformly bounded on  $L_p(\mathbb{R}^+)$ . We see

$$\int_0^\infty J_n(f, x) \psi(x) dx = \int_0^\infty \psi(x) \int_0^\infty f(t) H_n(t-x) dt dx.$$

For  $x, t \in [0, \infty)$  we can write

$$\psi(x) = \psi(t) + \psi'(t)(x-t) + \int_x^t (x-u) \psi''(u) du.$$

Hence,

$$\begin{aligned}
 &\int_0^\infty \int_0^\infty f(t) \psi(x) H_n(t-x) dx dt \\
 &= \int_0^\infty f(t) \psi(t) \int_0^\infty H_n(t-x) dx dt
 \end{aligned}$$

$$\begin{aligned}
 & + \int_0^\infty f(t) \psi'(t) \int_0^\infty (x-t) H_n(t-x) dx dt \\
 & + \int_0^\infty f(t) \int_0^\infty \int_x^t (x-u) \psi''(u) du H_n(t-x) dx dt \\
 & = I_1'' + I_2'' + I_3''.
 \end{aligned}$$

From (1.2)

$$I_1'' = \int_0^\infty f(t) \psi(t) \int_{-\infty}^t H_n(u) du dt = \int_0^\infty f(t) \psi(t) \left( 1 - \int_t^\infty H_n(u) du \right) dt.$$

Since  $\psi(t) = 0$  for  $t \notin [a, \infty)$  (so, we may take  $t \geq a$ ), by (2.1) we have

$$\begin{aligned}
 & \left| \int_0^\infty f(t) \psi(t) \int_t^\infty H_n(u) du dt \right| \\
 & \leq \left| \int_0^\infty f(t) \psi(t) \int_t^\infty \left(\frac{u}{t}\right)^\alpha H_n(u) du dt \right| \quad \because (u/t)^\alpha \geq 1 \\
 & \leq O(\mu_n^\beta) \int_a^\infty |f(t)\psi(t)| \frac{1}{t^\alpha} dt \quad \text{by (1.4)} \\
 & \leq O(\mu_n^\beta) \sup_{a \leq t < \infty} |\psi(t)| \int_a^\infty |f(t)| \frac{1}{t^\alpha} dt \\
 & \leq O(\mu_n^\beta) \sup_{a \leq t < \infty} |\psi(t)| \|f\|_{L_p([a, \infty))} \|t^{-\alpha}\|_{L_q([a, \infty))},
 \end{aligned}$$

where  $1/p + 1/q = 1$ . Thus, we have

$$I_1'' = \int_0^\infty f(t) \psi(t) dt + O(\mu_n^\beta) \|f\|_{L_p([a, \infty))}.$$

We estimate  $I_2''$ . We may  $a \leq t$ , because  $\psi(t) = 0$  for  $t \notin [a, \infty)$ . Noting (1.4) and  $\alpha < 3$ ,

$$\begin{aligned}
 |I_2''| &= \int_a^\infty |f(t)\psi'(t)| \int_t^\infty u H_n(u) du dt \\
 &\leq \int_a^\infty |f(t)\psi'(t)| \int_t^\infty \frac{u^\alpha}{t^{\alpha-1}} H_n(u) du dt \\
 &= O(\mu_n^\beta) \int_a^\infty |f(t)\psi'(t)| \frac{1}{t^{\alpha-1}} dt \\
 &= O(\mu_n^\beta) \|\psi'\|_{L_\infty([a, \infty))} \|f\|_{L_p([a, \infty))} \|t^{1-\alpha}\|_{L_q([a, \infty))} \\
 &= O(\mu_n^\beta) \|f\|_{L_p([a, \infty))}.
 \end{aligned}$$

Finally, we estimate  $I_3''$ . Let  $1 < p < \infty$  and  $1/p + 1/q = 1$ . As the estimation for  $I_{2,2}$ , we have by (1.3)

$$\begin{aligned}
 |I_3''| &= \left| \int_0^\infty f(t) \int_0^\infty \int_x^t (x-u) \psi''(u) du H_n(t-x) dx dt \right| \\
 &= \left| \int_0^\infty f(t) \int_0^\infty \frac{1}{t-x} \int_x^t \psi''(u) du (t-x)^2 H_n(t-x) dx dt \right|
 \end{aligned}$$

$$\begin{aligned}
 &\leq \int_a^\infty |f(t)| \theta(\psi'', t) \int_0^\infty (t-x)^2 H_n(t-x) dx dt \\
 &= O(\mu_n) \int_a^\infty |f(t)| \theta(\psi'', t) dt \leq O(\mu_n) \|f\|_{L_p([a, \infty))} \|\psi''\|_{L_q([a, \infty))} \\
 &= O(\mu_n) \|f\|_{L_p([a, \infty))}.
 \end{aligned}$$

We estimate  $I_3''$  for  $p = 1$ . There exists  $\eta$  between  $x$  and  $t$  such that

$$\begin{aligned}
 |I_3''| &\leq \left| \int_0^\infty f(t) \int_0^\infty \psi''(\eta)(t-x)^2 H_n(t-x) dx dt \right| \\
 &\leq \sup_{a \leq t < \infty} |\psi''(t)| \int_a^\infty |f(t)| \int_0^\infty (t-x)^2 H_n(t-x) dx dt \\
 &\leq O(\mu_n) \sup_{a \leq t < \infty} |\psi''(t)| \|f\|_{L_1([a, \infty))} = O(\mu_n) \|f\|_{L_1([a, \infty))}.
 \end{aligned}$$

Here we used (1.3). Consequently, it follows that  $|A_n(f, \psi)|$  is uniformly bounded on  $L_p(\mathbb{R}^+)$ . Next, from Theorem 1.1 we see that for  $f \in C_c^2(\mathbb{R}^+)$ ,

$$\lim_{n \rightarrow \infty} A_n(f, \psi) = \frac{1}{2} \int_a^\infty f''(x) \psi(x) dx = \frac{1}{2} \int_a^\infty f(x) \psi''(x) dx. \quad (3.14)$$

Since  $\{A_n(\cdot, \psi)\}$  is uniformly bounded on  $L_p(\mathbb{R}^+)$ , and  $C_c^2(\mathbb{R}^+)$  is dense in  $L_p(\mathbb{R}^+)$ , (3.14) yields

$$\lim_{n \rightarrow \infty} A_n(f, \psi) = \frac{1}{2} \int_a^\infty f(x) \psi''(x) dx \quad (3.15)$$

for any  $f \in L_p(\mathbb{R}^+)$ . Now, for any fixed  $f \in L_p(\mathbb{R}^+)$ , we consider the sequence of linear functional  $\{A_n(f, \cdot)\}$ . Since  $\|J_n(f) - f\|_{L_p([a, \infty))} = O(\mu_n)$ ,  $n \rightarrow \infty$ , there exist  $h \in L_p([a, \infty))$  ( $p > 1$ ) and  $h \in \text{BV}[a, \infty)$  ( $p = 1$ ) and a subsequence  $\{A_{n_j}(f, \cdot)\}$  such that

$$\lim_{j \rightarrow \infty} A_{n_j}(f, \psi) = \begin{cases} \int_0^\infty h(x) \psi(x) dx, & p > 1, \\ \int_0^\infty \psi(x) dh(x), & p = 1. \end{cases} \quad (3.16)$$

From (3.15) and (3.16) we obtain

$$\frac{1}{2} \int_0^\infty f(x) \psi''(x) dx = \begin{cases} \int_0^\infty h(x) \psi(x) dx, & p > 1, \\ \int_0^\infty \psi(x) dh(x), & p = 1. \end{cases} \quad (3.17)$$

A particular solution to (3.17) is

$$\frac{1}{2} f(x) = \begin{cases} \int_\eta^x \int_\eta^\xi h(\mu) d\mu d\xi, & p > 1, \\ \int_\eta^x \int_\eta^\xi dh(\mu) d\xi, & p = 1. \end{cases}$$

The homogeneous problem

$$\int_a^\infty f(x) \psi''(x) dx = 0$$

has the general solution  $f(x) = C_1x + C_2$  for  $a \leq x < \infty$ , since we can take  $\psi \in C^2([a, \infty))$  arbitrarily as  $\psi'' \in L_q([a, \infty)), \psi(a) = \psi'(a) = 0, \lim_{r \rightarrow \infty} \psi^{(i)}(r) = 0, i = 0, 1$ . Hence, if  $1 < p < \infty, f \in L_p^2([a, \infty)),$  and if  $p = 1$  then  $f' \in BV([a, \infty)).$  Hence, (i) and (ii) hold. We will show (iii). Now, if

$$\|J_n(f) - f\|_{L_p([a, \infty))} = o(\mu_n), \quad n \rightarrow \infty,$$

then

$$\begin{aligned} |A_n(f, \psi)| &\leq \frac{1}{\mu_n} \int_a^\infty |J_n(f, x) - f(x)| |\psi(x)| dx \\ &\leq \left( \sup_{a \leq x < \infty} |\psi(x)| \right) \frac{C_p}{\mu_n} \|J_n(f) - f\|_{L_p([a, \infty))}, \end{aligned}$$

where  $C_p > 0$  is independent of  $n.$  Hence

$$\lim_{n \rightarrow \infty} A_n(f, \psi) = 0. \quad (3.18)$$

Considering (3.15), (3.16), (3.17), and (3.18), we obtain

$$\int_a^\infty f(x) \psi''(x) dx = 0,$$

and so

$$\int_a^\infty f''(x) \psi(x) dx = 0,$$

consequently, we have  $f''(x) = 0$ , that is,  $f$  is linear on  $[a, \infty).$  □

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

All authors conceived of the study, participated in its design and coordination, drafted the manuscript and participated in the sequence alignment. All authors read and approved the final manuscript.

#### Author details

<sup>1</sup>Department of Mathematics Education, Sungkyunkwan University, Seoul, 110-745, Republic of Korea. <sup>2</sup>Department of Mathematics, Meijo University, Nagoya, 468-8502, Japan.

#### Acknowledgements

The authors thank the referees for many valuable comments and corrections.

Received: 27 December 2013 Accepted: 22 August 2014 Published: 02 Sep 2014

#### References

1. Swetits, JJ, Wood, B: Local  $L_p$ -saturation of positive linear convolution operators. *J. Approx. Theory* **34**(4), 348-360 (1982)
2. Swetits, JJ, Wood, B: Approximating functions of exponential type on the positive axis: direct results. *Bull. Inst. Math. Acad. Sin.* **23**(2), 167-180 (1995)
3. Jung, HS, Sakai, R: Approximation by Lupas-type operators and Szász-Mirakyan-type operators. *J. Appl. Math.* **2012**, Article ID 546784 (2012)
4. Deo, N, Jung, HS, Sakai, R: Degree of approximation by hybrid operators. *Abstr. Appl. Anal.* **2013**, Article ID 732069 (2013)
5. Mamedov, RG: The asymptotic value of the approximation of differentiable functions by linear positive operators. *Dokl. Akad. Nauk SSSR* **128**, 471-474 (1959) (Russian)
6. Stein, EM: Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals. Princeton University Press, Princeton (1993)

10.1186/1029-242X-2014-329

**Cite this article as:** Jung and Sakai: Local saturation of a positive linear convolution operator. *Journal of Inequalities and Applications* 2014, 2014:329

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

---