

RESEARCH

Open Access



The matching energy of graphs with given edge connectivity

Shengjin Ji^{1,3}, Hongping Ma^{2*} and Gang Ma¹

*Correspondence: hpma@163.com

²School of Mathematics and Statistics, Jiangsu Normal University, Xuzhou, Jiangsu 221116, China
Full list of author information is available at the end of the article

Abstract

Let G be a simple graph of order n and $\mu_1, \mu_2, \dots, \mu_n$ the roots of its matching polynomial. The matching energy of G is defined as the sum $\sum_{i=1}^n |\mu_i|$. Let $K_{n-1,1}^k$ be the graph obtained from $K_1 \cup K_{n-1}$ by adding k edges between $V(K_1)$ and $V(K_{n-1})$. In this paper, we show that $K_{n-1,1}^k$ has the maximum matching energy among the connected graphs with order n and edge connectivity k .

Keywords: matching energy; edge connectivity; graph energy; matching; extremal graph

1 Introduction

We use Bondy and Murty [1] for terminology and notations not defined in this paper and consider undirected and simple graphs only. Let $G = (V, E)$ be a graph with order n . Denote by $m(G, t)$ the number of t -matchings of G . Clearly, $m(G, 1) = e(G)$, the size of G , and $m(G, t) = 0$ for $t > \lfloor n/2 \rfloor$.

Recall that the *matching polynomial* of a graph G is defined as

$$\alpha(G) = \alpha(G, \lambda) = \sum_{t \geq 0} (-1)^t m(G, t) \lambda^{n-2t}$$

and its theory is well elaborated in [2–4].

The *energy* of G is defined as

$$E(G) = \sum_{i=1}^n |\lambda_i|,$$

where $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of the adjacency matrix $A(G)$ of G . The theory of graph energy is well developed nowadays; for details, see reviews [5, 6], the book [7], and the recent related results [8–23].

The Coulson integral formula [24] plays an important role in the study of the graph energy. For an acyclic graph T , its Coulson integral formula is as follows:

$$E(T) = \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \ln \left[\sum_{t \geq 0} m(T, t) x^{2t} \right] dx. \tag{1}$$

Motivated by equation (1), Gutman and Wagner [25] defined the *matching energy* of a graph G as

$$ME = ME(G) = \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \ln \left[\sum_{t \geq 0} m(G, t)x^{2t} \right] dx. \tag{2}$$

Energy and matching energy of graphs are closely related, and they are two quantities of relevance for chemical applications; for details see [26–28].

We now give an equivalent definition of matching energy.

Definition 1.1 [25] Let G be a graph of order n , and let $\mu_1, \mu_2, \dots, \mu_n$ be the roots of its matching polynomial. Then

$$ME(G) = \sum_{i=1}^n |\mu_i|.$$

The equation (2) induces a *quasi-order* relation over the set of all graphs on n vertices: if G_1 and G_2 are two graphs of order n , then

$$G_1 \leq G_2 \iff m(G_1, t) \leq m(G_2, t) \text{ for all } t = 0, 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor. \tag{3}$$

If $G_1 \leq G_2$ and there exists some i such that $m(G_1, i) < m(G_2, i)$, then we write $G_1 < G_2$. Clearly,

$$G_1 < G_2 \implies ME(G_1) < ME(G_2).$$

Recall that the *Hosoya index* of a graph G is defined as $Z(G) = \sum_{t \geq 0} m(G, t)$ [29]. So we also have

$$G_1 < G_2 \implies Z(G_1) < Z(G_2).$$

In [3, 4], the authors gave two fundamental identities for the number of t -matchings of a graph.


Lemma 1.2 Let G be a graph, $e = uv$ an edge of G , and $N(u) = \{v_1(= v), v_2, \dots, v_j\}$ the set of all neighbors of u in G . Then we have

$$m(G, t) = m(G - uv, t) + m(G - u - v, t - 1) \tag{4}$$

and

$$m(G, t) = m(G - u, t) + \sum_{i=1}^j m(G - u - v_i, t - 1). \tag{5}$$

From Lemma 1.2, it is easy to get the following result.



Lemma 1.3 [25] *Let G be a graph and e one of its edges. Let $G - e$ be the subgraph obtained from G by deleting the edge e . Then $G - e < G$ and $ME(G - e) < ME(G)$.*

By Lemma 1.3, among all graphs on n vertices, the empty graph E_n and the complete graph K_n have the minimum and the maximum matching energy [25], respectively. It follows from equations (1) and (2) that $ME(T) = E(T)$ for any tree T [25]. By using the quasi-order relation, some results have been obtained on extremal graphs with respect to matching energy among some classes of connected graphs with n vertices. For example, the extremal graphs in connected unicyclic, bicyclic graphs were determined in [25] and [30, 31], respectively; the maximal matching energy graphs of tricyclic graphs were obtained in [32], the minimal graphs among connected k -cyclic ($k \leq n - 3$) graphs and bipartite graphs were characterized in [33]; the maximal connected graph with given connectivity (resp. chromatic number) was determined in [34]. For more as regards the latest results on matching energy, we refer the reader to [30, 35–41].

We now introduce some notations which will be used in the next section.

Let $\mathcal{G}_{n,k}$ be the set of connected graphs of order n (≥ 2) with edge connectivity k ($1 \leq k \leq n - 1$). Let $K_{n-1,1}^k$ be the graph obtained from $K_1 \cup K_{n-1}$ by adding k edges between $V(K_1)$ and $V(K_{n-1})$; see Figure 1. In this paper, we show that $K_{n-1,1}^k$ is the unique graph with the maximum matching energy (resp. Hosoya index) in $\mathcal{G}_{n,k}$.

2 Main results

First we recall some notations. We denote by $\kappa'(G)$ and $\delta(G)$ the edge connectivity and the minimum degree of a graph G , respectively. Let S be a nonempty proper subset of V . We use $G[S]$ to denote the subgraph of G induced by S . The *edge cut* of G , denoted by $\partial(S)$, is a subset of $E(G)$ of the form $[S, \bar{S}]$, where $\bar{S} = V \setminus S$. An edge cut $\partial(v)$ ($v \in V$) is called a *trivial edge cut*. A *k -edge cut* is an edge cut of k elements. Let $G \in \mathcal{G}_{n,k}$. Then G must have a k -edge cut $\partial(S)$ with $1 \leq |S| \leq \lfloor \frac{n}{2} \rfloor$.

Lemma 2.1 *Let $G \not\cong K_{n-1,1}^k$ be a graph in $\mathcal{G}_{n,k}$ with a trivial k -edge cut. Then $G < K_{n-1,1}^k$.*

Proof Let $\partial(S)$ be a trivial k -edge cut of G with $|S| = 1$. Since $G \not\cong K_{n-1,1}^k$, $G[\bar{S}]$ is a proper subgraph of K_{n-1} . Hence G is a proper subgraph of $K_{n-1,1}^k$, and so the result follows from Lemma 1.3. □

Lemma 2.2 *Let $G \in \mathcal{G}_{n,k}$ be a graph without trivial k -edge cuts. Then for any k -edge cut $\partial(S)$ of G with $2 \leq |S| \leq \lfloor \frac{n}{2} \rfloor$, we have $|S| \geq k$.*

Proof For $k \leq 2$, the assertion is trivial, so suppose $k \geq 3$. Assume, to the contrary, that G has a k -edge cut $\partial(S)$ with $2 \leq |S| \leq k - 1$. By the facts that $\delta(G) \geq \kappa'(G) = k$ and G has no trivial k -edge cuts, we have $\delta(G) \geq k + 1$, and thus $\sum_{v \in S} d_G(v) \geq |S|(k + 1)$. On the




Figure 2 The graphs G_1 and G_2 of $\mathcal{G}_{n,k}$ in Operation I.

other hand, $\sum_{v \in S} d_G(v) = 2e(G[S]) + k \leq |S|(|S| - 1) + k$. Therefore, we have $|S|(k + 1) \leq |S|(|S| - 1) + k$, that is, $(|S| - 1)(k - |S|) + |S| \leq 0$, which is a contradiction. Therefore the result holds. \square

For $k \leq m \leq \lfloor \frac{n}{2} \rfloor$, let $K_{n-m,m}^k$ be a graph obtained from $K_{n-m} \cup K_m$ by adding k independent edges between $V(K_{n-m})$ and $V(K_m)$, as shown in Figure 1. It is easy to see that $\kappa'(K_{n-m,m}^k) = k$ and $\kappa'(K_{n-1,1}^k) = k$.

We next show that for a graph $G \in \mathcal{G}_{n,k}$ without trivial k -edge cuts, $G \leq K_{n-m,m}^k$ for some m . Before this, we introduce a new graph operation as follows.

Let G_1 be a graph in $\mathcal{G}_{n,k}$ such that G_1 has a k -edge cut $\partial(S)$ with $G[S] = K_m$, $G[\bar{S}] = K_{n-m}$, and $k \leq m \leq \lfloor \frac{n}{2} \rfloor$. Suppose that $u_1, u_2 \in \bar{S}$, $v_1, v_2 \in S$, $e_1 = u_1v_1$, $e_2 = u_1v_2$ are two edges of $\partial(S)$, and u_2 is not incident with any edge in $\partial(S)$. If G_2 is obtained from G_1 by deleting the edge e_2 and adding a new edge $e'_2 = u_2v_2$, we say that G_2 is obtained from G_1 by *Operation I*, as shown in Figure 2. Clearly, $G_2 \in \mathcal{G}_{n,k}$.

Lemma 2.3 *If G_2 is obtained from G_1 by Operation I, then $G_1 < G_2$.*

Proof By equation (4), we have

$$m(G_1, t) = m(G_1 - e_2, t) + m(G_1 - u_1 - v_2, t - 1)$$

and

$$m(G_2, t) = m(G_2 - e'_2, t) + m(G_2 - u_2 - v_2, t - 1).$$

Note that $G_1 - e_2 \cong G_2 - e'_2$, and $G_1 - u_1 - v_2$ is isomorphic to a proper subgraph of $G_2 - u_2 - v_2$. So, $m(G_1 - u_1 - v_2, t - 1) \leq m(G_2 - u_2 - v_2, t - 1)$ for all t and $m(G_1 - u_1 - v_2, 1) < m(G_2 - u_2 - v_2, 1)$. The result thus follows. \square

Lemma 2.4 *Let $G \in \mathcal{G}_{n,k}$ be a graph without trivial k -edge cuts. Then $G \leq K_{n-m,m}^k$ for some m with $\max\{k, 2\} \leq m \leq \lfloor \frac{n}{2} \rfloor$.*

Proof Let $\partial(S)$ be a k -edge cut of G with $2 \leq |S| \leq \lfloor \frac{n}{2} \rfloor$. Let $|S| = m$. Then $m \geq k$ by Lemma 2.2. Let G_1 be the graph obtained from G , by adding edges if necessary, such that $G[S]$ and $G[\bar{S}]$ are complete graphs. Therefore $G \leq G_1$ by Lemma 1.3. If $G_1 \not\leq K_{n-m,m}^k$, then by using Operation I repeatedly, we can finally get $K_{n-m,m}^k$ from G_1 . Hence $G_1 \leq K_{n-m,m}^k$ by Lemma 2.3. The proof is thus complete. \square

In the following, we show that $K_{n-m,m}^k < K_{n-1,1}^k$ for $m \geq 2$.

Lemma 2.5 Suppose $\max\{k, 2\} \leq m \leq \lfloor \frac{n}{2} \rfloor$. Then $e(K_{n-m,m}^k) < e(K_{n-1,1}^k)$.

Proof Note that

$$e(K_{n-m,m}^k) = \frac{m(m-1)}{2} + \frac{(n-m)(n-m-1)}{2} + k$$

and

$$e(K_{n-1,1}^k) = \frac{(n-1)(n-2)}{2} + k.$$

Hence we have

$$\begin{aligned} e(K_{n-1,1}^k) - e(K_{n-m,m}^k) &= \frac{n^2 - 3n + 2}{2} - \frac{n^2 + 2m^2 - 2mn - n}{2} \\ &= (m-1)(n-m-1) > 0. \end{aligned}$$

The proof is thus complete. □

Lemma 2.6 Let $m \geq 1$ be a positive integer. Then we have

$$m(K_{m,m}^1, t) \leq m(K_{2m-1,1}^1, t) \quad \text{for all } t = 0, 1, \dots, m \tag{6}$$

and

$$m(K_{m+1,m}^1, t) \leq m(K_{2m,1}^1, t) \quad \text{for all } t = 0, 1, \dots, m. \tag{7}$$

Proof We apply induction on m . For $m = 1$ and $m = 2$, the assertions are trivial since $K_{2,2}^1$ and $K_{3,2}^1$ are proper subgraphs of $K_{3,1}^1$ and $K_{4,1}^1$, respectively. So suppose that $m \geq 3$ and inequalities (6) and (7) hold for smaller values of m . By Lemma 1.2, we obtain

$$\begin{aligned} m(K_{m,m}^1, t) &= m(K_{m,m-1}^1, t) + (m-2)m(K_{m,m-2}^1, t-1) + m(K_m \cup K_{m-2}, t-1) \\ &= m(K_{m,m-1}^1, t) + (m-1)m(K_{m,m-2}^1, t-1) - m(K_{m-1} \cup K_{m-3}, t-2) \\ &= m(K_{m,m-1}^1, t) - m(K_{m-1} \cup K_{m-3}, t-2) + (m-1)[m(K_{m-1,m-2}^1, t-1) \\ &\quad + (m-1)m(K_{m-2,m-2}^1, t-2) - m(K_{m-3} \cup K_{m-3}, t-3)] \\ &\leq m(K_{m,m-1}^1, t) + (m-1)m(K_{m-1,m-2}^1, t-1) + (m-1)^2m(K_{m-2,m-2}^1, t-2) \\ &\quad - m(K_{m-1} \cup K_{m-3}, t-2) \end{aligned}$$

and

$$\begin{aligned} m(K_{2m-1,1}^1, t) &= m(K_{2m-2,1}^1, t) + (2m-3)m(K_{2m-3,1}^1, t-1) + m(K_{2m-3}, t-1) \\ &= m(K_{2m-2,1}^1, t) + m(K_{2m-3}, t-1) + (2m-3)[m(K_{2m-4,1}^1, t-1) \\ &\quad + (2m-5)m(K_{2m-5,1}^1, t-2) + m(K_{2m-5}, t-2)] \\ &\geq m(K_{2m-2,1}^1, t) + (2m-3)m(K_{2m-4,1}^1, t-1) \\ &\quad + (2m-3)(2m-5)m(K_{2m-5,1}^1, t-2). \end{aligned}$$

By the induction hypothesis, we obtain

$$\begin{aligned} m(K_{m,m-1}^1, t) &\leq m(K_{2m-2,1}^1, t), \\ m(K_{m-1,m-2}^1, t-1) &\leq m(K_{2m-4,1}^1, t-1), \\ m(K_{m-2,m-2}^1, t-2) &\leq m(K_{2m-5,1}^1, t-2). \end{aligned}$$

Since $m \geq 3$, we have $m-1 \leq 2m-3$ and $(m-1)^2 \leq (2m-3)(2m-5)$ when $m \geq 4$. Notice that for $m = 3$, $K_{m-2,m-2}^1 = K_{m-1} \cup K_{m-3}$, and $(m-1)^2 - 1 = (2m-3)(2m-5)$. Hence inequality (6) holds.

By Lemma 1.2, we get

$$\begin{aligned} m(K_{m+1,m}^1, t) &= m(K_{m,m}^1, t) + (m-1)m(K_{m-1,m}^1, t-1) + m(K_{m-1} \cup K_m, t-1) \\ &\leq m(K_{m,m}^1, t) + m \cdot m(K_{m-1,m}^1, t-1) \\ &= m(K_{m,m}^1, t) + m \cdot [m(K_{m-1,m-1}^1, t-1) \\ &\quad + (m-2)m(K_{m-1,m-2}^1, t-2) + m(K_{m-1} \cup K_{m-2}, t-2)] \\ &\leq m(K_{m,m}^1, t) + m \cdot [m(K_{m-1,m-1}^1, t-1) \\ &\quad + (m-1)m(K_{m-1,m-2}^1, t-2)] \\ &= m(K_{m,m}^1, t) + m \cdot m(K_{m-1,m-1}^1, t-1) + m(m-1)m(K_{m-1,m-2}^1, t-2) \end{aligned}$$

and

$$\begin{aligned} m(K_{2m,1}^1, t) &= m(K_{2m-1,1}^1, t) + (2m-2)m(K_{2m-2,1}^1, t-1) + m(K_{2m-2}, t-1) \\ &= m(K_{2m-1,1}^1, t) + m(K_{2m-2}, t-1) + (2m-2)[m(K_{2m-3,1}^1, t-1) \\ &\quad + (2m-4)m(K_{2m-4,1}^1, t-2) + m(K_{2m-4}, t-2)] \\ &\geq m(K_{2m-1,1}^1, t) + (2m-2)m(K_{2m-3,1}^1, t-1) \\ &\quad + (2m-2)(2m-4)m(K_{2m-4,1}^1, t-2). \end{aligned}$$

By the induction hypothesis and inequality (6), we have

$$\begin{aligned} m(K_{m,m}^1, t) &\leq m(K_{2m-1,1}^1, t), \\ m(K_{m-1,m-1}^1, t-1) &\leq m(K_{2m-3,1}^1, t-1), \\ m(K_{m-1,m-2}^1, t-2) &\leq m(K_{2m-4,1}^1, t-2). \end{aligned}$$

Notice that $m \leq 2m-2$ and $m(m-1) \leq (2m-2)(2m-4)$. Therefore inequality (7) holds.

The proof is thus complete. □

Lemma 2.7 Suppose $2 \leq m \leq \lfloor \frac{n}{2} \rfloor$. Then

$$m(K_{n-m,m}^1, t) \leq m(K_{n-1,1}^1, t) \quad \text{for all } t = 0, 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor.$$

Proof We apply induction on n . As the two cases $n = 2m$ and $n = 2m + 1$ were proved by Lemma 2.6, we proceed to the induction step. By Lemma 1.2 and the induction hypothesis, we have

$$\begin{aligned} m(K_{n-m,m}^1, t) &= m(K_{n-m,m-1}^1, t) + (m - 2)m(K_{n-m,m-2}^1, t - 1) + m(K_{n-m} \cup K_{m-2}, t - 1) \\ &\leq m(K_{n-m,m-1}^1, t) + (m - 1)m(K_{n-m,m-2}^1, t - 1) \\ &\leq m(K_{n-2,1}^1, t) + (m - 1)m(K_{n-3,1}^1, t - 1) \\ &= m(K_{n-2}, t) + m(K_{n-3}, t - 1) \\ &\quad + (m - 1)(m(K_{n-3}, t - 1) + m(K_{n-4}, t - 2)) \\ &= m(K_{n-2}, t) + m \cdot m(K_{n-3}, t - 1) + (m - 1)m(K_{n-4}, t - 2) \end{aligned}$$

and

$$\begin{aligned} m(K_{n-1,1}^1, t) &= m(K_{n-1}, t) + m(K_{n-2}, t - 1) \\ &= m(K_{n-2}, t) + (n - 2)m(K_{n-3}, t - 1) \\ &\quad + m(K_{n-3}, t - 1) + (n - 3)m(K_{n-4}, t - 2) \\ &= m(K_{n-2}, t) + (n - 1)m(K_{n-3}, t - 1) + (n - 3)m(K_{n-4}, t - 2). \end{aligned}$$

Thus the result follows by the fact that $m \leq n - 2$. □

Lemma 2.8 *Suppose $k \leq m \leq \lfloor \frac{n}{2} \rfloor$. Then*

$$m(K_{n-m,m}^k, t) \leq m(K_{n-1,1}^k, t) \quad \text{for all } t = 0, 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor.$$

Proof We apply induction on k . As the case $k = 1$ was proved by Lemma 2.7, we suppose that $k \geq 2$ and the assertion holds for smaller values of k . By equation (4), we have

$$m(K_{n-m,m}^k, t) = m(K_{n-m,m}^{k-1}, t) + m(K_{n-m-1,m-1}^{k-1}, t - 1)$$

and

$$m(K_{n-1,1}^k, t) = m(K_{n-1,1}^{k-1}, t) + m(K_{n-2}, t - 1).$$

By the induction hypothesis and Lemma 1.3, we obtain $m(K_{n-m,m}^{k-1}, t) \leq m(K_{n-1,1}^{k-1}, t)$ and $m(K_{n-m-1,m-1}^{k-1}, t - 1) \leq m(K_{n-2}, t - 1)$. Thus the result follows. □

Combining with Lemmas 2.5 and 2.8, we obtain the following result directly.

Corollary 2.9 *Suppose $\max\{k, 2\} \leq m \leq \lfloor \frac{n}{2} \rfloor$. Then $K_{n-m,m}^k < K_{n-1,1}^k$.*

Theorem 2.10 *Let G be a graph in $\mathcal{G}_{n,k}$. Then $ME(G) \leq ME(K_{n-1,1}^k)$. The equality holds if and only if $G \cong K_{n-1,1}^k$.*

Proof Notice that $K_{n-1,1}^k \in \mathcal{G}_{n,k}$. Let $G \not\cong K_{n-1,1}^k$ be a graph in $\mathcal{G}_{n,k}$. It suffices to show that $G \prec K_{n-1,1}^k$. If G has a trivial k -edge cut, then we have $G \prec K_{n-1,1}^k$ by Lemma 2.1. Otherwise, by Lemma 2.4 and Corollary 2.9, we obtain $G \prec K_{n-1,1}^k$ again. The proof is thus complete. \square

By the proof of Theorem 2.10 and the definition of the Hosoya index, we can get the following result on the Hosoya index.

Theorem 2.11 *Let G be a graph in $\mathcal{G}_{n,k}$. Then $Z(G) \leq Z(K_{n-1,1}^k)$. The equality holds if and only if $G \cong K_{n-1,1}^k$.*

3 Conclusion

Chen *et al.* [30] characterized the extremal graphs of the matching energy in unicyclic, bicyclic graphs with a given diameter, respectively. Li *et al.* [37] obtained the unique graph having extremal matching energy in unicyclic graphs with fixed girth and the graphs with given clique number. Xu *et al.* [41] got the extremal graph of the matching energy among all t -apex trees. In [38], the authors obtained the extremal graph with matching energy in the complement of graphs. So it is interesting to characterize a graph having an extremal value of the matching energy with some graph invariants. In the paper, we got the maximal graph of the matching energy in graphs with given edge connectivity.

Therefore, our next work is to continue the study of graphs having extremal values of the matching energy with some graph invariants.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally in the design and progress of the study. All authors modified and approved the final manuscript.

Author details

¹School of Science, Shangdong University of Technology, Zibo, Shandong 255049, China. ²School of Mathematics and Statistics, Jiangsu Normal University, Xuzhou, Jiangsu 221116, China. ³School of Mathematics, Shandong University, Jinan, Shandong 250100, China.

Acknowledgements

This work was supported by National Natural Science Foundation of China (Grant Nos. 11101351, 11326216, 11401348, and 11561032), Jiangsu Government Scholarship for Overseas Studies, Natural Science Foundation of the Jiangsu Higher Education Institutions (No. 11KJB110014) and China Postdoctoral Science Foundation.

Received: 20 October 2015 Accepted: 8 December 2015 Published online: 30 December 2015

References

1. Bondy, JA, Murty, USR: Graph Theory. Springer, Berlin (2008)
2. Cvetković, D, Doob, M, Gutman, I, Torgašev, A: Recent Results in the Theory of Graph Spectra. North-Holland, Amsterdam (1988)
3. Farrell, EJ: An introduction to matching polynomials. *J. Comb. Theory, Ser. B* **27**, 75-86 (1979)
4. Gutman, I: The matching polynomial. *MATCH Commun. Math. Comput. Chem.* **6**, 75-91 (1979)
5. Gutman, I: The energy of a graph: old and new results. In: Betten, A, Kohnert, A, Laue, R, Wassermann, A (eds.) *Algebraic Combinatorics and Applications*, pp. 196-211. Springer, Berlin (2001)
6. Gutman, I, Li, X, Zhang, J: Graph energy. In: Dehmer, M, Emmert-Streib, F (eds.) *Analysis of Complex Networks: From Biology to Linguistics*, pp. 145-174. VCH, Weinheim (2009)
7. Li, X, Shi, Y, Gutman, I: *Graph Energy*. Springer, New York (2012)
8. Dehmer, M, Grabner, M: The discrimination power of molecular identification numbers revisited. *MATCH Commun. Math. Comput. Chem.* **69**, 785-794 (2013)
9. Dehmer, M, Li, X, Shi, Y: Connections between generalized graph entropies and graph energy. *Complexity* **21**(1), 35-41 (2015)
10. Das, KC, Mojallal, SA: Relation between energy and (signless) Laplacian energy of graphs. *MATCH Commun. Math. Comput. Chem.* **74**(2), 359-366 (2015)

11. Gong, S, Li, X, Xu, G, Gutman, I, Furtula, B: Borderenergetic graphs. *MATCH Commun. Math. Comput. Chem.* **74**(2), 321-332 (2015)
12. Huo, B, Ji, S, Li, X, Shi, Y: Complete solution to a problem on the maximal energy of bicyclic bipartite graphs. *Linear Algebra Appl.* **435**, 804-810 (2011)
13. Huo, B, Ji, S, Li, X, Shi, Y: Complete solution to a conjecture on the fourth maximal energy tree. *MATCH Commun. Math. Comput. Chem.* **66**(3), 903-912 (2011)
14. Ji, S, Li, J: An approach to the problem of the maximal energy of bicyclic graphs. *MATCH Commun. Math. Comput. Chem.* **68**, 741-762 (2012)
15. Li, J, Guo, J, Shiu, WC: A note on Randić energy. *MATCH Commun. Math. Comput. Chem.* **74**(2), 389-398 (2015)
16. Li, X, Qin, Z, Wei, M, Gutman, I, Dehmer, M: Novel inequalities for generalized graph entropies - graph energies and topological indices. *Appl. Comput. Math.* **259**, 470-479 (2015)
17. Li, X, Shi, Y, Wei, M, Li, J: On a conjecture about tricyclic graphs with maximal energy. *MATCH Commun. Math. Comput. Chem.* **72**(1), 183-214 (2014)
18. Rengqian, S, Ge, Y, Huo, B, Ji, S, Diao, Q: On the tree with diameter 4 and maximal energy. *Appl. Comput. Math.* **268**, 364-374 (2015)
19. Maden, AD: New bounds on the incidence energy, Randić energy and Randić estrada index. *MATCH Commun. Math. Comput. Chem.* **74**(2), 367-387 (2015)
20. Ma, H, Bai, L, Ji, S: On the minimal energy of conjugated unicyclic graphs with maximum degree at most 3. *Discrete Appl. Math.* **186**, 186-198 (2015)
21. Marin, CA, Monsalve, J, Rada, J: Maximum and minimum energy trees with two and three branched vertices. *MATCH Commun. Math. Comput. Chem.* **74**(2), 285-306 (2015)
22. Rojo, O: Effects on the energy and estrada indices by adding edges among pendent vertices. *MATCH Commun. Math. Comput. Chem.* **74**(2), 343-358 (2015)
23. Ma, J, Shi, Y, Wang, Z, Yue, J: On Wiener polarity index of bicyclic networks. *Sci. Rep.* (in press)
24. Gutman, I, Polansky, OE: *Mathematical Concepts in Organic Chemistry*. Springer, Berlin (1986)
25. Gutman, I, Wagner, S: The matching energy of a graph. *Discrete Appl. Math.* **160**(15), 2177-2187 (2012)
26. Aihara, J: A new definition of Dewar-type resonance energies. *J. Am. Chem. Soc.* **98**, 2750-2758 (1976)
27. Gutman, I, Milun, M, Trinajstić, N: Topological definition of delocalisation energy. *MATCH Commun. Math. Comput. Chem.* **1**, 171-175 (1975)
28. Gutman, I, Milun, M, Trinajstić, N: Graph theory and molecular orbitals 19, nonparametric resonance energies of arbitrary conjugated systems. *J. Am. Chem. Soc.* **99**, 1692-1704 (1977)
29. Hosoya, H: A newly proposed quantity characterizing the topological nature of structural isomers of saturated hydrocarbons. *Bull. Chem. Soc. Jpn.* **44**, 2332-2339 (1971)
30. Chen, L, Liu, J, Shi, Y: Matching energy of unicyclic and bicyclic graphs with a given diameter. *Complexity* **21**(2), 224-238 (2015)
31. Ji, S, Li, X, Shi, Y: The extremal matching energy of bicyclic graphs. *MATCH Commun. Math. Comput. Chem.* **70**, 697-706 (2013)
32. Chen, L, Shi, Y: The maximal matching energy of tricyclic graphs. *MATCH Commun. Math. Comput. Chem.* **73**, 105-119 (2015)
33. Ji, S, Ma, H: The extremal matching energy of graphs. *Ars Comb.* **115**, 343-355 (2014)
34. Li, S, Yan, W: The matching energy of graphs with given parameters. *Discrete Appl. Math.* **162**, 415-420 (2014)
35. Chen, L, Liu, J, Shi, Y: Bounds on the matching energy of unicyclic odd-cycle graphs. *MATCH Commun. Math. Comput. Chem.* **75**(2), 315-330 (2016)
36. Chen, X, Li, X, Lian, H: The matching energy of random graphs. *Discrete Appl. Math.* **193**, 102-109 (2015)
37. Li, H, Zhou, Y, Su, L: Graphs with extremal matching energies and prescribed parameters. *MATCH Commun. Math. Comput. Chem.* **72**, 239-248 (2014)
38. So, W, Wang, W: Finding the least element of the ordering of graphs with respect to their Matching Numbers. *MATCH Commun. Math. Comput. Chem.* **73**, 225-238 (2015)
39. Wang, W, So, W: On minimum matching energy of graphs. *MATCH Commun. Math. Comput. Chem.* **74**(2), 399-410 (2015)
40. Wu, T, Yan, Y, Zhang, H: Extremal matching energy of complements of trees. arXiv:1411.7458 (2014)
41. Xu, K, Zheng, Z, Das, K: Extremal t -apex trees with respect to matching energy. *Complexity* (2015). doi:10.1002/cplx.21651

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
