# RESEARCH

## **Open Access**



# Padé approximant related to inequalities for Gauss lemniscate functions

Juan Liu and Chao-Ping Chen\*

\*Correspondence: chenchaoping@sohu.com School of Mathematics and Informatics, Henan Polytechnic University, Jiaozuo City, Henan Province 454000, China

### Abstract

Based on the Padé approximation method, we present new inequalities for Gauss lemniscate functions. We also solve a conjecture on inequalities for Gauss lemniscate functions proposed by Sun and Chen.

MSC: 26D07; 41A60

Keywords: lemniscate functions; inequality; approximation

### 1 Introduction

The lemniscate, also called the lemniscate of Bernoulli, is the locus of points (x, y) in the plane satisfying the equation  $(x^2 + y^2)^2 = x^2 + y^2$ . In polar coordinates  $(r, \theta)$ , the equation becomes  $r^2 = \cos(2\theta)$  and its arc length is given by the function

$$\operatorname{arcsl} x = \int_0^x \frac{1}{\sqrt{1 - t^4}} \, \mathrm{d}t, \quad |x| \le 1,$$
 (1.1)

where arcsl is called the arc lemniscate sine function studied by Gauss in 1797-1798. Another lemniscate function investigated by Gauss is the hyperbolic arc lemniscate sine function, defined as

$$\operatorname{arcslh} x = \int_0^x \frac{1}{\sqrt{1+t^4}} \, \mathrm{d}t, \quad x \in \mathbb{R}.$$
(1.2)

The functions (1.1) and (1.2) can be found (see [1], Chapter 1, [2], p. 259, and [3–11]).

Another pair of lemniscate functions, the arc lemniscate tangent arctl and the hyperbolic arc lemniscate tangent arctlh, have been introduced in [4], (3.1)-(3.2). Therein it has been proven that

$$\operatorname{arctl} x = \operatorname{arcsl}\left(\frac{x}{\sqrt[4]{1+x^4}}\right), \quad x \in \mathbb{R}$$
 (1.3)

and

$$\operatorname{arctlh} x = \operatorname{arcslh}\left(\frac{x}{\sqrt[4]{1-x^4}}\right), \quad |x| < 1$$
 (1.4)

(see [4], Proposition 3.1).



© The Author(s) 2016. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Recently, numerous inequalities have been given for the lemniscate functions [6, 9–11]. For example, Neuman [6] proved the following inequalities:

$$\left(\frac{5}{3+2(1-x^4)^{1/2}}\right)^{1/2} < \frac{\arcsin x}{x} < \left(1-x^4\right)^{-1/10}$$
(1.5)

and

$$\left(\frac{5}{3+2(1+x^4)^{1/2}}\right)^{1/2} < \frac{\operatorname{arcslh} x}{x} < \left(1+x^4\right)^{-1/10}$$
(1.6)

for 0 < |x| < 1.

Shafer [12] indicated several elementary quadratic approximations of selected functions without proof. Subsequently, Shafer [13] established these results as analytic inequalities. For example, Shafer [13] proved that, for x > 0,

$$\frac{8x}{3 + \sqrt{25 + \frac{80}{3}x^2}} < \arctan x.$$
(1.7)

The inequality (1.7) can also be found in [14]. Zhu [15] developed (1.7) to produce a symmetric double inequality

$$\frac{8x}{3+\sqrt{25+\frac{80}{3}x^2}} < \arctan x < \frac{8x}{3+\sqrt{25+\frac{256}{\pi^2}x^2}}, \quad x > 0,$$
(1.8)

where the constants 80/3 and 256/ $\pi^2$  are the best possible. In [15], (1.8) is called a Shafertype inequality.

Mortici and Srivastava [16] presented new bounds for  $\arctan x$ . Some inequalities for trigonometric functions were refined in [17].

Very recently, Sun and Chen [18] established the following Shafer-type inequalities for the lemniscate functions:

$$\frac{10}{5 + \sqrt{25 - 10x^4}} < \frac{\arcsin x}{x}, \quad 0 < x < 1, \tag{1.9}$$

$$\frac{10}{5 + \sqrt{25 - 15x^4}} < \frac{\operatorname{arctlh} x}{x}, \quad 0 < x < 1, \tag{1.10}$$

$$\frac{95}{80 + \sqrt{225 + 285x^4}} < \frac{\operatorname{arcslh} x}{x}, \quad x > 0, \tag{1.11}$$

and presented the following conjecture.

### **Conjecture 1.1** For x > 0,

$$\frac{\operatorname{arcslh} x}{x} < \frac{95 + \frac{931}{2925}x^{12}}{80 + \sqrt{225 + 285x^4}}$$
(1.12)

and

$$\frac{1210}{940 + 9\sqrt{900 + 1210x^4}} < \frac{\arctan x}{x} < \frac{1210 + \frac{2.078.417}{280.800}x^{12}}{940 + 9\sqrt{900 + 1210x^4}}.$$
(1.13)

Based on the Padé approximation method, in this paper we present new inequalities for Gauss lemniscate functions. We also prove Conjecture 1.1.

Some computations in this paper were performed using Maple software.

### 2 Padé approximant

For later use, we introduce the Padé approximant (see [19–21]). Let f be a formal power series,

$$f(t) = c_0 + c_1 t + c_2 t^2 + \cdots .$$
(2.1)

The Padé approximation of order (p,q) of the function f is the rational function, denoted by

$$[p/q]_f(t) = \frac{\sum_{j=0}^p a_j t^j}{1 + \sum_{j=1}^q b_j t^j},$$
(2.2)

where  $p \ge 0$  and  $q \ge 1$  are any given integers, the coefficients  $a_j$  and  $b_j$  are given by (see [19, 21])

$$a_{0} = c_{0},$$

$$a_{1} = c_{0}b_{1} + c_{1},$$

$$a_{2} = c_{0}b_{2} + c_{1}b_{1} + c_{2},$$

$$\vdots$$

$$a_{p} = c_{0}b_{p} + \dots + c_{p-1}b_{1} + c_{p},$$

$$0 = c_{p+1} + c_{p}b_{1} + \dots + c_{p-q+1}b_{q},$$

$$\vdots$$

$$0 = c_{p+q} + c_{p+q-1}b_{1} + \dots + c_{p}b_{q},$$
(2.3)

and we have

1

$$[p/q]_f(t) - f(t) = O(t^{p+q+1}).$$
(2.4)

Thus, the first p + q + 1 coefficients of the series expansion of  $[p/q]_f$  are identical to those of f. Moreover, we have (see [20])

$$[p/q]_{f}(t) = \frac{\begin{vmatrix} t^{q} f_{p-q}(t) & t^{q-1} f_{p-q+1}(t) & \cdots & f_{p}(t) \\ c_{p-q+1} & c_{p-q+2} & \cdots & c_{p+1} \\ \vdots & \vdots & \ddots & \vdots \\ c_{p} & c_{p+1} & \cdots & c_{p+q} \end{vmatrix}}{\begin{vmatrix} t^{q} & t^{q-1} & \cdots & 1 \\ c_{p-q+1} & c_{p-q+2} & \cdots & c_{p+1} \\ \vdots & \vdots & \ddots & \vdots \\ c_{p} & c_{p+1} & \cdots & c_{p+q} \end{vmatrix}},$$

$$(2.5)$$

with  $f_n(x) = c_0 + c_1 x + \dots + c_n x^n$ , the *n*th partial sum of the series  $f(f_n \text{ is identically zero for } n < 0)$ .

Chen [9] presented the following power-series expansions (for |x| < 1):

$$\frac{\arccos x}{x} = \sum_{n=0}^{\infty} \frac{\Gamma(n+\frac{1}{2})}{\sqrt{\pi}(4n+1) \cdot n!} x^{4n},$$
(2.6)

$$\frac{\operatorname{arcslh} x}{x} = \sum_{n=0}^{\infty} (-1)^n \frac{\Gamma(n+\frac{1}{2})}{\sqrt{\pi}(4n+1) \cdot n!} x^{4n},$$
(2.7)

$$\frac{\arctan x}{x} = \sum_{n=0}^{\infty} (-1)^n \frac{\Gamma(n+\frac{3}{4})}{\Gamma(\frac{3}{4}) \cdot (4n+1) \cdot n!} x^{4n}$$
(2.8)

and

$$\frac{\operatorname{arctlh} x}{x} = \sum_{n=0}^{\infty} \frac{\Gamma(n+\frac{3}{4})}{\Gamma(\frac{3}{4}) \cdot (4n+1) \cdot n!} x^{4n}.$$
(2.9)

We now consider the Padé approximant for the function  $\frac{\operatorname{arcsl} x}{x}$  at the point x = 0. Let

$$f(t) = \sum_{j=0}^{\infty} c_j t^j = 1 + \frac{1}{10}t + \frac{1}{24}t^2 + \frac{5}{208}t^3 + \frac{35}{2176}t^4 + \cdots,$$
(2.10)

with the coefficients  $c_i$  given by

$$c_j = \frac{\Gamma(j + \frac{1}{2})}{\sqrt{\pi}(4j + 1) \cdot j!}.$$
(2.11)

Let us find the (2, 2) Padé approximant for the function (2.10) at the point t = 0,

$$[2/2]_f(t) = \frac{\sum_{j=0}^2 a_j t^j}{1 + \sum_{j=1}^2 b_j t^j}.$$

Noting that

$$c_0 = 1,$$
  $c_1 = \frac{1}{10},$   $c_2 = \frac{1}{24},$   $c_3 = \frac{5}{208},$   $c_4 = \frac{35}{2176},$  (2.12)

holds, we have, by (2.3),

$$\begin{cases} a_0 = 1, \\ a_1 = b_1 + \frac{1}{10}, \\ a_2 = b_2 + \frac{1}{10}b_1 + \frac{1}{24}, \\ 0 = \frac{5}{208} + \frac{1}{24}b_1 + \frac{1}{10}b_2, \\ 0 = \frac{35}{2176} + \frac{5}{508}b_1 + \frac{1}{24}b_2, \end{cases}$$

that is,

$$a_0 = 1$$
,  $a_1 = -\frac{55}{68}$ ,  $a_2 = \frac{23,623}{265,200}$ ,  $b_1 = \frac{309}{340}$ ,  $b_2 = \frac{489}{3536}$ .

We thus obtain

$$[2/2]_{f}(t) = \frac{1 - \frac{55}{68}t + \frac{23,623}{265,200}t^{2}}{1 - \frac{309}{340}t + \frac{489}{3536}t^{2}},$$
(2.13)

and we have, by (2.4),

$$f(t) = [2/2]_f(t) + O(t^5).$$
(2.14)

That is

$$\sum_{j=0}^{\infty} \frac{\Gamma(j+\frac{1}{2})}{\sqrt{\pi}(4j+1) \cdot j!} t^{j} = \frac{1 - \frac{55}{68}t + \frac{23,623}{265,200}t^{2}}{1 - \frac{309}{340}t + \frac{489}{3536}t^{2}} + O(t^{5}).$$
(2.15)

Replacing *t* by  $x^4$  in (2.15) yields

$$\frac{\arccos x}{x} = \frac{1 - \frac{55}{68}x^4 + \frac{23,623}{265,200}x^8}{1 - \frac{309}{340}x^4 + \frac{489}{3536}x^8} + O(x^{20})$$
$$= \frac{265,200 - 214,500x^4 + 23,623x^8}{15(17,680 - 16,068x^4 + 2445x^8)} + O(x^{20}).$$
(2.16)

Remark 2.1 Using (2.5), we can also derive (2.13). Indeed, we have

$$[2/2]_{f}(t) = \frac{\begin{vmatrix} t^{2}f_{0}(t) & tf_{1}(t) & f_{2}(t) \\ c_{1} & c_{2} & c_{3} \\ c_{2} & c_{3} & c_{4} \end{vmatrix}}{\begin{vmatrix} t^{2} & t & 1 \\ c_{1} & c_{2} & c_{3} \\ c_{2} & c_{3} & c_{4} \end{vmatrix}} = \frac{\begin{vmatrix} t^{2} & t(1 + \frac{1}{10}t) & 1 + \frac{1}{10}t + \frac{1}{24}t^{2} \\ \frac{1}{10} & \frac{1}{24} & \frac{5}{208} \\ \frac{1}{24} & \frac{5}{208} & \frac{35}{2176} \end{vmatrix}}{\begin{vmatrix} t^{2} & t & 1 \\ c_{1} & c_{2} & c_{3} \\ c_{2} & c_{3} & c_{4} \end{vmatrix}} = \frac{\begin{vmatrix} t^{2} & t(1 + \frac{1}{10}t) & 1 + \frac{1}{10}t + \frac{1}{24}t^{2} \\ \frac{1}{10} & \frac{1}{24} & \frac{5}{208} \\ \frac{1}{24} & \frac{5}{208} & \frac{35}{2176} \end{vmatrix}}{\begin{vmatrix} t^{2} & t & 1 \\ \frac{1}{10} & \frac{1}{24} & \frac{5}{208} \\ \frac{1}{24} & \frac{5}{208} & \frac{35}{2176} \end{vmatrix}}$$
$$= \frac{1 - \frac{55}{68}t + \frac{23,623}{265,200}t^{2}}{1 - \frac{309}{340}t + \frac{489}{3536}t^{2}}.$$

Following the same method as used in the derivation of the formula (2.16), we find

$$\frac{\operatorname{arcslh} x}{x} = \frac{1 + \frac{55}{68}x^4 + \frac{23,623}{265,200}x^8}{1 + \frac{309}{340}x^4 + \frac{489}{3536}x^8} + O(x^{20})$$
$$= \frac{265,200 + 214,500x^4 + 23,623x^8}{15(17,680 + 16,068x^4 + 2445x^8)} + O(x^{20}), \tag{2.17}$$

$$\frac{\arctan x}{x} = \frac{1 + \frac{63}{130}x^4 - \frac{139}{6240}x^8}{1 + \frac{33}{52}x^4} + O(x^{16}),$$
(2.18)

$$\frac{\operatorname{arctl} x}{x} = \frac{1 + \frac{79,047}{94,520}x^4 + \frac{565,795}{5,898,048}x^8}{1 + \frac{18,645}{18,904}x^4 + \frac{336,105}{1,966,016}x^8} + O(x^{20})$$
$$= \frac{29,490,240 + 24,662,664x^4 + 2,828,975x^8}{15(1,966,016 + 1,939,080x^4 + 336,105x^8)} + O(x^{20})$$
(2.19)

and

$$\frac{\operatorname{arctlh} x}{x} = \frac{1 - \frac{79,047}{94,520}x^4 + \frac{565,795}{5,898,048}x^8}{1 - \frac{18,645}{18,904}x^4 + \frac{336,105}{1,966,016}x^8} + O(x^{20})$$
$$= \frac{29,490,240 - 24,662,664x^4 + 2,828,975x^8}{15(1,966,016 - 1,939,080x^4 + 336,105x^8)} + O(x^{20}). \tag{2.20}$$

In view of (2.16) and (2.17), we pose the following.

### Conjecture 2.1 Let

$$\frac{\arccos x}{x} = \frac{1 + \sum_{j=1}^{n} a_j x^{4j}}{1 + \sum_{j=1}^{n} b_j x^{4j}} + O(x^{8n+4})$$
(2.21)

and

$$\frac{\operatorname{arcslh} x}{x} = \frac{1 + \sum_{j=1}^{n} \alpha_j x^{4j}}{1 + \sum_{j=1}^{n} \beta_j x^{4j}} + O(x^{8n+4}).$$
(2.22)

Then the coefficients  $a_i$  and  $\alpha_i$  satisfy the following relation:

$$a_j = (-1)^j \alpha_j, \quad j = 1, 2, \dots, n,$$
 (2.23)

and the coefficients  $b_j$  and  $\beta_j$  satisfy the following relation:

$$b_j = (-1)^j \beta_j, \quad j = 1, 2, \dots, n.$$
 (2.24)

In view of (2.19) and (2.20), we pose the following.

# Conjecture 2.2 Let

$$\frac{\arctan x}{x} = \frac{1 + \sum_{j=1}^{n} p_j x^{4j}}{1 + \sum_{j=1}^{n} q_j x^{4j}} + O(x^{8n+4})$$
(2.25)

and

$$\frac{\operatorname{arctlh} x}{x} = \frac{1 + \sum_{j=1}^{n} r_j x^{4j}}{1 + \sum_{j=1}^{n} s_j x^{4j}} + O(x^{8n+4}).$$
(2.26)

*Then the coefficients*  $p_i$  *and*  $r_i$  *satisfy the following relation:* 

$$p_j = (-1)^j r_j, \quad j = 1, 2, \dots, n,$$
 (2.27)

and the coefficients  $q_i$  and  $s_i$  satisfy the following relation:

$$q_j = (-1)^j s_j, \quad j = 1, 2, \dots, n.$$
 (2.28)

# **3** Inequalities

Equations (2.16)-(2.20) motivate us to establish the following theorems.

**Theorem 3.1** *For* 0 < *x* < 1,

$$\frac{265,200 - 214,500x^4 + 23,623x^8}{15(17,680 - 16,068x^4 + 2445x^8)} < \frac{\arcsin x}{x}.$$
(3.1)

Proof Consider the function

$$f(x) = \arcsin x - \frac{x(265,200 - 214,500x^4 + 23,623x^8)}{15(17,680 - 16,068x^4 + 2445x^8)}, \quad 0 < x < 1.$$

Differentiation yields

$$f'(x) = \frac{1}{\sqrt{1 - x^4}} - \frac{312,582,400 - 411,873,280x^4 + 177,771,984x^8 - 21,634,288x^{12} + 3,850,549x^{16}}{(17,680 - 16,068x^4 + 2445x^8)^2}.$$

Elementary calculations reveal that

$$\begin{split} & \left(\frac{1}{\sqrt{1-t}}\right)^2 \\ & -\left(\frac{312,582,400-411,873,280t+177,771,984t^2-21,634,288t^3+3,850,549t^4}{(17,680-16,068t+2445t^2)^2}\right)^2 \\ & = \frac{t^5g(t)}{(1-t)(17,680-16,068t+2445t^2)^4}, \quad 0 < t < 1, \end{split}$$

where

$$g(t) = 1,744,280,123,040,000 - 2,406,774,938,256,000t$$
$$+ 1,064,272,682,007,600t^{2}$$
$$- 145,697,716,749,000t^{3} + 14,826,727,601,401t^{4}.$$

We now prove that f'(x) > 0 for 0 < x < 1. It suffices to show that g(t) > 0 for 0 < t < 1. Differentiation yields

$$g'(t) = -2,406,774,938,256,000 + 2,128,545,364,015,200t$$
$$-437,093,150,247,000t^{2}$$
$$+59,306,910,405,604t^{3}$$

and

$$g''(t) = 2,128,545,364,015,200 - 874,186,300,494,000t$$
$$+ 177,920,731,216,812t^2 > 0, \quad 0 < t < 1.$$

We then obtain, for 0 < t < 1,

$$g'(t) < g'(1) = -656,015,814,082,196 < 0 \implies$$
  
 $g(t) > g(1) = 270,906,877,644,001 > 0.$ 

Hence, f'(x) > 0 for 0 < x < 1, and we have

$$f(x) > f(0) = 0, \quad 0 < x < 1.$$

The proof is complete.

**Remark 3.1** *There is no strict comparison between the two lower bounds in* (1.5) *and* (3.1).

**Theorem 3.2** *For* x > 0,

$$\frac{\operatorname{arcslh} x}{x} < \frac{265,200 + 214,500x^4 + 23,623x^8}{15(17,680 + 16,068x^4 + 2445x^8)}.$$
(3.2)

*Proof* Consider the function

$$F(x) = \operatorname{arcslh} x - \frac{x(265,200+214,500x^4+23,623x^8)}{15(17,680+16,068x^4+2445x^8)}, \quad x > 0.$$

Differentiation yields

$$F'(x) = \frac{1}{\sqrt{1 + x^4}} - \frac{312,582,400 + 411,873,280x^4 + 177,771,984x^8 + 21,634,288x^{12} + 3,850,549x^{16}}{(17,680 + 16,068x^4 + 2445x^8)^2}.$$

Elementary calculations reveal that

$$\begin{split} &\left(\frac{1}{\sqrt{1+t}}\right)^2 \\ &\quad -\left(\frac{312,582,400+411,873,280t+177,771,984t^2+21,634,288t^3+3,850,549t^4}{(17,680+16,068t+2445t^2)^2}\right)^2 \\ &\quad = -\frac{t^5 G(t)}{(1+t)(17,680+16,068t+2445t^2)^4}, \end{split}$$

where

$$G(t) = 1,744,280,123,040,000 + 2,406,774,938,256,000t + 1,064,272,682,007,600t^{2}$$
$$+ 145,697,716,749,000t^{3} + 14,826,727,601,401t^{4}.$$

Hence, F'(x) < 0 for x > 0, and we have

$$F(x) < F(0) = 0, \quad x > 0.$$

The proof is complete.

### **Remark 3.2** *For* 0 < t < 1*, we find*

$$\begin{split} I(t) &:= \frac{1}{1+t} - \left(\frac{265,200+214,500t+23,623t^2}{15(17,680+16,068t+2445t^2)}\right)^{10} \\ &= \frac{t^2 P_{19}(t)}{576,650,390,625(1+t)(17,680+16,068t+2445t^2)^{10}} \end{split}$$

with

$$P_{19}(t) = P_{16}(t) + t^{17} P_2(t),$$

where

$$P_{16}(t) = 3,309,224,024,069,080,418,989,754,522,912,085,339,870,997,824,000t^{16}$$
  
+ ...  
+ 229,442,535,851,108,636,620,015,850,036,920,320,000,000,000,000,000

is a polynomial of the 16th degree, having all coefficients positive, and

$$P_2(t) = 77,541,624,086,159,498,428,020,328,992,837,339,887,447,064,000$$
  
- 565,686,157,207,722,134,655,870,693,904,642,976,763,301,024t  
- 54,119,091,759,561,776,058,592,767,712,571,305,215,681,649 $t^2 > 0$ 

for 0 < t < 1. So, I(t) > 0 for 0 < t < 1. We then see that the inequality (3.2) is sharper than the right side of (1.6).

**Theorem 3.3** *For x* > 0,

$$\frac{1+\frac{63}{130}x^4-\frac{139}{6240}x^8}{1+\frac{33}{52}x^4} < \frac{\arctan x}{x} < \frac{29,490,240+24,662,664x^4+2,828,975x^8}{15(1,966,016+1,939,080x^4+336,105x^8)}.$$
 (3.3)

*Proof* Consider the function

$$\lambda(x) = \arctan x - \frac{x(1 + \frac{63}{130}x^4 - \frac{139}{6240}x^8)}{1 + \frac{33}{52}x^4}.$$

Differentiation yields

$$\lambda'(x) = \frac{1}{(1+x^4)^{3/4}} + \frac{x^3(48,672+14,456x^4+4587x^8)}{30(52+33x^4)^2} > 0.$$

We then obtain

$$\lambda(x) > \lambda(0) = 0, \quad x > 0.$$

Hence the first inequality in (3.3) holds for x > 0.

Consider the function

$$T(x) = \arctan x - \frac{x(29,490,240+24,662,664x^4+2,828,975x^8)}{15(1,966,016+1,939,080x^4+336,105x^8)}, \quad x > 0.$$

Differentiation yields

$$T'(x) = \frac{1}{(1+x^4)^{3/4}} - \frac{P_{16}(x)}{(1,966,016+1,939,080x^4+336,105x^8)^2}$$

where

$$P_{16}(x) = 3,865,218,912,256 + 4,725,610,426,368x^4 + 1,899,763,315,008x^8 + 170,687,344,256x^{12} + 63,388,842,825x^{16}.$$

Elementary calculations reveal that

$$\begin{aligned} &\frac{1}{(1+x^4)^3} - \left(\frac{P_{16}(x)}{(1,966,016+1,939,080x^4+336,105x^8)^2}\right)^4 \\ &= -\frac{x^{20}P_{56}(x)}{(1+x^4)^3(1,966,016+1,939,080x^4+336,105x^8)^8}, \end{aligned}$$

where

 $P_{56}(x) = 13,193,567,461,486,862,074,082,196,527,146,063,235,598,765,785,088$   $+ 75,159,817,580,420,162,914,879,309,165,363,929,497,102,849,146,880x^4$   $+ 190,493,075,741,254,897,950,338,074,805,626,536,902,462,936,186,880x^8$   $+ 283,233,781,637,227,052,608,425,496,608,925,420,321,925,549,260,800x^{12}$   $+ 274,381,791,750,085,496,947,276,941,731,670,794,946,132,888,780,800x^{16}$   $+ 182,271,787,590,701,615,339,301,540,193,194,594,557,235,390,578,688x^{20}$   $+ 85,570,287,614,566,775,085,144,063,641,805,696,286,360,924,323,840x^{24}$   $+ 29,163,131,006,055,200,534,183,374,987,447,946,919,333,657,968,640x^{28}$   $+ 7,481,144,367,677,341,229,619,045,201,182,337,247,982,930,534,400x^{32}$   $+ 1,502,545,339,351,309,468,552,186,115,563,901,082,330,882,201,600x^{36}$   $+ 238,495,639,577,137,257,561,813,891,822,862,592,696,929,928,896x^{40}$   $+ 29,999,531,147,567,967,753,948,099,263,441,234,315,939,560,800x^{44}$   $+ 3,208,050,013,558,968,652,633,219,730,412,159,840,683,611,875x^{48}$   $+ 222,336,558,000,169,152,230,844,556,985,454,831,178,171,875x^{52}$ 

 $+\ 16,\!145,\!492,\!412,\!888,\!980,\!411,\!169,\!048,\!998,\!579,\!532,\!875,\!390,\!625x^{56}.$ 

Hence, T'(x) < 0 for x > 0, and we have

 $T(x) < T(0) = 0, \quad x > 0.$ 

Hence, the second inequality in (3.3) holds for x > 0. The proof is complete.

**Theorem 3.4** *For* 0 < *x* < 1,

$$\frac{29,490,240 - 24,662,664x^4 + 2,828,975x^8}{15(1,966,016 - 1,939,080x^4 + 336,105x^8)} < \frac{\operatorname{arctlh} x}{x}.$$
(3.4)

*Proof* Consider the function

$$H(x) = \operatorname{arctlh} x - \frac{x(29,490,240 - 24,662,664x^4 + 2,828,975x^8)}{15(1,966,016 - 1,939,080x^4 + 336,105x^8)}, \quad 0 < x < 1.$$

Differentiation yields

$$H'(x) = \frac{1}{(1-x^4)^{3/4}} - \frac{Q(x^4)}{(1,966,016-1,939,080x^4+336,105x^8)^2},$$

where

$$Q(t) = 3,865,218,912,256 - 4,725,610,426,368t + 1,899,763,315,008t^{2}$$
$$- 170,687,344,256t^{3} + 63,388,842,825t^{4}, \quad 0 < t < 1.$$

Elementary calculations reveal that

$$\frac{1}{(1-t)^3} - \left(\frac{Q(t)}{(1,966,016-1,939,080t+336,105t^2)^2}\right)^4$$
$$= \frac{t^5 R(t)}{(1-t)^3 (1,966,016-1,939,080t+336,105t^2)^8},$$

where

$$\begin{split} R(t) &= 301,748,693,573,399,407,094,173,717,482,883,533,487,653,121 \\ &+ 9,932,615,535,811,554,413,076,061,553,884,665,646,471,005,365(1-t) \\ &+ 151,766,449,766,787,034,704,161,483,173,419,604,277,111,004,855(1-t)^2 \\ &+ 680,433,563,535,649,162,659,902,808,405,182,093,255,793,778,810(1-t)^3 \\ &+ 1856,390,570,444,186,005,047,799,006,039,729,435,155,242,856,245(1-t)^4 \\ &+ 3,124,371,679,128,783,209,196,299,976,215,123,075,149,799,026,971(1-t)^5 \\ &+ 3,542,098,875,374,455,780,446,672,503,755,643,630,859,189,032,595(1-t)^6 \\ &+ 2,471,319,403,553,122,128,066,406,333,619,269,039,662,418,255,020(1-t)^7 \\ &+ 1,090,939,975,419,395,315,836,646,390,524,713,606,873,688,710,195(1-t)^8 \\ &+ 188,439,516,872,719,220,883,777,738,283,311,857,263,725,003,515(1-t)^9 \\ &+ 72,805,480,166,035,035,195,735,976,881,949,595,398,022,003,221(1-t)^{10} \\ &+ 2,968,223,270,581,948,926,689,804,107,877,843,092,991,437,050(1-t)^{11} \\ &+ (1,786,914,569,129,666,891,048,623,948,471,984,527,027,924,375 \\ &- 3,700,335,780,276,573,525,522,128,994,658,629,077,296,875(1-t))(1-t)^{12} \\ &+ 16,145,492,412,888,980,411,169,048,998,579,532,875,390,625(1-t)^{14}. \end{split}$$

Since R(t) > 0 for 0 < t < 1, we have H'(x) > 0 for 0 < x < 1. We then obtain

$$H(x) > H(0) = 0, \quad 0 < x < 1.$$

The proof is complete.

### 4 Proof of Conjecture 1.1

*Proof of* (1.12) It suffices to show by (3.2) that

$$\frac{265,200+214,500x^4+23,623x^8}{15(17,680+16,068x^4+2445x^8)} < \frac{95+\frac{931}{2925}x^{12}}{80+\sqrt{225+285x^4}}, \quad x > 0,$$

i.e.,

$$\frac{95 + \frac{931}{2925}x^{12}}{\frac{265,200 + 214,500x^4 + 23,623x^8}{15(17,680 + 16,068x^4 + 2445x^8)}} - 80 > \sqrt{225 + 285x^4}, \quad x > 0.$$
(4.1)

Elementary calculations show that

$$\begin{pmatrix} \frac{95 + \frac{931}{2925}x^{12}}{\frac{265,200+214,500x^4+23,623x^8}{15(17,680+16,068x^4+2445x^8)}} - 80 \end{pmatrix}^2 - (\sqrt{225 + 285x^4})^2 \\ = \frac{x^{16}P_{24}(x)}{(265,200 + 214,500x^4 + 23,623x^8)^2},$$

where

$$\begin{split} P_{24}(x) &= 2,214,994,396,680,000 + 2,167,794,625,751,625x^4 \\ &\quad + 771,850,648,332,400x^8 \\ &\quad + 100,410,388,038,870x^{12} + 15,721,941,655,056x^{16} + 3,584,399,789,880x^{20} \\ &\quad + 272,711,522,475x^{24}. \end{split}$$

We see from  $P_{24}(x) > 0$  that (4.1) holds. The proof is complete.

*Proof of* (1.13) First of all, we prove the second inequality in (1.13). It suffices to show by the right-hand side of (3.3) that

$$\frac{29,490,240+24,662,664x^4+2,828,975x^8}{15(1,966,016+1,939,080x^4+336,105x^8)} < \frac{1210+\frac{2,078,417}{280,800}x^{12}}{940+9\sqrt{900+1210x^4}}, \quad x > 0,$$

i.e.,

$$\frac{1210 + \frac{2,078,417}{280,800}x^{12}}{\frac{29,490,240 + 24,662,664x^4 + 2,828,975x^8}{15(1,966,016 + 1,939,080x^4 + 336,105x^8)} - 940 > 9\sqrt{900 + 1210x^4}, \quad x > 0.$$
(4.2)

Elementary calculations show that

$$\left(\frac{1210 + \frac{2,078,417}{280,800}x^{12}}{\frac{29,490,240 + 24,662,664x^4 + 2,828,975x^8}{15(1,966,016 + 1,939,080x^4 + 336,105x^8)} - 940\right)^2 - \left(9\sqrt{900 + 1210x^4}\right)^2$$
$$= \frac{P_{40}(x)}{350,438,400(29,490,240 + 24,662,664x^4 + 2,828,975x^8)^2},$$

where

$$P_{40}(x) = 423,992,204,507,234,653,175,808,000,000$$

- + 813,161,362,018,201,812,231,782,400,000x<sup>4</sup>
- $+516,869,957,972,853,387,975,720,960,000x^8$
- $+ 125,747,707,439,033,797,403,639,808,000x^{12}$
- $+\,18,\!812,\!522,\!309,\!950,\!882,\!139,\!627,\!520,\!000x^{16}$
- $+\ 6,902,\!175,\!873,\!182,\!801,\!970,\!021,\!120,\!000x^{20}$
- $+1,857,663,393,190,652,946,224,195,584x^{24}$
- + 192,485,925,752,231,924,989,587,840 $x^{28}$
- $+21,951,612,856,626,590,316,104,640x^{32}$
- $+5,630,747,696,194,377,041,485,200x^{36}$
- $+ 487,994,939,463,408,187,266,225x^{40}.$

We see from  $P_{40}(x) > 0$  that (4.2) holds. Hence, the second inequality in (1.13) holds.

Second, we prove the first inequality in (1.13). We consider two cases.

*Case* 1. 0 < *x* < 1.

It suffices to show by the left-hand side of (3.3) that

$$\frac{1210}{940 + 9\sqrt{900 + 1210x^4}} < \frac{1 + \frac{63}{130}x^4 - \frac{139}{6240}x^8}{1 + \frac{33}{57}x^4}, \quad 0 < x < 1,$$

i.e.,

$$9\sqrt{900 + 1210x^4} > \frac{1210}{\frac{1 + \frac{63}{130}x^4 - \frac{139}{6240}x^8}{1 + \frac{33}{52}x^4}} - 940, \quad 0 < x < 1.$$
(4.3)

Elementary calculations show that

$$(9\sqrt{900 + 1210x^4})^2 - \left(\frac{1210}{\frac{1 + \frac{63}{130}x^4 - \frac{139}{6240}x^8}{1 + \frac{33}{52}x^4}} - 940\right)^2$$
  
=  $\frac{1210x^{12}(128,621,376 - 81,039,502x^4 + 1,565,001x^8)}{(6240 + 3024x^4 - 139x^8)^2} > 0, \quad 0 < x < 1,$ 

which shows that (4.3) holds.

Case 2.  $x \ge 1$ . Consider the function U(x) defined by

$$U(x) = \arctan x - \frac{1210x}{940 + 9\sqrt{900 + 1210x^4}}$$

Differentiation yields

$$U'(x) = \frac{1}{(1+x^4)^{3/4}} + \frac{12,100(1089x^4 - 94\sqrt{900 + 1210x^4} - 810)}{(940 + 9\sqrt{900 + 1210x^4})^2\sqrt{900 + 1210x^4}}.$$
(4.4)

Noting that

$$1089x^4 - 94\sqrt{900 + 1210x^4} - 810 > 0, \quad x \ge 2,$$

holds, we obtain

$$U'(x) > 0, \quad x \ge 2.$$

We now show that U'(x) > 0 is also valid for  $1 \le x < 2$ . It suffices to show that

$$y(x) > 0, \quad 1 \le x < 2,$$

where

$$y(x) = y_1(x) + y_2(x),$$

with

$$y_1(x) = \frac{(940 + 9\sqrt{900 + 1210x^4})^2\sqrt{900 + 1210x^4}}{12,100(1 + x^4)^{3/4}} + 1089x^4 - 810$$

and

$$y_2(x) = -94\sqrt{900 + 1210x^4}.$$

Differentiation yields

$$y_1'(x) = \frac{x^3(940 + 9\sqrt{900 + 1210x^4})y_3(x)}{1210(1 + x^4)^{7/4}\sqrt{900 + 1210x^4}} + 32,596x^3,$$

where

$$y_{3}(x) = 4104\sqrt{900 + 1210x^{4}} + 3267x^{4}\sqrt{900 + 1210x^{4}} - 113,740x^{4} - 26,320$$
  
>  $4104\sqrt{1210x^{4}} + 3267x^{4}\sqrt{1210x^{4}} - 113,740x^{4} - 26,320$   
=  $81,081\sqrt{10} - 140,060 + (305,910\sqrt{10} - 454,960)(x - 1)$   
+  $(584,199\sqrt{10} - 682,440)(x - 1)^{2} + (718,740\sqrt{10} - 454,960)(x - 1)^{3}$ 

+ 
$$(539,055\sqrt{10} - 113,740)(x-1)^4$$
 +  $215,622\sqrt{10}(x-1)^5$  +  $35,937\sqrt{10}(x-1)^6$   
> 0 for  $1 \le x \le 2$ .

Hence, we have  $y'_1(x) > 0$  for  $1 \le x < 2$ .

Let  $1 \le r \le x \le s \le 2$ . Since  $y_1(x)$  is increasing and  $y_2(x)$  is decreasing for  $1 \le x \le 2$ , we obtain

$$y(x) \ge y_1(r) + y_2(s) =: \sigma_1(r, s).$$

We divide the interval [1,2] into 100 subintervals:

$$[1,2] = \bigcup_{k=0}^{99} \left[ 1 + \frac{k}{100}, 1 + \frac{k+1}{100} \right] > 0 \quad \text{for } k = 0, 1, 2, \dots, 99.$$

By direct computation we get

$$\sigma_1\left(1+\frac{k}{100},1+\frac{k+1}{100}\right) > 0 \quad \text{for } k = 0,1,2,\dots,99.$$

Hence,

$$y(x) > 0$$
 for  $x \in \left[1 + \frac{k}{100}, 1 + \frac{k+1}{100}\right]$  and  $k = 0, 1, 2, \dots, 99$ .

This proves U'(x) > 0 for  $1 \le x < 2$ .

We then obtain U'(x) > 0 for all  $x \ge 1$ , and we have

$$U(x) > U(1) = 0.00154438... > 0$$
 for  $x \ge 1$ ,

which shows the first inequality in (1.13) holds for  $x \ge 1$ . Thus, the first inequality in (1.13) holds for all x > 0. The proof is complete.

### **Competing interests**

The authors declare that they have no competing interests.

# Authors' contributions

All authors read and approved the final manuscript.

### Acknowledgements

The authors thank the referees for helpful comments.

### Received: 22 September 2016 Accepted: 29 November 2016 Published online: 08 December 2016

### References

- 1. Siegel, CL: Topics in Complex Function Theory, vol. 1. Wiley, New York (1969)
- 2. Borwein, JM, Borwein, PB: Pi and the AGM: A Study in the Analytic Number Theory and Computational Complexity. Wiley, New York (1987)
- 3. Carlson, BC: Algorithms involving arithmetic and geometric means. Am. Math. Mon. 78, 496-505 (1971)
- 4. Neuman, E: On Gauss lemniscate functions and lemniscatic mean. Math. Pannon. 18, 77-94 (2007)
- 5. Neuman, E: Two-sided inequalities for the lemniscate functions. J. Inequal. Spec. Funct. 1, 1-7 (2010)
- 6. Neuman, E: On Gauss lemniscate functions and lemniscatic mean II. Math. Pannon. 23, 65-73 (2012)
- 7. Neuman, E: Inequalities for Jacobian elliptic functions and Gauss lemniscate functions. Appl. Math. Comput. 218, 7774-7782 (2012)
- 8. Neuman, E: On lemniscate functions. Integral Transforms Spec. Funct. 24, 164-171 (2013)
- 9. Chen, CP: Wilker and Huygens type inequalities for the lemniscate functions. J. Math. Inequal. 6, 673-684 (2012)

- Chen, CP: Wilker and Huygens type inequalities for the lemniscate functions, II. Math. Inequal. Appl. 16, 577-586 (2013)
- 11. Deng, JE, Chen, CP: Sharp Shafer-Fink type inequalities for Gauss lemniscate functions. J. Inequal. Appl., **2014** 35 (2014) http://www.journalofinequalitiesandapplications.com/content/2014/1/35
- 12. Shafer, RE: On quadratic approximation. SIAM J. Numer. Anal. 11, 447-460 (1974)
- Shafer, RE: Analytic inequalities obtained by quadratic approximation. Publ. Elektroteh. Fak. Univ. Beogr., Ser. Mat. Fiz. 577-598, 96-97 (1977)
- 14. Shafer, RE: On quadratic approximation, II. Publ. Elektroteh. Fak. Univ. Beogr., Ser. Mat. Fiz. 602-633, 163-170 (1978)
- 15. Zhu, L: On a quadratic estimate of Shafer. J. Math. Inequal. 2, 571-574 (2008)
- 16. Mortici, C, Srivastava, HM: Estimates for the arctangent function related to Shafer's inequality. Colloq. Math. 136, 263-270 (2014)
- 17. Debnath, L, Mortici, C, Zhu, L: Refinements of Jordan-Stečkin and Becker-Stark inequalities. Results Math. 67, 207-215 (2015)
- Sun, JL, Chen, CP: Shafer-type inequalities for inverse trigonometric functions and Gauss lemniscate functions. J. Inequal. Appl., 2016 212 (2016) http://link.springer.com/article/10.1186/s13660-016-1157-2/fulltext.html
- Bercu, G: Padé approximant related to remarkable inequalities involving trigonometric functions. J. Inequal. Appl., 2016 99 (2016) http://www.doc88.com/p-0037658479714.html
- 20. Brezinski, C, Redivo-Zaglia, M: New representations of Padé, Padé-type, and partial Padé approximants. J. Comput. Appl. Math. 284, 69-77 (2015)
- 21. Bercu, G, Wu, S: Refinements of certain hyperbolic inequalities via the Padé approximation method. J. Nonlinear Sci. Appl. 9, 5011-5020 (2016)

### Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- ► Convenient online submission
- ► Rigorous peer review
- Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ► High visibility within the field
- ► Retaining the copyright to your article

Submit your next manuscript at > springeropen.com