

RESEARCH

Open Access



CrossMark

Some inequalities on the spectral radius of matrices

Linlin Zhao^{1*} and Qingbing Liu²

*Correspondence:

zhaolinlin0635@163.com

¹College of Mathematics Science,
Dezhou University, Dezhou,
Shandong 253023, People's
Republic of China
Full list of author information is
available at the end of the article

Abstract

Let A_1, A_2, \dots, A_k be nonnegative matrices. In this paper, some upper bounds for the spectral radius $\rho(A_1 \circ A_2 \circ \dots \circ A_k)$ are proposed. These bounds generalize some existing results, and comparisons between these bounds are also considered.

MSC: 15A06; 15A42; 15B34

Keywords: spectral radius; nonnegative matrix; Hadamard product

1 Introduction

Let M_n denote the set of all $n \times n$ complex matrices and $A = (a_{ij}), B = (b_{ij}) \in M_n$. If $a_{ij} - b_{ij} \geq 0$, we say that $A \geq B$, and if $a_{ij} \geq 0$, we say that A is nonnegative, denoted by $A \geq 0$. The symbol $\rho(A)$ stands for the spectral radius of A . If A is a nonnegative matrix, the Perron-Frobenius theorem guarantees that $\rho(A) \in \sigma(A)$, where $\sigma(A)$ denotes the spectrum of A .

If there does not exist a permutation matrix P such that

$$P^T AP = \begin{pmatrix} A_1 & A_{12} \\ 0 & A_2 \end{pmatrix},$$

where A_1, A_2 are square matrices, then A is called irreducible.

Let A be an irreducible nonnegative matrix. It is well known that there exists a positive vector u such that $Au = \rho(A)u$, u being called a right Perron eigenvector of A .

The Hadamard product of A, B is defined as $A \circ B = (a_{ij}b_{ij}) \in M_n$.

Let $A \geq 0, B \geq 0$. By using the Gersgorin theorem, Brauer theorem and Brualdi theorem, respectively, the authors of [1–5] have given some inequalities for the upper bounds of $\rho(A \circ B)$. Audenaert [6], Horn and Zhang [7] proved a beautiful inequality on $\rho(A \circ B)$ for nonnegative matrices A and B , that is, $\rho(A \circ B) \leq \rho(AB)$. Huang [8] generalized the above inequality to any k nonnegative matrices, that is, $\rho(A_1 \circ A_2 \circ \dots \circ A_k) \leq \rho(A_1 A_2 \cdots A_k)$. Motivated by [8] and [1–4, 9, 10], in this paper we propose some inequalities on the upper bounds for the spectral radius of the Hadamard product of any k nonnegative matrices. These bounds generalize some existing results, and some comparisons between these bounds are also considered.

2 Main results

First, we give some lemmas which are useful for obtaining the main results.

Lemma 2.1 ([11]) Let $A \in M_n$ be a nonnegative matrix. If A_k is a principal submatrix of A , then $\rho(A_k) \leq \rho(A)$. If A is irreducible and $A_k \neq A$, then $\rho(A_k) < \rho(A)$.

Lemma 2.2 ([11]) If $A \in M_n$ is an irreducible nonnegative matrix, and $Az \leq kz$ for a nonzero nonnegative vector z , then $\rho(A) \leq k$.

Lemma 2.3 ([12]) Let $A = (a_{ij}) \in M_n$ be a nonnegative matrix. Then

$$\rho(A) \leq \max_{i \neq j} \frac{1}{2} \left\{ a_{ii} + a_{jj} + \left[(a_{ii} - a_{jj})^2 + 4 \sum_{k \neq i} a_{ik} \sum_{k \neq j} a_{jk} \right]^{\frac{1}{2}} \right\}.$$

Lemma 2.4 Let $A_1, A_2, \dots, A_k \in M_n$ and D_1, D_2, \dots, D_k be diagonal matrices of order n , then

$$\begin{aligned} & D^{-1}(A_1 \circ A_2 \circ \cdots \circ A_k)D \\ &= (D_1^{-1}A_1D_1) \circ (D_2^{-1}A_2D_2) \circ \cdots \circ (D_k^{-1}A_kD_k), \end{aligned}$$

where D equals the product of the matrices D_k, D_{k-1}, \dots, D_1 , that is, $D = D_k \cdots D_2 D_1$.

Proof Let $d_{r,i}$ be the i th diagonal of D_r and $a_{r,ij}$ be the (i,j) entry of A_r ($r = 1, 2, \dots, k$). Then the (i,j) entry of $D^{-1}(A_1 \circ A_2 \circ \cdots \circ A_k)D$ is

$$\frac{1}{\prod_{r=1}^k d_{r,i}} \left(\prod_{r=1}^k a_{r,ij} \right) \prod_{r=1}^k d_{r,j} = \prod_{r=1}^k \left(\frac{1}{d_{r,i}} a_{r,ij} d_{r,j} \right),$$

which coincides with the (i,j) entry of $(D_1^{-1}A_1D_1) \circ (D_2^{-1}A_2D_2) \circ \cdots \circ (D_k^{-1}A_kD_k)$. The proof is completed. \square

Theorem 2.1 Let $A_1, A_2, \dots, A_k \in M_n$ and $A_1 = (a_{ij}) \geq 0, A_2 = (b_{ij}) \geq 0, \dots, A_k = (k_{ij}) \geq 0$. Then

$$\begin{aligned} & \rho(A_1 \circ A_2 \circ \cdots \circ A_k) \\ & \leq \max_{1 \leq i \leq n} \{ a_{ii}b_{ii} \cdots k_{ii} + (\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii}) \}. \end{aligned} \quad (2.1)$$

Proof If $A_1 \circ A_2 \circ \cdots \circ A_k$ is irreducible, then A_1, A_2, \dots, A_k are all irreducible. From Lemma 2.1, we have

$$\rho(A_1) - a_{ii} > 0, \quad \rho(A_2) - b_{ii} > 0, \quad \dots, \quad \rho(A_k) - k_{ii} > 0, \quad \forall i \in N.$$

Since A_1, A_2, \dots, A_k are nonnegative irreducible, there exist k positive vectors u, v, \dots, w such that $A_1u = \rho(A_1)u, A_2^T v = \rho(A_2)v, \dots, A_k^T w = \rho(A_k)w$. Thus, we have

$$a_{ii}u_i + \sum_{j \neq i} a_{ij}u_j = \rho(A_1)u_i, \quad \forall i \in N,$$

$$b_{jj}v_j + \sum_{i \neq j} b_{ij}v_i = \rho(A_2)v_j, \quad \forall j \in N,$$

...,

$$k_{jj}w_j + \sum_{i \neq j} k_{ij}w_i = \rho(A_k)w_j, \quad \forall j \in N.$$

Thus, we have

$$b_{ij} \leq \frac{[\rho(A_2) - b_{jj}]v_j}{v_i}, \quad \dots, \quad k_{ij} \leq \frac{[\rho(A_k) - k_{jj}]w_j}{w_i}.$$

Let z be the vector (z_i) , where

$$z_i = \frac{u_i}{(\rho(A_2) - b_{ii})v_i \cdots (\rho(A_k) - k_{ii})w_i} > 0, \quad \forall i \in N.$$

We define $P = A_1 \circ A_2 \circ \cdots \circ A_k$. For any $i \in N$,

$$\begin{aligned} (Pz)_i &= a_{ii}b_{ii} \cdots k_{ii}z_i + \sum_{i \neq j} a_{ij}b_{ij} \cdots k_{ij}z_j \\ &\leq a_{ii}b_{ii} \cdots k_{ii}z_i + \sum_{i \neq j} a_{ij} \frac{(\rho(A_2) - b_{jj})v_j}{v_i} \cdots \frac{(\rho(A_k) - k_{jj})w_j}{w_i} z_j. \end{aligned}$$

For

$$z_j = \frac{u_j}{(\rho(A_2) - b_{jj})v_j \cdots (\rho(A_k) - k_{jj})w_j},$$

we have

$$\begin{aligned} (Pz)_i &\leq a_{ii}b_{ii} \cdots k_{ii}z_i + \frac{1}{v_i \cdots w_i} \sum_{i \neq j} a_{ij}u_j \\ &= a_{ii}b_{ii} \cdots k_{ii}z_i + \frac{1}{v_i \cdots w_i} (\rho(A_1) - a_{ii})u_i \\ &= a_{ii}b_{ii} \cdots k_{ii}z_i + (\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii})(\rho(A_1) - a_{ii})z_i \\ &= \{a_{ii}b_{ii} \cdots k_{ii} + (\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii})\}z_i. \end{aligned}$$

By Lemma 2.2, this shows that

$$\rho(A_1 \circ \cdots \circ A_k) \leq \max_{1 \leq i \leq n} \{a_{ii}b_{ii} \cdots k_{ii} + (\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii})\}.$$

If $A_1 \circ A_2 \circ \cdots \circ A_k$ is reducible, we denote by $P = (p_{ij})$ the $n \times n$ permutation matrix with $p_{12} = p_{23} = \cdots = p_{n1} = 1$, the remaining $p_{ij} = 0$, then all $A_1 + tP, A_2 + tP, \dots, A_k + tP$ are nonnegative irreducible matrices for any chosen positive real numbers t . We substitute $A_1 + tP, A_2 + tP, \dots, A_k + tP$ for A_1, A_2, \dots, A_k , respectively, in the previous case, and then, letting $t \rightarrow 0$, the result follows by continuity. The proof is completed. \square

Setting $k = 2$ in Theorem 2.1, we have the following corollary.

Corollary 2.1 ([1]) Let $A_1, A_2 \in M_n$ and $A_1 \geq 0, A_2 \geq 0$. Then

$$\rho(A_1 \circ A_2) \leq \max_{1 \leq i \leq n} \{a_{ii}b_{ii} + (\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii})\}.$$

Theorem 2.2 Let $A_1, A_2, \dots, A_k \in M_n$ and $A_1 = (a_{ij}) \geq 0, A_2 = (b_{ij}) \geq 0, \dots, A_k = (k_{ij}) \geq 0$. Then

$$\begin{aligned} & \rho(A_1 \circ A_2 \circ \cdots \circ A_k) \\ & \leq \max_{i \neq j} \frac{1}{2} \{a_{ii}b_{ii} \cdots k_{ii} + a_{jj}b_{jj} \cdots k_{jj} + [(a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj})^2 \\ & \quad + 4(\rho(A_1) - a_{ii}) \cdots (\rho(A_k) - k_{ii})(\rho(A_1) - a_{jj}) \cdots (\rho(A_k) - k_{jj})]^{\frac{1}{2}}\}. \end{aligned} \quad (2.2)$$

Proof First we assume that $A_1 \circ A_2 \circ \cdots \circ A_k$ is irreducible. Obviously, A_1, A_2, \dots, A_k are all irreducible, from Lemma 2.1, we have

$$\rho(A_1) - a_{ii} > 0, \quad \rho(A_2) - b_{ii} > 0, \quad \dots, \quad \rho(A_k) - k_{ii} > 0, \quad \forall i \in N.$$

For the irreducibility of A_1, A_2, \dots, A_k , there exist k positive vectors $u = (u_i), v = (v_i), \dots, w = (w_i)$ such that $A_1u = \rho(A_1)u, A_2v = \rho(A_2)v, \dots, A_kw = \rho(A_k)w$. Thus, we have

$$a_{ii} + \sum_{j \neq i} \frac{a_{ij}u_j}{u_i} = \rho(A_1), \quad \forall i \in N,$$

$$b_{ii} + \sum_{j \neq i} \frac{b_{ij}v_j}{v_i} = \rho(A_2), \quad \forall i \in N,$$

\dots

$$k_{ii} + \sum_{j \neq i} \frac{k_{ij}w_j}{w_i} = \rho(A_k), \quad \forall i \in N.$$

Define

$$U = \text{diag}(u_1, u_2, \dots, u_n), \quad V = \text{diag}(v_1, v_2, \dots, v_n), \quad \dots,$$

$$W = \text{diag}(w_1, w_2, \dots, w_n).$$

Let

$$\hat{A}_1 = (\hat{a}_{ij}) = U^{-1}A_1U = \left(\frac{1}{u_i}a_{ij}u_j \right),$$

$$\hat{A}_2 = (\hat{b}_{ij}) = V^{-1}A_2V = \left(\frac{1}{v_i}b_{ij}v_j \right),$$

\dots

$$\hat{A}_k = (\hat{k}_{ij}) = W^{-1}A_kW = \left(\frac{1}{w_i}k_{ij}w_j \right).$$

It is easy to show that $\hat{A}_1, \hat{A}_2, \dots, \hat{A}_k$ are all nonnegative irreducible matrices, and all the row sums of $\hat{A}_1, \hat{A}_2, \dots, \hat{A}_k$ are equal to $\rho(A_1), \rho(A_2), \dots, \rho(A_k)$, respectively.

Let $D = W \cdots VU$ be the product of k nonsingular diagonal matrices U, V, \dots, W . According to Lemma 2.4, we have

$$\begin{aligned} & D^{-1}(A_1 \circ A_2 \circ \cdots \circ A_k)D \\ &= (U^{-1}A_1U) \circ (V^{-1}A_2V) \circ \cdots \circ (W^{-1}A_kW) \\ &= \hat{A}_1 \circ \hat{A}_2 \circ \cdots \circ \hat{A}_k. \end{aligned}$$

Thus, we have $\rho(A_1 \circ A_2 \circ \cdots \circ A_k) = \rho(\hat{A}_1 \circ \hat{A}_2 \circ \cdots \circ \hat{A}_k)$. From Lemma 2.3, we have

$$\begin{aligned} & \rho(\hat{A}_1 \circ \hat{A}_2 \circ \cdots \circ \hat{A}_k) \\ &\leq \max_{i \neq j} \frac{1}{2} \left\{ \hat{a}_{ii}\hat{b}_{ii} \cdots \hat{k}_{ii} + \hat{a}_{jj}\hat{b}_{jj} \cdots \hat{k}_{jj} + \left[(\hat{a}_{ii}\hat{b}_{ii} \cdots \hat{k}_{ii} - \hat{a}_{jj}\hat{b}_{jj} \cdots \hat{k}_{jj})^2 \right. \right. \\ &\quad \left. \left. + 4 \left(\sum_{k \neq i} \hat{a}_{ik}\hat{b}_{ik} \cdots \hat{k}_{ik} \right) \left(\sum_{k \neq j} \hat{a}_{jk}\hat{b}_{jk} \cdots \hat{k}_{jk} \right) \right]^{1/2} \right\} \\ &\leq \max_{i \neq j} \frac{1}{2} \left\{ a_{ii}b_{ii} \cdots k_{ii} + a_{jj}b_{jj} \cdots k_{jj} + \left[(a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj})^2 \right. \right. \\ &\quad \left. \left. + 4 \left(\sum_{k \neq i} \frac{a_{ik}u_k}{u_i} \sum_{k \neq i} \frac{b_{ik}v_k}{v_i} \cdots \sum_{k \neq i} \frac{k_{ik}w_k}{w_i} \right) \left(\sum_{k \neq j} \frac{a_{jk}u_k}{u_j} \sum_{k \neq j} \frac{b_{jk}v_k}{v_j} \cdots \sum_{k \neq j} \frac{k_{jk}w_k}{w_j} \right) \right]^{1/2} \right\} \\ &= \max_{i \neq j} \frac{1}{2} \left\{ a_{ii}b_{ii} \cdots k_{ii} + a_{jj}b_{jj} \cdots k_{jj} + \left[(a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj})^2 \right. \right. \\ &\quad \left. \left. + 4(\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii})(\rho(A_1) - a_{jj}) \right. \right. \\ &\quad \left. \left. \times (\rho(A_2) - b_{jj}) \cdots (\rho(A_k) - k_{jj}) \right]^{1/2} \right\}. \end{aligned}$$

If $A_1 \circ A_2 \circ \cdots \circ A_k$ is reducible, the proof is similar to Theorem 2.1. So, the proof is completed. \square

Setting $k = 2$ in Theorem 2.2, we have the following corollary.

Corollary 2.2 ([2]) *Let $A_1, A_2 \in M_n$ and $A_1 \geq 0, A_2 \geq 0$. Then*

$$\begin{aligned} \rho(A_1 \circ A_2) &\leq \max_{i \neq j} \frac{1}{2} \left\{ a_{ii}b_{ii} + a_{jj}b_{jj} + \left[(a_{ii}b_{ii} - a_{jj}b_{jj})^2 \right. \right. \\ &\quad \left. \left. + 4(\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii})(\rho(A_1) - a_{jj})(\rho(A_2) - b_{jj}) \right]^{1/2} \right\}. \end{aligned}$$

We next give a simple comparison between the upper bound in (2.1) and the upper bound in (2.2). Without loss of generality, for $i \neq j$, assume that

$$\begin{aligned} & a_{ii}b_{ii} \cdots k_{ii} + (\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii}) \\ &\geq a_{jj}b_{jj} \cdots k_{jj} + (\rho(A_1) - a_{jj})(\rho(A_2) - b_{jj}) \cdots (\rho(A_k) - k_{jj}). \end{aligned}$$

Let $\gamma = a_{ii}b_{ii} \cdots k_{ii} + a_{jj}b_{jj} \cdots k_{jj}$. From (2.2), we have

$$\begin{aligned}
& a_{ii}b_{ii} \cdots k_{ii} + a_{jj}b_{jj} \cdots k_{jj} + [(a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj})^2 \\
& + 4(\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii})(\rho(A_1) - a_{jj}) \\
& \times (\rho(A_2) - b_{jj}) \cdots (\rho(A_k) - k_{jj})]^{\frac{1}{2}} \\
& \leq \gamma + \{(a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj})^2 + 4(\rho(A_1) - a_{ii}) \cdots (\rho(A_k) - k_{ii}) \\
& \times [(\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii}) + a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj}]\}^{\frac{1}{2}} \\
& = \gamma + [(a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj} + 2(\rho(A_1) - a_{ii}) \cdots (\rho(A_k) - k_{ii}))^2]^{\frac{1}{2}} \\
& = 2a_{ii}b_{ii} \cdots k_{ii} + 2(\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii}).
\end{aligned}$$

Thus, we have

$$\begin{aligned}
& \rho(A_1 \circ A_2 \circ \cdots \circ A_k) \\
& \leq \max_{i \neq j} \frac{1}{2} \{ a_{ii} \cdots k_{ii} + a_{jj} \cdots k_{jj} + [(a_{ii} \cdots k_{ii} - a_{jj} \cdots k_{jj})^2 + 4(\rho(A_1) - a_{ii}) \\
& \times (\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii})(\rho(A_1) - a_{jj})(\rho(A_2) - b_{jj}) \cdots (\rho(A_k) - k_{jj})]^{\frac{1}{2}} \} \\
& \leq \max_{1 \leq i \leq n} \frac{1}{2} [2a_{ii}b_{ii} \cdots k_{ii} + 2(\rho(A_1) - a_{ii}) \cdots (\rho(A_k) - k_{ii})] \\
& = \max_{1 \leq i \leq n} [a_{ii}b_{ii} \cdots k_{ii} + (\rho(A_1) - a_{ii})(\rho(A_2) - b_{ii}) \cdots (\rho(A_k) - k_{ii})].
\end{aligned}$$

Hence, bound (2.2) is better than bound (2.1).

In [8], the author proved that

$$\rho(A_1 \circ A_2 \circ \cdots \circ A_k) \leq \rho(A_1 A_2 \cdots A_k). \quad (2.3)$$

At present, we cannot give the comparison between bounds (2.1) and (2.3) or bounds (2.2) and (2.3), but the following numerical example shows that bounds (2.1) and (2.2) are better than (2.3). Next, we give an example: Consider four 4×4 nonnegative matrices

$$\begin{aligned}
A &= \begin{pmatrix} 4 & 1 & 0 & 2 \\ 0 & 0.05 & 1 & 1 \\ 0 & 0 & 4 & 0.5 \\ 1 & 0.5 & 0 & 4 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \\
C &= \begin{pmatrix} 2 & 0 & 1 & 1 \\ 1 & 4 & 0.5 & 0.5 \\ 1 & 0 & 3 & 0.5 \\ 0.5 & 1 & 1 & 2 \end{pmatrix}, \quad D = \begin{pmatrix} 2 & 0.5 & 0.5 & 0.5 \\ 1 & 1 & 1 & 1 \\ 0.5 & 0 & 2 & 0.5 \\ 0 & 1 & 1 & 2 \end{pmatrix}.
\end{aligned}$$

(i) It is easy to calculate that $\rho(A \circ B) = 5.4983$. By inequalities (2.1) and (2.2), we have

$$\rho(A \circ B) \leq \max_{1 \leq i \leq 4} \{a_{ii}b_{ii} + (\rho(A) - a_{ii})(\rho(B) - b_{ii})\} = 16.3949,$$

and

$$\rho(A \circ B) \leq 11.6478.$$

By inequality (2.3), we have

$$\rho(A \circ B) \leq \rho(AB) = 19.05.$$

(ii) From calculation, we get $\rho(A \circ B \circ C) = 12.0014$. By inequalities (2.1) and (2.2), we have

$$\rho(A \circ B \circ C) \leq \max_{1 \leq i \leq 4} \{a_{ii}b_{ii}c_{ii} + (\rho(A) - a_{ii})(\rho(B) - b_{ii})(\rho(C) - c_{ii})\} = 20.8846,$$

and

$$\rho(A \circ B \circ C) \leq 17.8268.$$

By inequality (2.3), we have

$$\rho(A \circ B \circ C) \leq \rho(ABC) = 88.5.$$

(iii) Let $A \circ B \circ C \circ D = G = (g_{ij})$. Then

$$G = \begin{pmatrix} 16 & 0 & 0 & 1 \\ 0 & 0.2 & 0.5 & 0.5 \\ 0 & 0 & 24 & 0.075 \\ 0 & 0.5 & 0 & 16 \end{pmatrix}, \quad ABCD = \begin{pmatrix} 117.25 & 78.75 & 155.75 & 126 \\ 34.3375 & 23.0625 & 45.6125 & 36.9 \\ 75.375 & 50.625 & 100.125 & 81 \\ 92.125 & 61.875 & 122.375 & 99 \end{pmatrix}.$$

It is easy to calculate that $\rho(G) = 24.0001$. By inequalities (2.1) and (2.2), we have

$$\rho(G) \leq \max_{1 \leq i \leq 4} \{a_{ii}b_{ii}c_{ii}d_{ii} + (\rho(A) - a_{ii})(\rho(B) - b_{ii})(\rho(C) - c_{ii})(\rho(D) - d_{ii})\} = 36.6608$$

and

$$\begin{aligned} \rho(G) &\leq \max_{i \neq j} \frac{1}{2} \{g_{ii} + g_{jj} + [(g_{ii} - g_{jj})^2 + 4(\rho(A) - a_{ii})(\rho(B) - b_{ii})(\rho(C) - c_{ii})(\rho(D) - d_{ii}) \\ &\quad \times (\rho(D) - d_{ii})(\rho(A) - a_{jj})(\rho(B) - b_{jj})(\rho(C) - c_{jj})(\rho(D) - d_{jj})]^{1/2}\} \\ &= 32.4451. \end{aligned}$$

By inequality (2.3), we have $\rho(G) \leq \rho(ABCD) = 339.44$.

Next, we will give some other inequalities for $\rho(A_1 \circ A_2 \circ \cdots \circ A_k)$. For $A_1 \geq 0$, write $L_1 = A_1 - \text{diag}(a_{11}, \dots, a_{nn})$. We denote $J_{A_1} = D_1^{-1}L_1$ with $D_1 = \text{diag}(d_{ii})$, where

$$d_{ii} = \begin{cases} a_{ii}, & \text{if } a_{ii} \neq 0, \\ 1, & \text{if } a_{ii} = 0. \end{cases}$$

Then J_{A_1} is nonnegative.

For $A_2 \geq 0$, let $D_2 = \text{diag}(s_{ii}), \dots$, for $A_k \geq 0$, let $D_k = \text{diag}(t_{ii})$ with

$$s_{ii} = \begin{cases} b_{ii}, & \text{if } b_{ii} \neq 0, \\ 1, & \text{if } b_{ii} = 0, \end{cases}$$

\dots ,

$$t_{ii} = \begin{cases} k_{ii}, & \text{if } k_{ii} \neq 0, \\ 1, & \text{if } k_{ii} = 0, \end{cases}$$

respectively. Then the nonnegative matrix J_{A_2}, \dots, J_{A_k} can be similarly defined.

Theorem 2.3 *Let $A_1, A_2, \dots, A_k \in M_n$ and $A_1 \geq 0, A_2 \geq 0, \dots, A_k \geq 0$. Then*

$$\begin{aligned} & \rho(A_1 \circ A_2 \circ \dots \circ A_k) \\ & \leq \max_{1 \leq i \leq n} \{a_{ii}b_{ii} \dots k_{ii} + d_{ii}\rho(J_{A_1})s_{ii}\rho(J_{A_2}) \dots t_{ii}\rho(J_{A_k})\}. \end{aligned} \quad (2.4)$$

Proof Let $Q = A_1 \circ A_2 \circ \dots \circ A_k$. First assume that Q is irreducible. Obviously A_1, A_2, \dots, A_k are all irreducible, and then $J_{A_1}, J_{A_2}, \dots, J_{A_k}$ are all nonnegative irreducible, so there exist k positive vectors x, y, \dots, h such that $J_{A_1}x = \rho(J_{A_1})x, J_{A_2}y = \rho(J_{A_2})y, \dots, J_{A_k}h = \rho(J_{A_k})h$. So, we have

$$\sum_{j \neq i} \frac{a_{ij}x_j}{x_i} = d_{ii}\rho(J_{A_1}), \quad \sum_{j \neq i} \frac{b_{ij}y_j}{y_i} = s_{ii}\rho(J_{A_2}), \quad \dots, \quad \sum_{j \neq i} \frac{k_{ij}h_j}{h_i} = t_{ii}\rho(J_{A_k}).$$

Now let $z = (z_i)$ be the vector, where $z_i = (x_i y_i \dots h_i) > 0$ for all i . For the irreducible nonnegative matrix Q , we have

$$\begin{aligned} (Qz)_i &= a_{ii}b_{ii} \dots k_{ii}z_i + \sum_{i \neq j} a_{ij}b_{ij} \dots k_{ij}z_j \\ &\leq a_{ii}b_{ii} \dots k_{ii}z_i + \left(\sum_{i \neq j} a_{ij}x_j \right) \left(\sum_{i \neq j} b_{ij}y_j \right) \dots \left(\sum_{i \neq j} k_{ij}h_j \right) \\ &= a_{ii}b_{ii} \dots k_{ii}z_i + (d_{ii}x_i\rho(J_{A_1}))(s_{ii}y_i\rho(J_{A_2})) \dots (t_{ii}h_i\rho(J_{A_k})) \\ &= \{a_{ii}b_{ii} \dots k_{ii} + d_{ii}\rho(J_{A_1})s_{ii}\rho(J_{A_2}) \dots t_{ii}\rho(J_{A_k})\}z_i. \end{aligned}$$

By Lemma 2.2, this shows that

$$\rho(A_1 \circ A_2 \circ \dots \circ A_k) \leq \max_{1 \leq i \leq n} \{a_{ii}b_{ii} \dots k_{ii} + d_{ii}\rho(J_{A_1})s_{ii}\rho(J_{A_2}) \dots t_{ii}\rho(J_{A_k})\}.$$

The proof is completed. \square

Setting $k = 2$ in Theorem 2.3, we have the following corollary.

Corollary 2.3 ([4]) *Let $A_1, A_2 \in M_n$ and $A_1 \geq 0, A_2 \geq 0$. Then*

$$\rho(A_1 \circ A_2) \leq \max_{1 \leq i \leq n} \{a_{ii}b_{ii} + d_{ii}\rho(J_{A_1})s_{ii}\rho(J_{A_2})\}.$$

Theorem 2.4 Let $A_1, A_2, \dots, A_k \in M_n$ and $A_1 \geq 0, A_2 \geq 0, \dots, A_k \geq 0$. Then

$$\begin{aligned} & \rho(A_1 \circ A_2 \circ \cdots \circ A_k) \\ & \leq \max_{i \neq j} \frac{1}{2} \left\{ a_{ii} b_{ii} \cdots k_{ii} + a_{jj} b_{jj} \cdots k_{jj} + [(a_{ii} b_{ii} \cdots k_{ii} - a_{jj} b_{jj} \cdots k_{jj})^2 \right. \\ & \quad \left. + 4(d_{ii} s_{ii} \cdots t_{ii})(d_{jj} s_{jj} \cdots t_{jj})(\rho^2(J_{A_1}) \rho^2(J_{A_2}) \cdots \rho^2(J_{A_k}))^{\frac{1}{2}}] \right\}. \end{aligned} \quad (2.5)$$

Proof First we assume that $A_1 \circ A_2 \circ \cdots \circ A_k$ is irreducible. Obviously, $J_{A_1}, J_{A_2}, \dots, J_{A_k}$ are all nonnegative irreducible, then there exist k positive vectors x, y, \dots, h such that $J_{A_1}x = \rho(J_{A_1})x, J_{A_2}y = \rho(J_{A_2})y, \dots, J_{A_k}h = \rho(J_{A_k})h$. Thus, we have

$$\sum_{j \neq i} \frac{a_{ij} x_j}{x_i} = d_{ii} \rho(J_{A_1}), \quad \sum_{j \neq i} \frac{b_{ij} y_j}{y_i} = s_{ii} \rho(J_{A_2}), \quad \dots, \quad \sum_{j \neq i} \frac{k_{ij} h_j}{h_i} = t_{ii} \rho(J_{A_k}).$$

Define

$$\begin{aligned} X &= \text{diag}(x_1, x_2, \dots, x_n), & Y &= \text{diag}(y_1, y_2, \dots, y_n), & \dots, \\ H &= \text{diag}(h_1, h_2, \dots, h_n). \end{aligned}$$

Let

$$\tilde{A}_1 = (\tilde{a}_{ij}) = X^{-1} A_1 X, \quad \tilde{A}_2 = (\tilde{b}_{ij}) = Y^{-1} A_2 Y, \quad \dots, \quad \tilde{A}_k = (\tilde{k}_{ij}) = H^{-1} A_k H.$$

From Lemma 2.4, we have

$$\begin{aligned} & (X^{-1} Y^{-1} \cdots H^{-1})(A_1 \circ A_2 \circ \cdots \circ A_k)(H \cdots Y X) \\ & = (X^{-1} A_1 X) \circ (Y^{-1} A_2 Y) \circ \cdots \circ (H^{-1} A_k H) \\ & = \tilde{A}_1 \circ \tilde{A}_2 \circ \cdots \circ \tilde{A}_k. \end{aligned}$$

Thus, $\rho(A_1 \circ A_2 \circ \cdots \circ A_k) = \rho(\tilde{A}_1 \circ \tilde{A}_2 \circ \cdots \circ \tilde{A}_k)$. From Lemma 2.3, we have

$$\begin{aligned} & \rho(\tilde{A}_1 \circ \tilde{A}_2 \circ \cdots \circ \tilde{A}_k) \\ & \leq \max_{i \neq j} \frac{1}{2} \left\{ a_{ii} b_{ii} \cdots k_{ii} + a_{jj} b_{jj} \cdots k_{jj} + [(a_{ii} b_{ii} \cdots k_{ii} - a_{jj} b_{jj} \cdots k_{jj})^2 \right. \\ & \quad \left. + 4 \left(\sum_{k \neq i} \frac{a_{ik} x_k}{x_i} \frac{b_{ik} y_k}{y_i} \cdots \frac{k_{ik} h_k}{h_i} \right) \left(\sum_{k \neq j} \frac{a_{jk} x_k}{x_j} \frac{b_{jk} y_k}{y_j} \cdots \frac{k_{jk} h_k}{h_j} \right) \right]^{\frac{1}{2}} \right\} \\ & \leq \max_{i \neq j} \frac{1}{2} \left\{ a_{ii} b_{ii} \cdots k_{ii} + a_{jj} b_{jj} \cdots k_{jj} + [(a_{ii} b_{ii} \cdots k_{ii} - a_{jj} b_{jj} \cdots k_{jj})^2 \right. \\ & \quad \left. + 4 \left(\sum_{k \neq i} \frac{a_{ik} x_k}{x_i} \sum_{k \neq i} \frac{b_{ik} y_k}{y_i} \cdots \sum_{k \neq i} \frac{k_{ik} h_k}{h_i} \right) \left(\sum_{k \neq j} \frac{a_{jk} x_k}{x_j} \sum_{k \neq j} \frac{b_{jk} y_k}{y_j} \cdots \sum_{k \neq j} \frac{k_{jk} h_k}{h_j} \right) \right]^{\frac{1}{2}} \right\} \\ & = \max_{i \neq j} \frac{1}{2} \left\{ a_{ii} b_{ii} \cdots k_{ii} + a_{jj} b_{jj} \cdots k_{jj} + [(a_{ii} b_{ii} \cdots k_{ii} - a_{jj} b_{jj} \cdots k_{jj})^2 \right. \\ & \quad \left. + 4(d_{ii} s_{ii} \cdots t_{ii})(d_{jj} s_{jj} \cdots t_{jj})(\rho^2(J_{A_1}) \rho^2(J_{A_2}) \cdots \rho^2(J_{A_k}))^{\frac{1}{2}}] \right\}. \end{aligned}$$

If $A_1 \circ A_2 \circ \cdots \circ A_k$ is reducible, then substituting $A_1 + tP, A_2 + tP, \dots, A_k + tP$ for A_1, A_2, \dots, A_k , respectively, in the previous case, letting $t \rightarrow 0$, the result is derived. \square

Setting $k = 2$ in Theorem 2.4, we have the following corollary.

Corollary 2.4 ([1]) *Let $A_1, A_2 \in M_n$ and $A_1 \geq 0, A_2 \geq 0$. Then*

$$\rho(A_1 \circ A_2) \leq \max_{i \neq j} \frac{1}{2} \left\{ a_{ii}b_{ii} + a_{jj}b_{jj} + [(a_{ii}b_{ii} - a_{jj}b_{jj})^2 + 4d_{ii}s_{ii}d_{jj}s_{jj}\rho^2(J_{A_1})\rho^2(J_{A_2})]^{1/2} \right\}.$$

We next give a comparison between the upper bound in (2.4) and the upper bound in (2.5). Without loss of generality, for $i \neq j$, assume that

$$\begin{aligned} & a_{ii}b_{ii} \cdots k_{ii} + d_{ii}s_{ii} \cdots t_{ii}\rho(J_{A_1})\rho(J_{A_2}) \cdots \rho(J_{A_k}) \\ & \geq a_{jj}b_{jj} \cdots k_{jj} + d_{jj}s_{jj} \cdots t_{jj}\rho(J_{A_1})\rho(J_{A_2}) \cdots \rho(J_{A_k}). \end{aligned}$$

Let $\gamma = a_{ii}b_{ii} \cdots k_{ii} + a_{jj}b_{jj} \cdots k_{jj}$. From (2.5), we have

$$\begin{aligned} & a_{ii}b_{ii} \cdots k_{ii} + a_{jj}b_{jj} \cdots k_{jj} + [(a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj})^2 \\ & \quad + 4(d_{ii}s_{ii} \cdots t_{ii})(d_{jj}s_{jj} \cdots t_{jj})(\rho^2(J_{A_1})\rho^2(J_{A_2}) \cdots \rho^2(J_{A_k}))]^{1/2} \\ & \leq \gamma + \{(a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj})^2 + 4d_{ii}s_{ii} \cdots t_{ii}\rho(J_{A_1})\rho(J_{A_2}) \cdots \rho(J_{A_k}) \\ & \quad \times [d_{ii}s_{ii} \cdots t_{ii}\rho(J_{A_1})\rho(J_{A_2}) \cdots \rho(J_{A_k}) + a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj}]\}^{1/2} \\ & = \gamma + [(a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj} + 2d_{ii}s_{ii} \cdots t_{ii}\rho(J_{A_1})\rho(J_{A_2}) \cdots \rho(J_{A_k}))^2]^{1/2} \\ & = 2a_{ii}b_{ii} \cdots k_{ii} + 2d_{ii}s_{ii} \cdots t_{ii}\rho(J_{A_1})\rho(J_{A_2}) \cdots \rho(J_{A_k}). \end{aligned}$$

Thus, from (2.5) and the above inequality, we have

$$\begin{aligned} & \rho(A_1 \circ A_2 \circ \cdots \circ A_k) \\ & \leq \max_{i \neq j} \frac{1}{2} \left\{ a_{ii}b_{ii} \cdots k_{ii} + a_{jj}b_{jj} \cdots k_{jj} + [(a_{ii}b_{ii} \cdots k_{ii} - a_{jj}b_{jj} \cdots k_{jj})^2 \right. \\ & \quad \left. + 4(d_{ii}s_{ii} \cdots t_{ii})(d_{jj}s_{jj} \cdots t_{jj})(\rho^2(J_{A_1})\rho^2(J_{A_2}) \cdots \rho^2(J_{A_k}))]^{1/2} \right\} \\ & \leq \max_{1 \leq i \leq n} \frac{1}{2} (2a_{ii}b_{ii} \cdots k_{ii} + 2d_{ii}s_{ii} \cdots t_{ii}\rho(J_{A_1})\rho(J_{A_2}) \cdots \rho(J_{A_k})) \\ & = \max_{1 \leq i \leq n} (a_{ii}b_{ii} \cdots k_{ii} + d_{ii}s_{ii} \cdots t_{ii}\rho(J_{A_1})\rho(J_{A_2}) \cdots \rho(J_{A_k})). \end{aligned}$$

Hence, the upper bound (2.5) is better than bound (2.4). Here too, we could not give the comparison between (2.4) and (2.3) or (2.5) and (2.3). Next, we give an example which shows that the results obtained in Theorems 2.3 and 2.4 are better than inequalities (2.3).

Let

$$A = \begin{pmatrix} 2 & 0 & 1 & 1 \\ 1 & 4 & 0.5 & 0.5 \\ 1 & 0 & 3 & 0.5 \\ 0.5 & 1 & 1 & 2 \end{pmatrix}, \quad B = \begin{pmatrix} 2 & 0.5 & 0.5 & 0.5 \\ 1 & 1 & 1 & 1 \\ 0.5 & 0 & 2 & 0.5 \\ 0 & 1 & 1 & 2 \end{pmatrix},$$

$$C = \begin{pmatrix} 4 & 1 & 1 & 1 \\ 2 & 2 & 1 & 1 \\ 0 & 2 & 2 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \quad D = \begin{pmatrix} 1 & 0.5 & 2 & 0.5 \\ 0.5 & 1 & 0.5 & 0 \\ 0 & 0.5 & 1 & 0.5 \\ 0 & 1 & 0.5 & 1 \end{pmatrix}.$$

Let $A \circ B \circ C \circ D = P = (p_{ij})$. Then

$$P = \begin{pmatrix} 16 & 0 & 1 & 0.25 \\ 1 & 8 & 0.25 & 0 \\ 0 & 0 & 12 & 0.075 \\ 0 & 1 & 0.5 & 4 \end{pmatrix}, \quad ABCD = \begin{pmatrix} 35.5 & 55.75 & 86 & 35.75 \\ 57.75 & 91.875 & 139 & 57.875 \\ 30.25 & 57.25 & 78 & 34.75 \\ 34.875 & 64.125 & 87.5 & 38.875 \end{pmatrix}.$$

It is easy to calculate that $\rho(P) = 16.0028$. By inequalities (2.4) and (2.5), we have

$$\rho(P) \leq \max_{1 \leq i \leq 4} \{p_{ii}(1 + \rho(J_A)\rho(J_B)\rho(J_C)\rho(J_D))\} = 36.2262$$

and

$$\begin{aligned} \rho(P) &\leq \max_{i \neq j} \frac{1}{2} \{p_{ii} + p_{jj} + [(p_{ii} - p_{jj})^2 + 4p_{ii}p_{jj}\rho^2(J_A)\rho^2(J_B)\rho^2(J_C)\rho^2(J_D)]^{\frac{1}{2}}\} \\ &= 29.6605. \end{aligned}$$

By inequality (2.3) and Lemma 2.1, we have $\rho(P) \leq 91.875$.

3 Conclusions

In this paper, we have proposed some upper bounds for $\rho(A_1 \circ A_2 \circ \cdots \circ A_k)$ of the Hadamard product of matrices. These bounds generalize some corresponding results of [1–4].

Acknowledgements

This research is financed by the Natural Science Foundation of Shandong Province ZR2017MA050; Natural Science Foundation of Zhejiang Province (LY14A010007) and Ningbo Natural Science Foundation (2015A610173).

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

Author details

¹College of Mathematics Science, Dezhou University, Dezhou, Shandong 253023, People's Republic of China.

²Department of Mathematics, Zhejiang Wanli University, Ningbo, Zhejiang 315100, People's Republic of China.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 1 October 2017 Accepted: 19 December 2017 Published online: 05 January 2018

References

1. Fang, MZ: Bounds on eigenvalues of the Hadamard product and the Fan product of matrices. *Linear Algebra Appl.* **425**, 7–15 (2007)
2. Liu, QB, Chen, GL: On two inequalities for the Hadamard product and the Fan product of matrices. *Linear Algebra Appl.* **431**, 974–984 (2009)
3. Liu, QB, Chen, GL, Zhao, LL: Some new bounds on the spectral radius of matrices. *Linear Algebra Appl.* **432**, 936–948 (2010)

4. Huang, R: Some inequalities for the Hadamard product and the Fan product of matrices. *Linear Algebra Appl.* **428**, 1551-1559 (2008)
5. Zhao, LL: Two inequalities for the Hadamard product of matrices. *J. Inequal. Appl.* **2012**, 122 (2012)
6. Audenaert, KMR: Spectral radius of Hadamard product versus conventional product for non-negative matrices. *Linear Algebra Appl.* **432**, 366-368 (2010)
7. Horn, RA, Zhang, FZ: Bounds on the spectral radius of a Hadamard product of nonnegative or positive semidefinite matrices. *Electron. J. Linear Algebra* **20**, 90-94 (2010)
8. Huang, ZJ: On the spectral radius and the spectral norm of Hadamard products of nonnegative matrices. *Linear Algebra Appl.* **434**, 457-462 (2011)
9. Zhou, DM, Chen, GL, Wu, GX, Zhang, XY: Some new bounds for eigenvalues of the Hadamard product and the Fan product of matrices. *Linear Algebra Appl.* **438**, 1415-1426 (2013)
10. Zhou, DM, Chen, GL, Wu, GX, Zhang, XY: Some inequalities for the Hadamard product of an *M*-matrix and an inverse *M*-matrix. *J. Inequal. Appl.* **2013**, 16 (2013)
11. Berman, A, Plemmons, RJ: *Nonnegative Matrices in the Mathematical Sciences*. SIAM, Philadelphia (1994)
12. Brauer, A: Limits for the characteristic roots of a matrix. *Duke Math. J.* **14**, 21-26 (1947)

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com